

References

- Ajchenbaum, F., Ando, K., DeCaprio, J. A., and Griffin, J. D. (1993). Independent regulation of human D-type cyclin gene expression during G1 phase in primary human T lymphocytes. *J Biol Chem* 268, 4113-9.
- Angel, P., and Karin, M. (1991). The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta* 1072, 129-57.
- Assoian, R. K. (1997). Anchorage-dependent cell cycle progression. *J Cell Biol* 136, 1-4.
- Assoian, R. K., and Zhu, X. (1997). Cell anchorage and the cytoskeleton as partners in growth factor dependent cell cycle progression. *Curr Opin Cell Biol* 9, 93-8.
- Bachmann, M. F., McKall-Faienza, K., Schmits, R., Bouchard, D., Beach, J., Speiser, D. E., Mak, T. W., and Ohashi, P. S. (1997). Distinct Roles for LFA-1 and CD28 during Activation of Naive T cells: Adhesion versus Costimulation. *Immunity* 7, 549-557.
- Bartek, J., Bartkova, J., and Lukas, J. (1996). The retinoblastoma protein pathway and the restriction point. *Curr Opin Cell Biol* 8, 805-14.
- Berg, N. N., and Ostergaard, H. L. (1995). Characterization of intercellular adhesion molecule-1 (ICAM-1)-augmented degranulation by cytotoxic T cells. ICAM-1 and anti-CD3 must be co-localized for optimal adhesion and stimulation. *J Immunol* 155, 1694-702.
- Bohmer, R. M., Scharf, E., and Assoian, R. K. (1996). Cytoskeletal integrity is required throughout the mitogen stimulation phase of the cell cycle and mediates the anchorage-dependent expression of cyclin D1. *Mol Biol Cell* 7, 101-111.
- Boussiotis, V. A., Barber, D. L., Lee, B. J., Gribben, J. G., Freeman, G. J., and Nadler, L. M. (1996). Differential association of protein tyrosine kinases with the T cell receptor is linked to the induction of anergy and its prevention by B7 family-mediated costimulation. *J Exp Med* 184, 365-76.

Cai, Z., Kishimoto, H., Brunmark, A., Jackson, M. R., Peterson, P. A., and Sprent, J. (1997). Requirements for peptide-induced T cell receptor downregulation on naive CD8+ T cells. *J Exp Med* 185, 641-51.

Cano, E., and Mahadevan, L. C. (1995). Parallel signal processing among mammalian MAPKs. *Trends Biochem Sci* 20, 117-22.

Chen, C. S., Mrksich, M., Huang, S., Whitesides, G. M., and Ingber, D. E. (1997). Geometric control of cell life and death. *Science* 276, 1425-8.

Chen, Q., Kinch, M. S., Lin, T. H., Burridge, K., and Juliano, R. L. (1994). Integrin-mediated cell adhesion activates mitogen-activated protein kinases. *J Biol Chem* 269, 26602-5.

Chen, Q., Lin, T. H., Der, C. J., and Juliano, R. L. (1996). Integrin-mediated activation of mitogen-activated protein (MAP) or extracellular signal-related kinase kinase (MEK) and kinase is independent of Ras. *J Biol Chem* 271, 18122-7.

Chen, Y. R., Meyer, C. F., and Tan, T. H. (1996). Persistent activation of c-Jun N-terminal kinase 1 (JNK1) in gamma radiation-induced apoptosis. *J Biol Chem* 271, 631-4.

Clark, E. A., and Brugge, J. S. (1995). Integrins and signal transduction pathways: the road taken. *Science* 268, 233-9.

Corbi, A. L. (1996). Leukocyte integrins: Structure, expression and function (Austin, Texas,U.S.A.: R.G. Landes company).

Corr, M., Slanetz, A. E., Boyd, L. F., Jelonek, M. T., Khilko, S., al-Ramadi, B. K., Kim, Y. S., Maher, S. E., Bothwell, A. L., and Margulies, D. H. (1994). T cell receptor-MHC class I peptide interactions: affinity, kinetics, and specificity. *Science* 265, 946-9.

Crawley, J. B., Rawlinson, L., Lali, F. V., Page, T. H., Saklatvala, J., and Foxwell, B. M. (1997). T cell proliferation in response to interleukins 2 and 7 requires p38MAP kinase activation. *J Biol Chem* 272, 15023-7.

Crawley, J. B., Willcocks, J., and Foxwell, B. M. (1996). Interleukin-7 induces T cell proliferation in the absence of Erk/MAP kinase activity. *Eur J Immunol* *26*, 2717-23.

Croft, M., and Dubey, C. (1997). Accessory molecule and costimulation requirements for CD4 T cell response. *Crit Rev Immunol* *17*, 89-118.

Damle, N. K., Klussman, K., Linsley, P. S., and Aruffo, A. (1992). Differential costimulatory effects of adhesion molecules B7, ICAM-1, LFA-3, and VCAM-1 on resting and antigen-primed CD4+ T lymphocytes. *J Immunol* *148*, 1985-92.

Daum, G., Eisenmann-Tappe, I., Fries, H. W., Troppmair, J., and Rapp, U. R. (1994). The ins and outs of Raf kinases. *Trends Biochem Sci* *19*, 474-80.

DeCaprio, J. A., Furukawa, Y., Ajchenbaum, F., Griffin, J. D., and Livingston, D. M. (1992). The retinoblastoma-susceptibility gene product becomes phosphorylated in multiple stages during cell cycle entry and progression. *Proc Natl Acad Sci U S A* *89*, 1795-8.

Dong, C., Yang, D. D., Wysk, M., Whitmarsh, A. J., Davis, R. J., and Flavell, R. A. (1998). Defective T cell differentiation in the absence of Jnk1. *Science* *282*, 2092-5.

Dubey, C., Croft, M., and Swain, S. L. (1995). Costimulatory requirements of naive CD4+ T cells. ICAM-1 or B7-1 can costimulate naive CD4 T cell activation but both are required for optimum response. *J Immunol* *155*, 45-57.

Dustin, M. L., and Springer, T. A. (1989). T-cell receptor cross-linking transiently stimulates adhesiveness through LFA-1. *Nature* *341*, 619-24.

Evans, G. A., Wahl, L. M., and Farrar, W. L. (1992). Interleukin-2-dependent phosphorylation of the retinoblastoma-susceptibility-gene product p110-115RB in human T-cells. *Biochem J* *282*, 759-64.

Fang, F., Orend, G., Watanabe, N., Hunter, T., and Ruoslahti, E. (1996). Dependence of cyclin E-CDK2 kinase activity on cell anchorage. *Science* 271, 499-502.

Fero, M. L., Rivkin, M., Tasch, M., Porter, P., Carow, C. E., Firpo, E., Polyak, K., Tsai, L. H., Broudy, V., Perlmutter, R. M., Kaushansky, K., and Roberts, J. M. (1996). A syndrome of multiorgan hyperplasia with features of gigantism, tumorigenesis, and female sterility in p27(Kip1)-deficient mice. *Cell* 85, 733-44.

Ferrell, J. E., Jr., and Bhatt, R. R. (1997). Mechanistic studies of the dual phosphorylation of mitogen-activated protein kinase. *J Biol Chem* 272, 19008-16.

Finco, T. S., and Baldwin, A. S. (1995). Mechanistic aspects of NF-kappa B regulation: the emerging role of phosphorylation and proteolysis. *Immunity* 3, 263-72.

Firpo, E. J., Koff, A., Solomon, M. J., and Roberts, J. M. (1994). Inactivation of a Cdk2 inhibitor during interleukin 2-induced proliferation of human T lymphocytes. *Mol Cell Biol* 14, 4889-901.

Garrity, P. A., Chen, D., Rothenberg, E. V., and Wold, B. J. (1994). Interleukin-2 transcription is regulated in vivo at the level of coordinated binding of both constitutive and regulated factors. *Mol Cell Biol* 14, 2159-69.

Ghosh, P., Sica, A., Cippitelli, M., Subleski, J., Lahesmaa, R., Young, H. A., and Rice, N. R. (1996). Activation of nuclear factor of activated T cells in a cyclosporin A-resistant pathway. *J Biol Chem* 271, 7700-4.

Ginsberg, M. H., Du, X., and Plow, E. F. (1992). Inside-out integrin signalling. *Curr Opin Cell Biol* 4, 766-71.

Gould, G. W., Cuenda, A., Thomson, F. J., and Cohen, P. (1995). The activation of distinct mitogen-activated protein kinase cascades is required for the stimulation of 2-deoxyglucose uptake by interleukin-1 and insulin-like growth factor-1 in KB cells. *Biochem J* 311, 735-8.

Green, J. M., Noel, P. J., Sperling, A. I., Walunas, T. L., Gray, G. S., Bluestone, J. A., and Thompson, C. B. (1994). Absence of B7-dependent responses in CD28-deficient mice. *Immunity* 1, 501-8.

Guadagno, T. M., Ohtsubo, M., Roberts, J. M., and Assoian, R. K. (1993). A link between cyclin A expression and adhesion-dependent cell cycle progression [published erratum appears in *Science* 1994 Jan 28;263(5146):455]. *Science* 262, 1572-5.

Hansen, L. K., Mooney, D. J., Vacanti, J. P., and Ingber, D. E. (1994). Integrin binding and cell spreading on extracellular matrix act at different points in the cell cycle to promote hepatocyte growth. *Mol Biol Cell* 5, 967-75.

Henics, T., Sanfridson, A., Hamilton, B. J., Nagy, E., and Rigby, W. F. (1994). Enhanced stability of interleukin-2 mRNA in MLA 144 cells. Possible role of cytoplasmic AU-rich sequence-binding proteins. *J Biol Chem* 269, 5377-83.

Hernandez-Caselles, T., Rubio, G., Campanero, M. R., del Pozo, M. A., Muro, M., Sanchez-Madrid, F., and Aparicio, P. (1993). ICAM-3, the third LFA-1 counterreceptor, is a co-stimulatory molecule for both resting and activated T lymphocytes. *Eur J Immunol* 23, 2799-806.

Hibi, M., Lin, A., Smeal, T., Minden, A., and Karin, M. (1993). Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain. *Genes Dev* 7, 2135-48.

Hodge, M. R., Ranger, A. M., Charles de la Brousse, F., Hoey, T., Grusby, M. J., and Glimcher, L. H. (1996). Hyperproliferation and dysregulation of IL-4 expression in NF-ATp-deficient mice. *Immunity* 4, 397-405.

Hughes, C. C., and Pober, J. S. (1996). Transcriptional regulation of the interleukin-2 gene in normal human peripheral blood T cells. Convergence of costimulatory signals and differences from transformed T cells. *J Biol Chem* 271, 5369-77.

Hulskens, J., Behrens, J., and Birchmeier, W. (1994). Tumor-suppressor gene products in cell contacts: the cadherin-APC-armadillo connection. *Curr Opin Cell Biol* 6, 711-6.

Ichijo, H., Nishida, E., Irie, K., ten Dijke, P., Saitoh, M., Moriguchi, T., Takagi, M., Matsumoto, K., Miyazono, K., and Gotoh, Y. (1997). Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* 275, 90-4.

Jain, J., Loh, C., and Rao, A. (1995). Transcriptional regulation of the IL-2 gene. *Curr Opin Immunol* 7, 333-42.

June, C. H., Fletcher, M. C., Ledbetter, J. A., Schieven, G. L., Siegel, J. N., Phillips, A. F., and Samelson, L. E. (1990). Inhibition of tyrosine phosphorylation prevents T-cell receptor-mediated signal transduction. *Proc Natl Acad Sci U S A* 87, 7722-6.

Kanner, S. B. (1996). Focal adhesion kinase-related fakB is regulated by the integrin LFA-1 and interacts with the SH3 domain of phospholipase C gamma 1. *Cell Immunol* 171, 164-9.

Kanner, S. B., Grosmaire, L. S., Ledbetter, J. A., and Damle, N. K. (1993). Beta 2-integrin LFA-1 signaling through phospholipase C-gamma 1 activation. *Proc Natl Acad Sci U S A* 90, 7099-103.

Karin, M. (1996). The regulation of AP-1 activity by mitogen-activated protein kinases. *Philos Trans R Soc Lond B Biol Sci* 351, 127-34.

Kishimoto, H., Cai, Z., Brunmark, A., Jackson, M. R., Peterson, P. A., and Sprent, J. (1996). Differing roles for B7 and intercellular adhesion molecule-1 in negative selection of thymocytes [see comments]. *J Exp Med* 184, 531-7.

Klippel, A., Reinhard, C., Kavanaugh, W. M., Apell, G., Escobedo, M. A., and Williams, L. T. (1996). Membrane localization of phosphatidylinositol 3-kinase is sufficient to activate multiple signal-transducing kinase pathways. *Mol Cell Biol* 16, 4117-27.

Kumagai, N., Benedict, S. H., Mills, G. B., and Gelfand, E. W. (1987). Requirements for the simultaneous presence of

phorbol esters and calcium ionophores in the expression of human T lymphocyte proliferation-related genes. *J Immunol* 139, 1393-9.

Kundig, T. M., Schorle, H., Bachmann, M. F., Hengartner, H., Zinkernagel, R. M., and Horak, I. (1993). Immune responses in interleukin-2-deficient mice. *Science* 262, 1059-61.

Lai, J. H., Horvath, G., Subleski, J., Bruder, J., Ghosh, P., and Tan, T. H. (1995). RelA is a potent transcriptional activator of the CD28 response element within the interleukin 2 promoter. *Mol Cell Biol* 15, 4260-71.

Lavoie, J. N., L'Allemand, G., Brunet, A., Muller, R., and Pouyssegur, J. (1996). Cyclin D1 expression is regulated positively by the p42/p44MAPK and negatively by the p38/HOGMAPK pathway. *J Biol Chem* 271, 20608-16.

Lenschow, D. J., Walunas, T. L., and Bluestone, J. A. (1996). CD28/B7 system of T cell costimulation. *Annu Rev Immunol* 14, 233-58.

Liu, J. (1993). FK506 and cyclosporin, molecular probes for studying intracellular signal transduction [published erratum appears in *Immunol Today* 1993 Aug;14(8):399]. *Immunol Today* 14, 290-5.

Lucas, J. J., Szepesi, A., Modiano, J. F., Domenico, J., and Gelfand, E. W. (1995). Regulation of synthesis and activity of the PLSTIRE protein (cyclin-dependent kinase 6 (cdk6)), a major cyclin D-associated cdk4 homologue in normal human T lymphocytes. *J Immunol* 154, 6275-84.

Maguire, J. E., Danahey, K. M., Burkly, L. C., and van Seventer, G. A. (1995). T cell receptor- and beta 1 integrin-mediated signals synergize to induce tyrosine phosphorylation of focal adhesion kinase (pp125FAK) in human T cells. *J Exp Med* 182, 2079-90.

Mainiero, F., Murgia, C., Wary, K. K., Curatola, A. M., Pepe, A., Blumemberg, M., Westwick, J. K., Der, C. J., and Giancotti, F. G. (1997). The coupling of alpha β 4 integrin to Ras-MAP

kinase pathways mediated by Shc controls keratinocyte proliferation. *Embo J* 16, 2365-75.

Marano, N., Holowka, D., and Baird, B. (1989). Bivalent binding of an anti-CD3 antibody to Jurkat cells induces association of the T cell receptor complex with the cytoskeleton. *J Immunol* 143, 931-8.

Marienfeld, R., Neumann, M., Chuvpilo, S., Escher, C., Kneitz, B., Avots, A., Schimpl, A., and Serfling, E. (1997). Cyclosporin A interferes with the inducible degradation of NF-kappa B inhibitors, but not with the processing of p105/NF-kappa B1 in T cells. *Eur J Immunol* 27, 1601-9.

Miyamoto, S., Teramoto, H., Coso, O. A., Gutkind, J. S., Burbelo, P. D., Akiyama, S. K., and Yamada, K. M. (1995). Integrin function: molecular hierarchies of cytoskeletal and signaling molecules. *J Cell Biol* 131, 791-805.

Miyamoto, S., Teramoto, H., Gutkind, J. S., and Yamada, K. M. (1996). Integrins can collaborate with growth factors for phosphorylation of receptor tyrosine kinases and MAP kinase activation: roles of integrin aggregation and occupancy of receptors. *J Cell Biol* 135, 1633-42.

Mochly-Rosen, D. (1995). Localization of protein kinases by anchoring proteins: a theme in signal transduction. *Science* 268, 247-51.

Modiano, J. F., Domenico, J., Szepesi, A., Lucas, J. J., and Gelfand, E. W. (1994). Differential requirements for interleukin-2 distinguish the expression and activity of the cyclin-dependent kinases Cdk4 and Cdk2 in human T cells. *J Biol Chem* 269, 32972-8.

Monks, C. R., Freiberg, B. A., Kupfer, H., Sciaky, N., and Kupfer, A. (1998). Three-dimensional segregation of supramolecular activation clusters in T cells. *Nature* 395, 82-6.

Morino, N., Mimura, T., Hamasaki, K., Tobe, K., Ueki, K., Kikuchi, K., Takehara, K., Kadokawa, T., Yazaki, Y., and Nojima, Y. (1995). Matrix/integrin interaction activates the mitogen-

activated protein kinase, p44 erk-1 and p42 erk-2. *J Biol Chem* 270, 269-73.

Nourse, J., Firpo, E., Flanagan, W. M., Coats, S., Polyak, K., Lee, M. H., Massague, J., Crabtree, G. R., and Roberts, J. M. (1994). Interleukin-2-mediated elimination of the p27Kip1 cyclin-dependent kinase inhibitor prevented by rapamycin. *Nature* 372, 570-3.

Nunes, J. A., Collette, Y., Truneh, A., Olive, D., and Cantrell, D. A. (1994). The role of p21ras in CD28 signal transduction: triggering of CD28 with antibodies, but not the ligand B7-1, activates p21ras. *J Exp Med* 180, 1067-76.

Pardi, R., Bender, J. R., Dettori, C., Giannazza, E., and Engleman, E. G. (1989). Heterogeneous distribution and transmembrane signaling properties of lymphocyte function-associated antigen (LFA-1) in human lymphocyte subsets. *J Immunol* 143, 3157-66.

Pardi, R., Inverardi, L., Rugarli, C., and Bender, J. R. (1992). Antigen-receptor complex stimulation triggers protein kinase C-dependent CD11a/CD18-cytoskeleton association in T lymphocytes. *J Cell Biol* 116, 1211-20.

Park, J. H., and Levitt, L. (1993). Overexpression of mitogen-activated protein kinase (ERK1) enhances T-cell cytokine gene expression: role of AP1, NF-AT, and NF-KB. *Blood* 82, 2470-7.

Pastor, M. I., Woodrow, M., and Cantrell, D. (1995). Regulation and function of p21ras in T lymphocytes. *Cancer Surv* 22, 75-83.

Perlmutter, R. M., Levin, S. D., Appleby, M. W., Anderson, S. J., and Alberola-Ila, J. (1993). Regulation of lymphocyte function by protein phosphorylation. *Annu Rev Immunol* 11, 451-99.

Petruzzelli, L., Takami, M., and Herrera, R. (1996). Adhesion through the interaction of lymphocyte function-associated antigen-1 with intracellular adhesion molecule-1 induces tyrosine phosphorylation of p130cas and its association with c-CrkII. *J Biol Chem* 271, 7796-801.

Price, M. A., Cruzalegui, F. H., and Treisman, R. (1996). The p38 and ERK MAP kinase pathways cooperate to activate Ternary Complex Factors and c-fos transcription in response to UV light. *Embo J* 15, 6552-63.

Ranger, A. M., Hodge, M. R., Gravallese, E. M., Oukka, M., Davidson, L., Alt, F. W., de la Brousse, F. C., Hoey, T., Grusby, M., and Glimcher, L. H. (1998). Delayed lymphoid repopulation with defects in IL-4-driven responses produced by inactivation of NF-ATc. *Immunity* 8, 125-34.

Renshaw, M. W., Ren, X.-D., and Schwartz, M. A. (1997). Growth factor activation of MAP kinase requires cell adhesion. *The EMBO Journal* 16, 5592-5599.

Rincon, M., and Flavell, R. A. (1994). AP-1 transcriptional activity requires both T-cell receptor-mediated and co-stimulatory signals in primary T lymphocytes. *Embo J* 13, 4370-81.

Rozdzial, M. M., Malissen, B., and Finkel, T. H. (1995). Tyrosine-phosphorylated T cell receptor zeta chain associates with the actin cytoskeleton upon activation of mature T lymphocytes. *Immunity* 3, 623-33.

Rudd, C. E. (1996). Upstream-downstream: CD28 cosignaling pathways and T cell function. *Immunity* 4, 527-34.

Rudd, C. E., Janssen, O., Cai, Y. C., da Silva, A. J., Raab, M., and Prasad, K. V. (1994). Two-step TCR zeta/CD3-CD4 and CD28 signaling in T cells: SH2/SW3 domains, protein-tyrosine and lipid kinases. *Immunol Today* 15, 225-34.

Ruoslahti, E., and Reed, J. C. (1994). Anchorage dependence, integrins, and apoptosis. *Cell* 77, 477-8.

Salio, M., Valitutti, S., and Lanzavecchia, A. (1997). Agonist-induced T cell receptor down-regulation: molecular requirements and dissociation from T cell activation. *Eur J Immunol* 27, 1769-73.

Sastry, S. K., and Horwitz, A. F. (1993). Integrin cytoplasmic domains: mediators of cytoskeletal linkages and extra- and

intracellular initiated transmembrane signaling. *Curr Opin Cell Biol* 5, 819-31.

Schaller, M. D., Otey, C. A., Hildebrand, J. D., and Parsons, J. T. (1995). Focal adhesion kinase and paxillin bind to peptides mimicking beta integrin cytoplasmic domains. *J Cell Biol* 130, 1181-7.

Schaller, M. D., and Parsons, J. T. (1994). Focal adhesion kinase and associated proteins. *Curr Opin Cell Biol* 6, 705-10.

Schlaepfer, D. D., Hanks, S. K., Hunter, T., and van der Geer, P. (1994). Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase. *Nature* 372, 786-91.

Schmits, R., Kundig, T. M., Baker, D. M., Shumaker, G., Simard, J. J., Duncan, G., Wakeham, A., Shahinian, A., van der Heiden, A., Bachmann, M. F., Ohashi, P. S., Mak, T. W., and Hickstein, D. D. (1996). LFA-1-deficient mice show normal CTL responses to virus but fail to reject immunogenic tumor. *J Exp Med* 183, 1415-26.

Semnani, R. T., Nutman, T. B., Hochman, P., Shaw, S., and van Seventer, G. A. (1994). Costimulation by purified intercellular adhesion molecule 1 and lymphocyte function-associated antigen 3 induces distinct proliferation, cytokine and cell surface antigen profiles in human "naive" and "memory" CD4+ T cells. *J Exp Med* 180, 2125-35.

Shier, P., Otulakowski, G., Ngo, K., Panakos, J., Chourmouzis, E., Christjansen, L., Lau, C. Y., and Fung-Leung, W. P. (1996). Impaired immune responses toward alloantigens and tumor cells but normal thymic selection in mice deficient in the beta2 integrin leukocyte function-associated antigen-1. *J Immunol* 157, 5375-86.

Sloan-Lancaster, J., Shaw, A. S., Rothbard, J. B., and Allen, P. M. (1994). Partial T cell signaling: altered phospho-zeta and lack of zap70 recruitment in APL-induced T cell anergy [see comments]. *Cell* 79, 913-22.

Stewart, M., Thiel, M., and Hogg, N. (1995). Leukocyte integrins. *Curr Opin Cell Biol* 7, 690-6.

Stewart, M. P., Cabanas, C., and Hogg, N. (1996). T cell adhesion to intercellular adhesion molecule-1 (ICAM-1) is controlled by cell spreading and the activation of integrin LFA-1. *J Immunol* 156, 1810-7.

Stokoe, D., Macdonald, S. G., Cadwallader, K., Symons, M., and Hancock, J. F. (1994). Activation of Raf as a result of recruitment to the plasma membrane. *Science* 264, 1463-7.

Su, B., Jacinto, E., Hibi, M., Kallunki, T., Karin, M., and Ben-Neriah, Y. (1994). JNK is involved in signal integration during costimulation of T lymphocytes. *Cell* 77, 727-36.

Szamel, M., Bartels, F., and Resch, K. (1993). Cyclosporin A inhibits T cell receptor-induced interleukin-2 synthesis of human T lymphocytes by selectively preventing a transmembrane signal transduction pathway leading to sustained activation of a protein kinase C isoenzyme, protein kinase C-beta. *Eur J Immunol* 23, 3072-81.

Tam, S. W., Shay, J. W., and Pagano, M. (1994). Differential expression and cell cycle regulation of the cdk4 inhibitor p16INK4. *Cancer Res* 54, 5816-20.

Timmerman, L. A., Clipstone, N. A., Ho, S. N., Northrop, J. P., and Crabtree, G. R. (1996). Rapid shuttling of NF-AT in discrimination of Ca²⁺ signals and immunosuppression. *Nature* 383, 837-40.

Treisman, R. (1996). Regulation of transcription by MAP kinase cascades. *Curr Opin Cell Biol* 8, 205-15.

Turcovski-Corrales, S. M., Fenton, R. G., Peltz, G., and Taub, D. D. (1995). CD28:B7 interactions promote T cell adhesion. *Eur J Immunol* 25, 3087-93.

Turner, J. M. (1993). IL-2-dependent induction of G1 cyclins in primary T cells is not blocked by rapamycin or cyclosporin A. *Int Immunol* 5, 1199-209.

Udagwa, T., Woodside, D. G., and McIntyre, B. W. (1996). Alpha 4 beta 1 (CD49d/CD29) integrin costimulation of human T cells enhances transcription factor and cytokine induction in the absence of altered sensitivity to anti-CD3 stimulation. *J Immunol* 157, 1965-72.

Ullman, K. S., Northrop, J. P., Verweij, C. L., and Crabtree, G. R. (1990). Transmission of signals from the T lymphocyte antigen receptor to the genes responsible for cell proliferation and immune function: the missing link. *Annu Rev Immunol* 8, 421-52.

Valitutti, S., Dessing, M., Aktories, K., Gallati, H., and Lanzavecchia, A. (1995). Sustained signaling leading to T cell activation results from prolonged T cell receptor occupancy. Role of T cell actin cytoskeleton. *J Exp Med* 181, 577-84.

Valitutti, S., Muller, S., Cella, M., Padovan, E., and Lanzavecchia, A. (1995). Serial triggering of many T-cell receptors by a few peptide-MHC complexes [see comments]. *Nature* 375, 148-51.

Valitutti, S., Muller, S., Salio, M., and Lanzavecchia, A. (1997). Degradation of T cell receptor (TCR)-CD3-zeta complexes after antigenic stimulation [see comments]. *J Exp Med* 185, 1859-64.

Van Gool, S. W., de Boer, M., and Ceuppens, J. L. (1993). CD28 ligation by monoclonal antibodies or B7/BB1 provides an accessory signal for the cyclosporin A-resistant generation of cytotoxic T cell activity. *J Immunol* 150, 3254-63.

Van Seventer, G. A., Bonvini, E., Yamada, H., Conti, A., Stringfellow, S., June, C. H., and Shaw, S. (1992). Costimulation of T cell receptor/CD3-mediated activation of resting human CD4+ T cells by leukocyte function-associated antigen-1 ligand intercellular cell adhesion molecule-1 involves prolonged inositol phospholipid hydrolysis and sustained increase of intracellular Ca²⁺ levels. *J Immunol* 149, 3872-80.

Viola, A., and Lanzavecchia, A. (1996). T cell activation determined by T cell receptor number and tunable thresholds [see comments]. *Science* 273, 104-6.

Ward, S. G., Parry, R. V., Matthews, J., and O'Neill, L. (1997). A p38 MAP kinase inhibitor SB203580 inhibits CD28-dependent T cell proliferation and IL-2 production. *Biochem Soc Trans* 25, 304S.

Weinberg, R. A. (1995). The retinoblastoma protein and cell cycle control. *Cell* 81, 323-30.

Whitmarsh, A. J., Shore, P., Sharrocks, A. D., and Davis, R. J. (1995). Integration of MAP kinase signal transduction pathways at the serum response element. *Science* 269, 403-7.

Xia, Z., Dickens, M., Raingeaud, J., Davis, R. J., and Greenberg, M. E. (1995). Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270, 1326-31.

Xing, J., Ginty, D. D., and Greenberg, M. E. (1996). Coupling of the RAS-MAPK pathway to gene activation by RSK2, a growth factor-regulated CREB kinase. *Science* 273, 959-63.

Zenner, G., Dirk zur Hausen, J., Burn, P., and Mustelin, T. (1995). Towards unraveling the complexity of T cell signal transduction. *Bioessays* 17, 967-75.

Zhu, X., and Assoian, R. K. (1995). Integrin-dependent activation of MAP kinase: a link to shape-dependent cell proliferation. *Mol Biol Cell* 6, 273-82.

Zhu, X., Ohtsubo, M., Bohmer, R. M., Roberts, J. M., and Assoian, R. K. (1996). Adhesion-dependent cell cycle progression linked to the expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of the retinoblastoma protein. *J Cell Biol* 133, 391-403.