

Literaturverzeichnis

- [Abr70] ABRAGAM, A.; BLEANEY, B.: *Electron paramagnetic resonance of transition ions.* Oxford : Clarendon Press, 1970
- [Adk80] ADKIN, R.J.; FOX, N.: *An introduction to the theory of elasticity.* New York : Longman, 1980, S. 56–122
- [Ahn86] AHN, C.B.; KIM, J.H.; CHO, Z.H.: High speed spiral scan echo planar imaging. In: *IEEE Trans. Med. Imaging* 5 (1986), S. 2–5
- [Aim92] AIME, S.; BOTTA, M.; ERMONDI, G.: NMR study of solution structures and dynamics of lanthanide(III) complexes of DOTA. In: *Inorg. Chem* 31 (1992), S. 4291–4299
- [Aim96] AIME, S.; BOTTA, M.; M., Fasano; TERRENO, E.; KINCESH, P.; CALABI, L.; PALEARI, L.: A new ytterbium chelate as contrast agent in chemical shift imaging and temperature sensitive probe for MR spectroscopy. In: *Magn. Reson. Med.* 35 (1996), S. 648–651
- [Ata93] ATALAY, M.K.; FORDER, J.R.; CHACKO, V.P.; KAWAMOTO, S.; ZERHOUNI, E.A.: Oxygenation in the rabbit myocardium: Assessment with susceptibility-dependent MR imaging. In: *Radiology* 189 (1993), S. 759–764
- [Atk91] ATKINSON, D.J.; EDELMAN, R.R.: Cineangiography of the heart in a single breath hold with a segmented TurboFLASH sequence. In: *Radiology* 178 (1991), S. 357–360
- [Axe89] AXEL, L.; DOUGHERTY, L.: MR imaging of motion with spatial modulation of magnetization. In: *Radiology* 171 (1989), S. 841–845
- [Axe92] AXEL, L.; GONCALVES, R.C.; BLOOMGARDEN, D.: Regional heart wall motion: Two-dimensional analysis and functional imaging with MR imaging. In: *Radiology* 183 (1992), S. 745–750
- [Bau97] BAUES, S.: *Vermessung und Optimierung der Homogenität eines 3-Tesla-Ganzkörper-NMR-Tomographen, Überprüfung der Linearität der Gradientenmagnetfelder.* Berlin, Technische Fachhochschule, Diplomarbeit, 1997

- [Baz94] BAZILLE, A.; GUTTMAN, M.A.; MCVEIGH, E.R.; ZERHOUNI, E.A.: Impact of semiautomated versus manual image segmentation errors on myocardial strain calculation by magnetic resonance tagging. In: *Invest. Radiol.* 29 (1994), S. 427–433
- [Bee92] DE BEER, R.; VAN ORMONDT, D.: *In-Vivo Magnetic Resonance Spectroscopy. Bd. 1: Analysis of NMR Data Using Time Domain Fitting.* Berlin : Springer-Verlag, 1992, S. 202–248
- [Ble72] BLEANEY, B.: Nuclear magnetic Resonance shifts in solution due to lanthanide ions. In: *J. Magn. Reson.* 8 (1972), S. 91–100
- [Blo46] BLOCH, F.; HANSEN, W.W.; PACKARD, M.E.: Nuclear induction. In: *Phys. Rev.* 69 (1946), S. 127
- [Bol90] BOLSTER, B.D. J.; MCVEIGH, E.R.; ZERHOUNI, E.A.: Myocardial tagging in polar coordinates with use of striped tags. In: *Radiology* 177 (1990), S. 769–772
- [Boo94a] VAN DEN BOOGAART, A.; ALA-KORPELA, M.; HOWE, F.A.; RODRIGUES, L.M.; STUBBS, M.; GRIFFITHS, J.R.: Magnetic resonance spectroscopy data analysis: Time or frequency domain? In: *MAGMA* 2 (1994), S. 479–482
- [Boo94b] VAN DEN BOOGAART, A.; ALA-KORPELA, M.; JOKISAARI, J.; GRIFFITHS, J.R.: Time and frequency domain analysis of NMR data compared: An application to 1D ^1H spectra of lipoproteins. In: *Magn. Reson. Med.* 31 (1994), S. 347–358
- [Boo96] VAN DEN BOOGART, A.; VAN HUFFEL, S.; GRAVERON-DEMILLY, D.; VAN ORMONDT, D.; DE BEER, R.: MRUI: A graphical user interface for accurate routine MRS data analysis. In: *Book of abstracts: 13th Annual Meeting of ESMRMB* ESMRMB, 1996. – Siehe auch: URL <http://www.mrui.uab.es/mrui/mruiHomePage.html>, S. 318
- [Bot94] BOTTOMLEY, P.A.: MR spectroscopy of the human heart: the status and the challenges. In: *Radiology* 191 (1994), S. 593–612
- [Bot97] BOTTOMLEY, P.A.; LUGO OLIVIERI, C.H.; GIAQUINTO, R.: What is the optimum phased array coil design for cardiac and torso magnetic resonance? In: *Magn. Reson. Med.* 37 (1997), S. 591–599
- [Bru99] Bruker Medical GmbH: *ParaVision MANUAL Version 2.0.* 1999
- [Bry84] BRYANT, D.J.; PAYNE, J.A.; FIRMIN, D.N.; LONGMORE, D.B.: Measurement of flow with NMR imaging using a gradient pulse and phase difference technique. In: *J. Comput. Assist. Tomogr.* 8 (1984), S. 588–593
- [Bue95] BUECHLER, D.N.; ROPER, D.H.; DURNEY, C.H.; CHRISTENSEN, D.A.: Modelling source in the FDTD formulation and their use in quantifying sources and boundary condition errors. In: *IEEE Trans. Microwave Theory Tech.* 43 (1995), S. 810–814

- [Bur97] BURAI, L.; HIETAPELTO, V.; KIRÁLY, R.; TÓTH, É.; BÜCHER, E.: Stability constants and ^1H relaxation effects of ternary complexes formed between Gd-DTPA, Gd-DTPA-BMA, Gd-DOTA, and Gd-EDTA and citrate, phosphate, and carbonate ions. In: *Magn. Reson. Med.* 38 (1997), S. 146–150
- [Bur99] BURGESS, R.E.; YU, Y.; ABDULJALIL, A.M.; KANGARLU, A.; ROBITAILLE, P.-M. L.: High signal-to-noise FLASH imaging at 8 Tesla. In: *Magn. Reson. Imaging* 17 (1999), S. 1099–1103
- [Cad95] CADY, E.B.; D'SOUZA, P.C.; PENRICE, J.; LOREK, A.: The estimation of local brain temperature by in vivo ^1H magnetic resonance spectroscopy. In: *Magn. Reson. Med.* 33 (1995), S. 862–867
- [Cet84] CETAS, T.C.: Will thermometric tomography become practical for hyperthermia treatment monitoring? In: *Cancer Res. (suppl)* 44 (1984), S. 4805–4808
- [Con95] CONSTANTINIDIS, C.D.; WESTGATE, C.R.; O'DELL, W.G.; ZERHOUNI, E.A.; MCVEIGH, E.R.: A phased array coil for human cardiac imaging. In: *Magn. Reson. Med.* 34 (1995), S. 92–98
- [Cor95] CORBETT, R.J.; LAPTOOK, A.R.; TOLLEFSBOL, G.; KIM, B.: Validation of a noninvasive method to measure brain temperature in vivo using ^1H NMR spectroscopy. In: *J. Neurochem.* 64 (1995), S. 1224–1230
- [Cra88] CRAWLEY, A.P.; WOOD, M.L.; HENKELMAN, R.M.: Elimination of transverse coherence in FLASH MRI. In: *Magn. Reson. Med.* 8 (1988), S. 248–260
- [Dam71] DAMADIAN, R.: Tumor detection by nuclear magnetic resonance. In: *Science* 171 (1971), S. 1151–1153
- [Dei99] DEICHMANN, R.; HAHN, D.; HAASE, A.: Fast T_1 mapping on a whole-body scanner. In: *Magn. Reson. Med.* 42 (1999), S. 206–209
- [Del91] DELANNOY, J.; CHEN, C.-N.; TURNER, R.; LEVIN, R.L.; LE BIHAN, D.: Noninvasive temperature imaging using diffusion MRI. In: *Magn. Reson. Med.* 19 (1991), S. 333–339
- [Dic50] DICKINSON, W.C.: Dependence of the ^{19}F nuclear resonance position on chemical compound. In: *Phys. Rev.* 77 (1950), S. 736
- [Don94] DONG, S.J.; MACGREGOR, J.H.; CRAWLEY, A.P.; MCVEIGH, E.; BELENKIE, I.; SMITH, E.R.; TYBERG, J.V.; BEYAR, R.: Left ventricular wall thickness and regional systolic function in patients with hypertrophic cardiomyopathy. A three-dimensional tagged magnetic resonance imaging study. In: *Circulation* 90 (1994), S. 1200–1209
- [Don95] DONG, S.J.; CRAWLEY, A.P.; MACGREGOR, J.H.; PETRANK, Y.F.; BERGMAN, D.W.; BELENKIE, I.; SMITH, E.R.; TYBERG, J.V.; BEYAR, R.: Regional left ventricular systolic function in relation to the cavity geometry in patients with chronic

- right ventricular pressure overload. A three-dimensional tagged magnetic resonance imaging study. **In:** *Circulation* 91 (1995), S. 2359–2370
- [DP95a] DE POORTER, J.: Noninvasive MRI thermometry with the proton resonance frequency method: Study of susceptibility effects. **In:** *Magn. Reson. Med.* 34 (1995), S. 359–367
- [DP95b] DE POORTER, J.; DE WAGTER, C.; DE DEENE, Y.; THOMSON, C.; STÅHLBERG, F.; ACHTEN, E.: Noninvasive MRI thermometry with the proton resonance frequency (PRF) method: *In vivo* results in human muscle. **In:** *Magn. Reson. Med.* 33 (1995), S. 74–81
- [Dra74] DRAGO, R.S.; ZINK, J.I.; RICHMAN, R.M.; PERRY, W.D.: Theory of isotropic shifts in the NMR of paramagnetic materials: Part I. **In:** *J. Chem. Educ.* 51 (1974), S. 371–376
- [Ern66] ERNST, R.R.; ANDERSON, W.A.: Application of Fourier transform spectroscopy to magnetic resonance. **In:** *Rev. Sci. Instrum.* 37 (1966), S. 93–102
- [Fay95] FAYAD, Z.A.; CONNICK, T.J.; AXEL, L.: An improved quadrature or phased-array coil for MR cardiac imaging. **In:** *Magn. Reson. Med.* 34 (1995), S. 186–193
- [Fay98] FAYAD, Z.A.; FERRARI, V.A.; KRAITCHMAN, D.L.; YOUNG, A.A.; PALEVSKY, H.I.; BLOOMGARDEN, D.C.; AXEL, L.: Right ventricular regional function using MR tagging: Normals versus chronic pulmonary hypertension. **In:** *Magn. Reson. Med.* 39 (1998), S. 116–123
- [Fis93] FISCHER, S.E.; MCKINNON, G.C.; MAIER, S.E.; BOESIGER, P.: Improved myocardial tagging contrast. **In:** *Magn. Reson. Med.* 30 (1993), S. 191–200
- [Fis94] FISCHER, S.E.; MCKINNON, G.C.; SCHEIDECKER, M.B.; PRINS, W.; MEIER, D.; BOESIGER, P.: True myocardial motion tracking. **In:** *Magn. Reson. Med.* 31 (1994), S. 401–413
- [Fre96] FRENZEL, T.; ROTH, K.; KOSSLER, S.; RADÜCHEL, B.; BAUER, H.; PLATZEK, J.; WEINMANN, H.J.: Noninvasive temperature measurement *in vivo* using a temperature-sensitive lanthanide complex and ^1H magnetic resonance spectroscopy. **In:** *Magn. Reson. Med.* 35 (1996), S. 364–369
- [Haa91] HAACKE, E.M.; LINDSKOG, E.D.; LIN, W.: A fast, iterative, partial-fourier technique capable of local phase recovery. **In:** *J. Magn. Reson.* 92 (1991), S. 126–145
- [Han86] HAND, J.W. (Hrsg.); JAMES, J.R. (Hrsg.): *Physical Techniques in Clinical Hyperthermia*. Hertfordshire : Research Studies Press Ltd, 1986
- [Hen85] HENKELMAN, R.M.: Measurement of signal intensities in the presence of noise in MR images. **In:** *Med. Phys.* 12 (1985), S. 232–233

- [Hen93] HENNEL, J.W.; KLINOWSKI, J.: *Fundamentals of Nuclear Magnetic Resonance*. 1. Aufl. Harlow : Longman Scientific & Technical, 1993
- [Hen95] HENNEL, F.; NÉDÉLEC, J.-F.: Interleaved asymmetric echo-planar imaging. In: *Magn. Reson. Med.* 34 (1995), S. 520–524
- [Hen99] HENTSCHEL, M.; DREHER, W.; WUST, P.; RÖLL, S.; LEIBFRITZ, D.; FELIX, R.: Fast spectroscopic imaging for non-invasiv thermometry using the Pr[MOE-DO3A] complex. In: *Phys. Med. Biol.* 44 (1999), S. 2397–2408
- [Hin66] HINDMAN, J.C.: Proton resonance shift of water in the gas and liquid states. In: *J. Chem. Phys.* 44 (1966), S. 4582–4592
- [Hoe93] HOEFT, S.; ROTH, K.: Struktur und Dynamik von Lanthanoid-Tetraazacyclododecantetraacetat-(DOTA-)Komplexen in Lösung. In: *Chem. Ber.* 126 (1993), S. 869–873
- [IEC95] Norm IEC 60601-2-33 1995. Particular requirements for the safety of magnetic resonance equipment for medical diagnostics
- [Il'98] IL'YASOV, K.A.; HENNIG, J.: Single-shot diffusion-weighted RARE sequence: Application for temperature monitoring during hyperthermia session. In: *J. Magn. Reson. Imaging* 8 (1998), S. 1296–1305
- [Ing75] INGELS, N.B.; DAUGHTERS, G.T.; STINSON, E.B.; ALDERMAN, E.L.: Measurement of midwall myocardial dynamics in intact man by radiography of surgically implanted markers. In: *Circulation* 52 (1975), S. 859–867
- [Jac91] JACKSON, J.I.; MEYER, C.H.; NISHIMURA, D.G.; MACOVSKI, A.: Selection of a convolution function for Fourier inversion using gridding. In: *IEEE Trans. Med. Imaging* 10 (1991), S. 473–478
- [Jaf96] JAFFER, F.A.; WEN, H.; BALABAN, R.S.; WOLFF, S.D.: A method to improve the B_0 homogeneity of the heart in vivo. In: *Magn. Reson. Med.* 36 (1996), S. 375–383
- [Jak95] JAKOB, P.M.; ZIEGLER, A.; DORAN, S.J.; DÉCORPS, M.: Echo-time-encoded burst imaging (EBI): aAnovel technique for spectroscopic imaging. In: *Magn. Reson. Med.* 33 (1995), S. 573–578
- [Jol91] JOLIOT, M.; MAZOYER, B.M.; HUESMAN, R.H.: In vivo NMR spectral parameter estimation: A comparison between time and frequency domain methods. In: *Magn. Reson. Med.* 18 (1991), S. 358–370
- [Ker97] KERR, A.B.; PAULY, J.M.; HU, B.S.; LI, K.C.; HARDY, C.J.; MEYER, C.H.; MACOVSKI, A.; NISHIMURA, D.G.: Real-time interactive MRI on a conventional scanner. In: *Magn. Reson. Med.* 38 (1997), S. 355–367
- [Kie98] VON KIENLIN, M.: Cardiac ^1H -MR spectroscopy. In: *MAGMA* 6 (1998), S. 107–108

- [Kni92] KNIJN, A.; DE BEER, R.; VAN ORMONDT, D.: Frequency-selective quantification in the time domain. In: *J. Magn. Reson.* 97 (1992), S. 444–450
- [Knu86] KNUTTEL, B.; JURETSCHKE, H.P.: Temperature measurements by nuclear magnetic resonance and its possible use as a means of in vivo noninvasive temperature measurement and for hyperthermia treatment assessment. In: *Recent Results Cancer Res.* 101 (1986), S. 109–118
- [Kon97] KONSTANCZAK, P.; WUST, P.; SANDER, B.; SCHRÜNDER, S.; FRENZEL, T.; WŁODARCZYK, W.; VOGL, T.; MÜLLER, G.; FELIX, R.: Thermometrie durch Messung der chemischen Verschiebung eines Lanthanidenkomplexes. In: *Strahlenther. Onkol.* 173 (1997), S. 106–116
- [Kum94] KUMAR, S.; GOLDGOF, D.: Automatic tracking of SPAMM grid and estimation of deformation parameters from cardiac MR images. In: *IEEE Trans. Med. Imaging* 13 (1994), S. 122–132
- [Kur94] KURODA, K.; ABE, K.; TSUTSUMI, S.; ISHIHARA, Y.; SUZUKI, Y.; SATOH, K.: Water proton magnetic resonance spectroscopic imaging. In: *Biomed. Thermol.* 13 (1994), S. 43–62
- [Kur96] KURODA, K.; SUZUKI, Y.; ISHIHARA, Y.; OKAMOTO, K.; SUZUKI, Y.: Temperature mapping using water proton chemical shift obtained with 3D-MRSI: Feasibility in vivo. In: *Magn. Reson. Med.* 35 (1996), S. 20–29
- [Kur00] KURODA, K.; MULKERN, R.V.; OSHIO, K.; PANYCH, L.P.; NAKAI, T.; MORIYA, T.; OKUDA, S.; HYNNEN, K.; JOLES, F.A.: Temperature mapping using the water proton chemical shift: Self-referenced method with echo-planar spectroscopic imaging. In: *Magn. Reson. Med.* 43 (2000), S. 220–225
- [Lau73] LAUTERBUR, P.C.: Image formation by induced local interactions: examples employing nuclear magnetic resonance. In: *Nature* 242 (1973), S. 190–191
- [LB89] LE BIHAN, D.; DELANOY, J.; LEVIN, R.L.: Temperature mapping with MR imaging of molecular diffusion: Application of hyperthermia. In: *Radiology* 171 (1989), S. 853–857
- [Leu99] LEUSSLER, C.; RÖSCHMANN, P.: 3D simulation of electromagnetic fields inside the human body for application of MR and EPR: Effects of object size and frequency on RF field homogeneity. In: *Proceedings of the 7th Annual Meeting of ISMRM* ISMRM, 1999, S. 2050
- [Lju83] LJUNGGREN, S.: A simple graphical representation of Fourier-based imaging methods. In: *J. Magn. Reson.* 54 (1983), S. 338–343
- [Lof98] LOFFLER, R.; SAUTER, R.; KOLEM, H.; HAASE, A.; VON KIENLIN, M.: Localized spectroscopy from anatomically matched compartments: improved sensitivity and

- localization for cardiac ^{31}P MRS in humans. In: *J. Magn. Reson.* 134 (1998), S. 287–299
- [Lor99] LORENZ, C.H.; JOHANSSON, L.O.M.: Contrast-enhanced coronary MRA. In: *J. Magn. Reson. Imaging* 10 (1999), S. 703–708
- [Mac95] MACFALL, J.; PRESCOTT, D.M.; FULLAR, E.; SAMULSKI, T.V.: Temperature dependence of canine brain tissue diffusion coefficient measured *in vivo* with magnetic resonance echo-planar imaging. In: *Int. J. Hyperthermia* 11 (1995), S. 73–86
- [Mai92] MAIER, S.E.; FISCHER, S.E.; MCKINNON, G.C.; HESS, O.M.; KRAYENBUEHL, H.P.; BOESIGER, P.: Evaluation of left ventricular segmental wall motion in hypertrophic cardiomyopathy with myocardial tagging. In: *Circulation* 86 (1992), S. 1919–1928
- [Man77] MANSFIELD, P.: Multi-planar image formation using NMR spin echoes. In: *J. Phys. C* 10 (1977), S. L55–L58
- [McK93] MCKINNON, G.C.: Ultrafast interleaved gradient-echo-planar imaging on a standard scanner. In: *Magn. Reson. Med.* 30 (1993), S. 609–616
- [Mey86] MEYERS, J.H.; STIRLING, M.C.; CHOY, M.; BUDA, A.J.; GALLAGHER, K.P.: Direct measurement of inner and outer wall thickening dynamics with epicardial echocardiography. In: *Circulation* 74 (1986), S. 164–172
- [Mey90] MEYER, C.H.; PAULY, J.M.; MACOVSKI, A.; NISHIMURA, D.G.: Simultaneous spatial and spectral selective excitation. In: *Magn. Reson. Med.* 15 (1990), S. 287–304
- [Mor95] MORNEBURG, H. (Hrsg.): *Bildgebende Systeme für die medizinische Diagnostik*. 3. Aufl. München : Publicis-MCD-Verlag, 1995, S. 152–190, 501–621
- [Mou96] MOULTON, M.J.; CRESWELL, L.L.; DOWNING, S.W.; ACTIS, R.L.; SZABO, B.A.; VANNIER, M.W.; PASQUE, M.K.: Spline surface interpolation for calculating 3-D ventricular strains from MRI tissue tagging. In: *Am. J. Physiol.* 270 (1996), S. H281–H297
- [Nag98] NAGEL, E.; UNDERWOOD, R.; PENNELL, D.; SECHTEM, U.P.; NEUBAUER, S.; WICKLINE, S.; HESS, O.M.; SCHWAIGER, M.; FLECK, E.: Neue Entwicklungen bei der nicht-invasiven Herzdarstellung. In: *Dtsch. Med. Wschr.* 123 (1998), S. 1018–1021
- [Nag00] NAGEL, E.; STUBER, M.; BURKHARD, B.; FISCHER, S.E.; SCHEIDECKER, M.B.; BOESIGER, P.: Cardiac rotation and relaxation in patients with aortic valve stenosis. In: *Eur. Heart J.* 21 (2000), S. 582–589
- [Nie96] NIEMI, P.; PONCELET, B.P.; KWONG, K.K.; WEISSKOFF, R.M.; ROSEN, B.R.; BRADY, T.J.; KANTOR, H.L.: Myocardial intensity changes associated with flow

- simulation in blood oxygenation sensitive magnetic resonance imaging. In: *Magn. Reson. Med.* 36 (1996), S. 78–82
- [Noe00] NOESKE, R.; SEIFERT, F.; RHEIN, K.-H.; RINNEBERG, H.: Human cardiac imaging at 3 T using phased array coils. In: *Magn. Reson. Med.* (2000). – im Druck
- [O'D95] O'DELL, W.G.; MOORE, C.C.; HUNTER, W.C.; ZERHOUNI, E.A.; MCVEIGH, E.R.: Three-dimensional myocardial deformations: Calculation with displacement field fitting to tagged MR images. In: *Radiology* 195 (1995), S. 829–835
- [Par83] PARKER, D.L.; SMITH, V.; SHELDON, P.; CROOKS, L.E.; FUSSELL, L.: Temperature distribution measurements in two-dimensional NMR imaging. In: *Med. Phys.* 10 (1983), S. 321–325
- [Pat95] Patentschrift DE 43 18 369 C 1 1995. Verwendung von makrocyclischen Metallkomplexen als Temperatursonden
- [Pau89] PAULY, J.; NISHIMURA, D.; MACOVSKI, A.: A k-space analysis of small-tip-angle excitation. In: *J. Magn. Reson.* 81 (1989), S. 43–56
- [Pel99] PELLER, M.; LÖFFLER, R.; BAUR, A.; TURNER, P.; ABDEL-RAHMAN, S.; FUTSCHIK, G.; SANTL, M.; HIDDEMANN, W.; REISER, M.; ISSELS, R.: MRT-gesteuerte regionale Tiefenhyperthermie. In: *Der Radiologe* 9 (1999), S. 756–763
- [Pet00] PETERS, R.D.; HENKELMAN, R.M.: Proton-resonance frequency shift MR thermometry is affected by changes in the electrical conductivity of tissue. In: *Magn. Reson. Med.* 43 (2000), S. 62–71
- [Pij92] PIJNAPPEL, W.W.F.; VAN DEN BOOGART, A.; DE BEER, R.; VAN ORMONDT, D.: SVD-based quantification of magnetic resonance signals. In: *J. Magn. Reson.* 97 (1992), S. 122–134
- [Poh97] POHMANN, R.; VON KIENLIN, M.; HAASE, A.: Theoretical evaluation and comparison of fast chemical shift imaging methods. In: *J. Magn. Reson.* 129 (1997), S. 145–160
- [Pos94] POSSE, S.; DECARLI, C.; LE BIHAN, D.: Three-dimensional echo-planar MR spectroscopic imaging at short echo times in the human brain. In: *Radiology* 192 (1994), S. 733–738
- [Pro50] PROCTER, W.G.; YU, F.C.: The dependence of a nuclear magnetic resonance frequency upon chemical compound. In: *Phys. Rev.* 77 (1950), S. 717
- [Pro93] PROVENCHER, S.W.: Estimation of metabolite concentration from localized in vivo proton NMR spectra. In: *Magn. Reson. Med.* 30 (1993), S. 672–679
- [Pur46] PURCELL, E.M.; TORREY, H.C.; POUND, R.V.: Resonance absorption by nuclear magnetic moments in a solid. In: *Phys. Rev.* 69 (1946), S. 37–38

- [Rau98] RAU, B.; WUST, P.; HOHENBERGER, P.; LÖFFEL, J.; HÜNERBEIN, M.; BELOW, C.; GELLERMANN, J.; SPEIDEL, A.; VOGL, T.; RIESS, H.; FELIX, R.; SCHLAG, P.M.: Preoperative hyperthermia combined with radiochemotherapy in locally advanced rectal cancer: a phase II clinical trial. In: *Ann. Surg.* 227 (1998), S. 380–389
- [Ree93] REEDER, S.B.: *Development of High Speed, High Resolution Magnetic Resonance Imaging and Tagging Techniques*. Baltimore, The Johns Hopkins University, Master Thesis, 1993
- [Ree98] REEDER, S.B.; FARANESH, A.Z.; BOXERMAN, J.L.; MCVEIGH, E.R.: In vivo measurement of T_2^* and field inhomogeneity maps in the human heart at 1.5 T. In: *Magn. Reson. Med.* 39 (1998), S. 988–998
- [Rin97] RINCK, P.A.: Current status & perspectives of biomedical MR. In: *Book of abstracts: 14th Annual Meeting of ESMRMB* ESMRMB, 1997, S. 6–7
- [Sch99] SCHMIDT, M.; THEISSEN, P.; CRNAC, J.; JOCHIMS, M.; VOTH, E.; BAER, F.M.; SCHICHA, H.; ERDMANN, E.: Klinischer Nutzen der Magnetresonanztomographie zur Darstellung von Koronarstenosen. In: *Dtsch. Med. Wschr.* 124 (1999), S. 1294–1300
- [Sin97] SINGERMAN, R.W.; DENISON, T.J.; WEN, H.; BALABAN, R.S.: Simulation of B_1 field distribution and intrinsic signal-to-noise in cardiac MRI as a function of static magnetic field. In: *J. Magn. Reson.* 125 (1997), S. 72–83
- [Stu99] STUBER, M.; FISCHER, S.E.; SCHEIDECKER, M.B.; BOESIGER, P.: Toward high-resolution myocardial tagging. In: *Magn. Reson. Med.* 41 (1999), S. 639–643
- [Sul90] SULLIVAN, D.M.: Three-dimensional computer simulation in deep regional hyperthermia using the FDTD method. In: *IEEE Trans. Microwave Theory Tech.* 38 (1990), S. 204–211
- [Tak95] TAKAHASHI, A.; PETERS, T.: Compensation of multi-dimensional selective excitation pulses using measured k -space trajectories. In: *Magn. Reson. Med.* 34 (1995), S. 446–456
- [Tan95] TANG, C.; MCVEIGH, E.R.; ZERHOUNI, E.A.: Multi-shot EPI for improvement of myocardial tag contrast: Comparison with segmented SPGR. In: *Magn. Reson. Med.* 33 (1995), S. 443–447
- [Twi83] TWIEG, D.B.: The k -space trajectory formulation of the NMR imaging process with application in the analysis and synthesis of imaging methods. In: *Med. Phys.* 10 (1983), S. 610–621
- [Van97] VANHAMME, L.; VAN DEN BOOGART, S.: Improved method for accurate and efficient quantification of MRS data with use of prior knowledge. In: *J. Magn. Reson.* 129 (1997), S. 35–43

- [Var00] Varian, Inc: *Produktinformation*. 2000. – Siehe auch:
URL <http://www.varianinc.com/nmr/900>
- [Vee88] VAN DER VEEN, J.W.; DE BEER, R.; LUYTEN, P.R.; ORMONDT, D.: Accurate quantification of *in vivo* ^{31}P NMR signals using the variable projection method and prior knowledge. In: *Magn. Reson. Med.* 6 (1988), S. 92–98
- [Wan90] WANG, H.Z.; RIEDERER, S.J.: A spoiling sequence for suppression of residual transverse magnetization. In: *Magn. Reson. Med.* 15 (1990), S. 175–191
- [Wan95] WANG, J.; REYKOWSKI, A.; DICKAS, J.: Calculation of signal-to-noise ratio for simple surface coils and arrays of coils. In: *IEEE Trans. Biomed. Eng.* 42 (1995), S. 908–917
- [Web93] WEBB, A.G.; WONG, M.; WILMES, L.J.; KOLBECK, K.J.; MAGIN, R.L.; SUSLICK, K.S.: Cobalt-59 functional agents for localized *in-vivo* temperature measurements. In: *Proceedings of the 12th Annual Meeting of SMRM* SMRM, 1993, S. 245
- [Wen97] WEN, H.; DENISON, T.J.; SINGERMAN, R.W.; BALABAN, R.S.: The intrinsic signal-to-noise ratio in human cardiac imaging at 1.5, 3 and 4 T. In: *J. Magn. Reson.* 125 (1997), S. 65–71
- [Wlo98] WŁODARCZYK, W.; BOROSCHEWSKI, R.; HENTSCHEL, M.; WUST, P.; MÖNICH, G.; FELIX, R.: Three-dimensional monitoring of small temperature changes for therapeutic hyperthermia using MR. In: *J. Magn. Reson. Imaging* 8 (1998), S. 165–174
- [Wus98] WUST, P.; GELLERMANN, J.; HARDER, C.; TILLY, W.; B., Rau.; DINGES, S.; SCHLAG, P.; BUDACH, V.; FELIX, R.: Rationale for using invasive thermometry for regional hyperthermia of pelvic tumors. In: *Int. J. Radiat. Oncol. Biol. Phys.* 41 (1998), S. 1129–1137
- [You93] YOUNG, A.A.; AXEL, L.; DOUGHERTY, L.; BOGEN, D.K.; PARENTEAU, C.S.: Validation of tagging with MR imaging to estimate material deformation. In: *Radiology* 188 (1993), S. 101–108
- [You94] YOUNG, I.R.; HAND, J.W.; OATRIDGE, A.; PRIOR, M.V.; FORSE, G.R.: Further observations on the measurement of tissue T_1 to monitor temperature *in vivo* by MRI. In: *Magn. Reson. Med.* 31 (1994), S. 342–345
- [You95] YOUNG, A.A.; KRAITCHMAN, D.L.; DOUGHERTY, L.; AXEL, L.: Tracking and finite element analysis of stripe deformation in magnetic resonance tagging. In: *IEEE Trans. Med. Imaging* 14 (1995), S. 413–421
- [Yun91] YUN, K.L.; MICZYPORUK, M.A.; DAUGHTERS, G.T.; INGELS, N.B.; STINSON, E.B.; ALDERMAN, E.L.; HANSEN, D.E.; MILLER, D.C.: Alterations in left ventricular diastolic twist mechanics during acute human cardiac allograft rejection. In: *Circulation* 83 (1991), S. 962–973

- [Zer88] ZERHOUNI, E.A.; PARISH, D.M.; ROGERS, W.J.; SHAPIRO, E.P.: Human heart: Tagging with MR imaging – a method for noninvasive assessment of myocardial motion. In: *Radiology* 169 (1988), S. 59–63
- [Zha92] ZHANG, Y.; SAMULSKI, T.V.; JOINES, W.T.; MATTIELLO, J.; LEVIN, R.L.; LE BIHAN, D.: On the accuracy of noninvasive thermometry using molecular diffusion magnetic resonance imaging. In: *Int. J. Hyperthermia* 8 (1992), S. 263–274
- [Zuo96] ZUO, C.S.; BOWERS, J.L.; METZ, K.R.; NOSAKA, T.; SHERRY, A.D.; CLOUSE, M.E.: TmDOTA⁵⁻: A substance for NMR temperature measurements *in vivo*. In: *Magn. Reson. Med.* 36 (1996), S. 955–959
- [Zur91] ZUR, Y.; WOOD, M.L.; NEURINGER, L.J.: Spoiling of transverse magnetization in steady-state sequences. In: *Magn. Reson. Med.* 21 (1991), S. 251–263

