

IV. Diskussion	68
V. Schlussfolgerung.....	74
VI. Zusammenfassung	76
VII. Summary.....	78
VIII. Literaturverzeichnis	80
IX. Anhang.....	122

VIII. Literaturverzeichnis

AGNEESSENS, J.; ENGELEN, S.; DEBEVER, P.; VERCROYSSE, J. (1998):

Gasterophilus intestinalis infections in horses in Belgium.

Vet. Parasitol., 77, 199-204

AHMED, Z.G.; HOSSNY, G.A.; EL-SEIFY, M.A.; HASSLINGER, M.-A. (1994):

Tissue reactions to the Dictyocaulus arnfieldi in Egyptian donkeys.

Beni Suef vet. med. Res., 4, 75-85

ANDERSON, R.R. (1984):

The use of ivermectin in horses: Research and clinical observations.

Comp. Cont. Educ. Pract. Vet., 6, 516-520

ANONYM (1990):

Doramectin (UK-67, 994) -Injectable Solution

Pfizer Central Research, Sandwich, Kent, England, Herstellerinformation 27. November 1990

ARBUCKLE, J.B.R.; KHALIL, L.F. (1978):

Thelazia lacrymalis in the eyelids of British horses.

Vet. Rec., 103, 158-159

AREF, S. (1982):

A random walk model for the migration of Strongylus vulgaris in the intestinal arteries of the horse.

Cornell. Vet., 72, 64-75

ARENA, J.P.; LIU, K.K.; PARESS, P.S.; FRAZIER, E.G.; CULLY, D.F.; MROZIK, H.;

SCHAEFFER, J.M. (1995):

The mechanism of action of avermectin in Caenorhabditis elegans: Correlation between activation of glutamate-sensitive chlorid current, membrane binding and biological activity.

J. Parasitol., 81, 286-294

ARUNDEL, J.H. (1978):

Parasitic disease of the horse.

Veterinary Review No. 18, University of Sydney Post Graduate Foundation in Veterinary Science, Sydney, New South Wales, 83

AUSTIN, S.M.; DI PIETRO, J.A.; FOREMAN, J.H.; BAKER, G.J.; TODD, K.S. (1991):

Comparison of the efficacy of ivermectin, oxibendazole and pyrantel pamoate against 28-day Parascaris equorum larvae in the intestine of pony foals.

J. Amer. vet. med. Ass., 198, 1946-1949

BAIN, A.M.; ROFE, J.C.; HOTSON, I.K.; MURPHY, S. (1969):

Habronema megastoma larvae associated with pulmonary abscesses in a foal.

Aust. vet. J., 45, 101-102

BAIN, S.A.; KELLY, J.D. (1977):

Prevalence and pathogenicity of Anoplocephala perfoliata in a horse population in South Auckland.

N. Z. vet. J., 25, 27-28

BARCLAY, W.P.; PHILLIPS, T.N.; FOERNER, J.J. (1982):

Intussusception associated with Anoplocephala perfoliata infection in five horses.

J. Amer. vet. med. Ass., 180, 752-753

BARNES, E.H.; DOBSON, R.J.; BARGER, I.A. (1995):

Worm control and anthelmintic resistance: Adventures with a model.

Parasitol. Today, 11, 56-63

BARRAGRY, T. (1984):

Anthelmintics - A review. Part 1.

N. Z. vet. J., 32, 161-164

BARTH, D.; HEINZE-MUTZ, E.M.; RONCALLI, R.A.; SCHLUTER, D.; GROSS, S.J. (1993):

The degradation of dung produced by cattle treated with an ivermectin slow-release bolus.

Vet. Parasitol., 48, 215-228

BAUER, C. (1986a):

Ciliates as possible food source for adult Oxyuris equi Schrank 1788.

Z. Parasitenk., 72, 279-280

BAUER, C. (1986b):

Befall mit Magenparasiten in Norddeutschland.

Dtsch. tierärztl. Wschr., 93, 388-389

BAUER, C. (1986c):

Filariosen.

In: Handlexikon der tierärztlichen Praxis, I-IV, Medical Book Company, Bjaeverskov / DK,

Lief. 182, 260g-260m

BAUER, C.; GANDRAS, R.; STOYE, M.; BÜRGER, H.-J. (1983):

Eine Feldstudie zur Anthelminthika-Resistenz von Strongyliden beim Pferd.

Berl. Münch. tierärztl. Wschr., 96, 312-316

BAUER, C.; STOYE, M. (1984):

Ergebnisse parasitologischer Kotuntersuchungen von Equiden, Hunden, Katzen und Igeln der Jahre 1974 - 1983.

Dtsch. tierärztl. Wschr., 91, 255-258

- BAUER, C.; MERKT, J.C.; JANKE-GRIMM, G.; BÜRGER, H.-J. (1986):
Prevalence and control of benzimidazole-resistant small strongyles on German thoroughbred studs.
Vet. Parasitol., 21, 189-203
- BAUER, C.; CIRAK, V.Y.; HERMOSILLA, C.; OKORO, H. (1998):
Efficacy of a 2 % moxidectin gel against gastrointestinal parasites of ponies.
Vet. Rec., 143, 558-561
- BELLO, T.R. (1989):
*Efficacy of ivermectin against experimental and natural infections of *Gasterophilus* spp. in ponies.*
Amer. J. vet. Res., 50, 2120-2123
- BELLO, T.R.; LANINGHAM, J.E.T. (1992):
Evaluation of daily pyrantel tartrate treatment of horses for prevention of intestinal penetration and development of small strongyle larvae resulting from experimentally induced infections.
J. Equine vet. Sci., 12, 280-286
- BELLO, T.R.; LANINGHAM, J.E.T. (1994):
A controlled trial evaluation of three oral dosages of moxidectin against equine parasites.
J. Equine vet. Sci., 14, 483-488
- BEMRICK, W.J. (1978):
Tolerance of equine strongylid larvae to desiccation and freezing.
Cryobiology, 15, 214-218
- BENNETT, D.G. (1972):
Predisposition to abdominal crisis in the horse.
J. Amer. vet. med. Ass., 161, 1189-1194
- BENNETT, D.G. (1986):
Clinical pharmacology of ivermectin.
J. Amer. vet. med. Ass., 189, 100-104
- BEROZA, G.A.; BARCLAY, W.P.; PHILLIPS, T.N.; DONAWICK, W.J.; FOERNER, J.J. (1983):
*Caecal perforation and peritonitis associated with *Anoplocephala perfoliata* infection in three horses.*
J. Amer. Ass. Equine Pract., 32, 435-439
- BEROZA, G.A.; WILLIAMS, R.; MARCUS, L.C.; MILLE, P. (1985):
Prevalence of tapeworm infection and associated large bowel disease in horses.
Proc. 2nd Equ. Colic. Symp., 2, 21-25

BEROZA, G.A.; MARCUS, L.C.; WILLIAMS, R.; BAUER, S.M. (1987):

Laboratory diagnosis of Anoplocephala perfoliata infection in horses.

Proc. Amer. Ass. Equine Pract., 32, 435-439

BIRD, J.; HERD, R.P. (1995):

In vitro assessment of two species of nematophagous fungi (Arthrobotrys oligospora and Arthrobotrys flagrans) to control the development of infective cyathostome larvae from naturally infected horses.

Vet. Parasitol., 56, 181-187

BITTNER, G. (1983):

Ein Beitrag zur Epizootologie der Strongylideninfektion des Pferdes in einem bayerischen Gestütsareal.

Vet. med. Diss., München

BOCH, J.; SUPPERER, R. (1992):

Veterinärmedizinische Parasitologie.

4. Aufl. Paul Parey Verlag, Berlin und Hamburg

BOERSEMA, J.H. (1982):

Possibilities and limitations in the detection of anthelmintic resistance.

In: Facts and reflections IV. C.E.C. Workshop „Resistance of parasites to anthelmintics“.

(9.-10.12., Lelystad/ Niederlande)

BOERSEMA, J.H.; EYSKER, M.; VAN DER AAR, W.M. (1998):

The reappearance of strongyle eggs in the faeces of horses after treatment with moxidectin.

Vet. Quarterly, 20, 15-17

BRAUER; E.; BRAUER, U. (1997):

Klinischer Fall einer generalisierten Demodikose bei einem Pferd und deren Behandlung mit Doramectin.

Tierärztl. Umschau, 52, 131-133

BRITT, D.P.; PRESTON, J.M. (1985):

Efficacy of ivermectin against Dictyocaulus arnfieldi in ponies.

Vet. Rec., 116, 343-345

BROCARD, P.; PFISTER, K. (1993):

Efficacité de l'antiparasitaire oral Mebendazole plus Trichlorfon (Telmin® plus Trichlorfon) contre les gasterophiles chez le cheval.

Schweiz. Arch. Tierheilk., 135, 356-359

BRUCE, A.H.; VAN DEN HEUVEL, W.J.H.; WISLOCKI, P.G. (1993):

Environmental effects of the usage of avermectins in livestock.

Vet. Parasitol., 48, 109-125

BUCKNELL, D.G.; GASSER, R.B.; BEVERIDGE, I. (1995):

The prevalence and epidemiology of gastrointestinal parasites of horses in Victoria, Australia.

Int. J. Parasitol., 25, 711-724

BUENO, L.; RUCKEBUSCH, Y.; DORCHIES, P. (1979):

Disturbances of digestive motility in horses associated with strongyle infection.

Vet. Parasitol., 5, 253-260

BÜRGER, H.J. (1986):

Auftreten benzimidazolresistenter kleiner Strongyliden beim Pferd.

Prakt. Tierarzt, 67, 643-646

BURKHARDT, E. (1978):

Zur Pathologie der Komplikationen bei der Strongylose.

6. Arbeitstagung der Fachgruppe Pferdekrankheiten der DVG, (28.-30.10., Berlin/BRD),
77-81

BURKHARDT, E. (1983):

*Zur Pathologie des *Strongylus (Delafondia) vulgaris*-Befalls beim Pferd - eine Übersicht.*

Berl. Münch. tierärztl. Wschr., 96, 37-43

CAIRNS, G.C.; HOLMDEN, J.H. (1977):

The anthelmintic efficacy of cambendazole in horses.

N. Z. vet. J., 25, 35-37

CARTER, G.T.; NIETSCH, J.A.; BORDERS, D.B. (1987):

*Structure determination of LL-F28249-alpha, beta, gamma and lambda, potent antiparasitic macrolides from *Streptomyces cyaneoegriseus* spp. noncyanogenus.*

J. Chem. Soc. Chem. Commun., 6, 402-404

CHIEJINA, S.N.; MASON, S.A. (1977):

*Immature stages of *Trichonema* spp. as a cause of diarrhoea in adult horses in spring.*

Vet. Rec., 100, 360-361

CHIU, S.-H.L.; LU, A.Y.H. (1989):

Metabolism and tissue residues.

In: W.C. Campbell (ed.), *Ivermectin and Abamectin*.

Springer Verlag, New York, 131-143

CHRISTL, H. (1971):

*Darmruptur nach Massenbefall mit *Anoplocephala perfoliata* bei einer Stute.*

Berl. Münch. tierärztl. Wschr., 84, 305-307

CLAYTON, H.M. (1986):

Askarids. Recent advances.

Vet. Clin. North Amer. Equine Pract., 2, 313-328

CLAYTON, H.M.; DUNCAN, J.L. (1978):

Clinical signs associated with Parascaris equorum infection in wormfree foals and yearlings.

Vet. Parasitol., 4, 69-78

CLAYTON, H.M.; DUNCAN, J.L. (1979):

The migration and development of Parascaris equorum in the horse.

Int. J. Parasitol., 9, 285-292

CLAYTON, H.M.; NEAVE, R.M.S. (1979):

*Efficacy of mebendazole against *D. arnfieldi* in the donkey.*

Vet. Rec., 104, 571-572

CLAYTON, H.M.; DUNCAN, J.L.; DARGIE, J.D. (1980):

Pathophysiological changes associated with Parascaris equorum infection in the foal.

Equine vet. J., 12, 23-25

COGLEY, T.P. (1989):

*Effects of migrating *Gasterophilus intestinalis* larvae (Diptera: Gasterophoridae) on the mouth of the horse.*

Vet. Parasitol., 31, 317-331

COGLEY, T.P.; ANDERSON, J.R.; COGLEY, L.J. (1982):

*Migration of *Gasterophilus intestinalis* larvae (Diptera: Gasterophoridae) in the equine oral cavity.*

Int. J. Parasitol., 12, 473-480

COLES, G.C.; BAUER, C; BORGSTEEDE, F.H.M.; GEERTS, S.; KLEI, T.R.; TAYLOR, M.A.;

WALLER, P.J. (1992):

World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) methods for the detection of anthelmintic resistance in nematodes of veterinary importance.

Vet. Parasitol., 44, 35-44

COLES, G.C.; HILLYER, M.H.; TAYLOR, F.G.R.; PARKER, L.D. (1998):

Activity of moxidectin against bots and lungworm in equids.

Vet. Rec., 143, 169-170

COLGAZIER, M.L.; ENZIE, F.D.; KATES, K.C. (1977):

Critical anthelmintic trials in ponies with four benzimidazoles: mebendazole, cambendazole, fenbendazole and albendazole.

J. Parasitol., 63, 724-727

CONDER, G.A.; THOMPSON, D.P.; JOHNSON, S.S. (1993):

Demonstration of co-resistance of Haemonchus contortus to ivermectin and moxidectin.

Vet. Rec., 132, 651-652

CONDER, G.A.; CAMPBELL, W.C. (1995):

Chemotherapy of nematode infections of veterinary importance, with special reference to drug resistance.

Adv. Parasitol., 35, 1-84

CONNAN, R.M. (1985):

Askaridoses of domesticated animals.

In: GAFAAR, S.M.; HOWARD, E.W.; MARSH, R.E. (eds.), Parasites, Pets and Predators, 253-269

COOK, R.W. (1975):

Cambendazole paste as an anthelmintic in horses.

Vet. Med. / Small Anim. Clin., 70, 1451-1453

CORBA, J.; PRASLICKA, J.; VARADY, M.; ANDRASKO, H.; HOLAKOVSKY, P. (1995):

Efficacy of Moxidectin 2 % Equine Gel and Eqvalan 1 % paste against naturally acquired internal parasite infections in horses.

Helminthologia, 32, 215-218

CORLOUER, J.P. (1980):

Strongylus vulgaris, données bibliographiques récentes. Diagnostic de l'infestation larvaire par la méthode d'immunofluorescence indirecte.

Thesis, École Nationale Vétérinaire, Toulouse, France

COSGROVE, J.S.; SHEERAN, J.J.; SAINTY, T.J. (1986):

Intussusception associated with infection with Anoplocephala perfoliata in a two-year-old thoroughbred.

Irish vet. J., 40, 35-36

COSTA, A.J.; BARBOSA, O.F.; MORAES, F.R.; ACUNA, A.H.; ROCHA, U.F.; SOARES, V.E.; PAULLILO, A.C.; SANCHES, A. (1998):

Comparative efficacy evaluation of moxidectin gel and ivermectin paste against internal parasites of equines in Brazil.

Vet. Parasitol., 80, 29-36

CRAIG, T.M.; HATCHFIELD, T.A.; PANKAVICH, J.A.; WANG, G.T. (1992):

Efficacy of moxidectin against an ivermectin-resistant strain of Haemonchus contortus in sheep.

Vet. Parasitol., 41, 329-333

CRAIG, T.M.; SCRUTCHFIELD, W.L.; MARTIN, M.T. (1993):

Comparison of prophylactic pyrantel and suppressive ivermectin anthelmintic programs in young horses.

Equine Pract., 15, 24-29

CRAVEN, J.; BJØRN, H.; HENDRICKSEN, S.A.; NANSEN, P.; LARSEN, M.; LENDAL, S.

(1996):

A survey of anthelmintic resistance on Danish horse farms.

Parasitologia, 38, 232

CZOSNEK, T. (1988):

Gasterophilus intestinalis, the cause of death in a mare.

Medycyna Weterynaryjna, 44, 6, 346

DAHME, E.; WEISS, E. (1988):

Grundriß der speziellen pathologischen Anatomie der Haustiere.

4. Aufl., Ferdinand Enke Verlag, Stuttgart

DAURIO, C.P.; LEANING, W.H.D. (1989):

The effect of oral ivermectin on immature ascarids in foals.

J. Equine vet. Sci., 9, 312-315

DEEGEN, E.; FREYTAG, K.; HERTSCH, B. (1976):

Diagnose der Schlundkrankungen beim Pferd.

Tierärztl. Prax., 4, 207-221

DEMEULENAERE, D.; DORNY, P.; VERCROYSSE, J.; CLAEREBOUT, E. (1996):

Comparative studies of ivermectin and moxidectin in the control of naturally acquired cyathostome infections in horses.

Abstr. Joint Meeting of Parasitology (13.-15.5., Bruxelles / Belgium), 40

DEMEULENAERE, D.; VERCROYSSE, J.; DORNY, P.; CLAEREBOUT, E. (1997):

Comparative studies of ivermectin and moxidectin in the control of naturally acquired cyathostome infections in horses.

Vet. Rec., 141, 383-386

DERKMANN, K.; HASSLINGER, M.-A. (1977):

Zur Bekämpfung des Pferdespulwurmes, Parascaris equorum (Goeze, 1782), mit Panacur®.

Berl. Münch. tierärztl. Wschr., 90, 95-98

DEWES, H.F.; TOWNSEND, K.G. (1990):

Further observations on Strongyloides westeri dermatitis: Recovery of larvae from soil and bedding, an survival in treated sites.

N. Z. vet. J., 38, 34-37

DI PIETRO, J.A.; TODD, K.S. (1989):

Anthelmintics used in treatments of parasitic infections of horses.

Equine Pract., 11, 5-15

DI PIETRO, J.A.; LOCK, T.F.; TODD, K.S.; DAVIS, J. (1989):

Efficacy of ivermectin in the treatment of induced Parascaris equorum infection in pony foals.

J. Amer. med. Ass., 195, 1712-1714

DI PIETRO, J.A.; KLEI, T.R.; FRENCH, D.D. (1990):

Contemporary topics in equine parasitology.

Comp. Cont. Educ. Pract. Vet., 12, 713-716

DI PIETRO, J.A.; EWERT; K.E.; TODD, K.S. (1993a):

Ivermectin treatment of horses: Effect on the distribution of lawns and roughs in horse pastures.

Vet. Parasitol., 48, 241-246

DI PIETRO, J.A.; PAUL, A.J.; EWERT, K.M.; TODD; K.S., Jr.; LOCK, T.F.; AGUILAR, R.

(1993b):

Moxidectin gel: A new equine endectocide.

Proc. 38th Ann. Conf. Amer. Ass. Equine Pract. (30.11-2.12., Orlando / Florida), 311- 316

DI PIETRO, J.A.; HUTCHENS, D.E.; LOCK, T.F.; WALKER, K.; PAUL, A.J.; SHIPLEY, C.; RULLI, D. (1997):

Clinical trial of moxidectin oral gel in horses.

Vet. Parasitol., 72, 167-177

DOHERTY, W.M.; STEWART, N.P.; COBB, R.M.; KEIRAN, P.J. (1994):

In-vitro comparison of the larvicidal activity of moxidectin and abamectin against Onthophagus gazella Fabricius (Coleoptera: Scarabaeidae) and Haemotobia irritans exigua De Meijere (Diptera: Muscidae).

J. Aust. ent. Soc., 33, 71-74

DORCHIES, P; DUCOS DE LAHITTE, J.; FLOCHLAY, A.; BLOND-RIOU, F. (1998):

Efficacy of moxidectin 2 % equine gel against natural nematode infections in ponies.

Vet. Parasitol., 74, 85-89

DRÖSSIGK, U.; SCHUSTER, R. (1997):

Bandwurmbefall bei Pferden - Probleme und Erfahrungen.

Prakt. Tierarzt, 78:7, 564-567

DRUDGE, J.H.; ELAM, G. (1961):

Preliminary observations on the resistance of horse strongyles to phenothiazine.

J. Parasitol., 47, (Suppl.), 38-39

DRUDGE, J.H.; SZANTO, J.; WYANT, Z.N. (1962):

Studies on the anthelmintic thiabendazole in the horse: Critical tests.

Proc. IV. Pan-Am Congress of Veterinary Medicine and Zootechnics „*Seminar on parasitic diseases with special reference to thiabendazole*“ (13.11., Mexico City), 79-81

DRUDGE, J.H.; LYONS, E.T. (1965):

Newer development in helminth control and Strongylus vulgaris research.

Proc. 11th Ann. Meet. Amer. Ass. Equine Pract. (6.-8.12., Miami Beach / Florida), 381-389

DRUDGE, J.H.; LYONS, E.T. (1970):

Activity of dichlorvos against horse bots.

Mod. vet. Pract., 51, 45-46

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C. (1975):

Critical and controlled tests of the antiparasitic activity of liquid and paste formulations of trichlorfon in the horse.

Vet. Med. / Small Anim. Clin., 70, 537-540

DRUDGE, J.H.; LYONS, E.T. (1976):

Zur Pathologie des Endoparasitenbefalls beim Pferd.

Die blauen Hefte, 55, 219-227

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C. (1978):

Critical tests of the anthelmintic febantel in the horse: Activity of a paste formulation alone or with a trichlorfon paste.

Amer. J. vet. Res., 39, 1419-1421

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C. (1979):

Benzimidazole resistance of equine strongyles - Critical tests of six compounds against population B.

Amer. J. vet. Res., 40, 590-594

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C. (1981a):

Parasite control in horses: A summary of contemporary drugs.

Vet. Med. / Small Anim. Clin., 76, 1479-1489

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C.; KUBIS, J.E. (1981b):

*Clinical trials with fenbendazole and oxicabendazole for *Strongyloides westeri* infection in foals.*

Amer. J. vet. Res., 42, 526-527

DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C.; KUBIS, J.E. (1981c):

Clinical trials of oxicabendazole for control of equine internal parasites.

Mod. vet. Pract., 62, 679-682

- DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C.; KUBIS, J.E. (1984):
Febantel in horses: Activity in clinical trials featuring benzimidazole-resistant strongyles.
Equine Pract., 6, 19-28
- DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C.; SWERCZEK, T.W. (1985):
Use of oxicabendazole for control of cambendazole-resistant small strongyles in a band of ponies: A six year study.
Amer. J. vet. Res., 46, 2507-2511
- DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C.; LOWRY, S.R.; FALLON, E.H. (1988):
Piperazine resistance in population-B equine strongyles: A study of selection in thoroughbreds in Kentucky from 1966 through 1983.
Amer. J. vet. Res., 49, 986-994
- DRUDGE, J.H.; LYONS, E.T.; TOLLIVER, S.C. (1990):
Phenothiazine in the origin of benzimidazole resistance in population-B equine strongyles.
Vet. Parasitol., 35, 117-130
- DUNCAN, J.L (1974):
Field studies on the epidemiology of mixed strongyle infection in the horse.
Vet. Rec., 94, 337-345
- EDWARDS, G.B. (1986):
Surgical management of intussusception in the horse.
Equine vet. J., 18, 313-321
- EGERTON, J.R.; SEWARD, R.L.; ROBIN, B. (1984):
L'ivermectin: un agent antiparasitaire pour les chevaux.
Rec. Méd. Vét., 160, 595-599
- EISENMENGER, E.; KREISEL, A.; SCHÖNBAUER, M.; FREILER, I. (1986):
*Filarien (*Onchocerca reticulata*) als Ursache von granulomatösen Wundheilungsstörungen an den Extremitäten von Pferden.*
Wien. tierärztl. Mschr., 73, 244-247
- EL-SEIFY, M.A.; HASSLINGER, M.-A. (1991):
*Natural infections of equids with *Dictyocaulus arnfieldi* in Egypt.*
J. Egypt. vet. med. Ass., 51, 519-528
- ENDE, H.; STOYE, M. (1976):
Banminth-Paste zur planmäßigen Strongylidenbekämpfung beim Pferd.
Dtsch. tierärztl. Wschr., 83, 431-432

ENGLISH, A.W. (1979a):

The epidemiology of equine strongylosis in southern Queensland. 1. The bionomics of the free-living stages in faeces and on pasture.

Aust. vet. J., 55, 299-305

ENGLISH, A.W. (1979b):

The epidemiology of equine strongylosis in southern Queensland. 2. The survival and migration of infective larvae in herbage.

Aust. vet. J., 55, 306-309

ENGLISH, A.W. (1979c):

*The epidemiology of equine strongylosis in southern Queensland. 3. Seasonal variation in arterial populations of *Strongylus vulgaris*, and the prevalence of some helminths.*

Aust. vet. J., 55, 310-314

ENGLISH, A.W. (1979d):

The effect of dung beetles (Coleoptera - Scarabaeinae) on the free-living stages of strongylid nematodes of the horse.

Aust. vet. J., 55, 315-321

ENIGK, K. (1970):

Gesundheitsschädlichkeit und Vorbeuge des Parasitenbefalles beim Pferd: I. Der Strongylidenbefall.

Vollblut, Zucht und Rennen, 42, 149-151

ENIGK, K. (1971):

Gesundheitsschädlichkeit und Vorbeuge des Parasitenbefalles beim Pferd: III. Bandwürmer, Biesfliegenlarven und Bremsen.

Vollblut, Zucht und Rennen, 45, 42-44

ENIGK, K. (1983):

*Anmerkungen zur endogenen Entwicklung von *Strongylus vulgaris*.*

Berl. Münch. tierärztl. Wschr., 96, 228-231

ENIGK, K.; WEINGÄRTNER, E. (1973):

Zur Verbreitung und Behandlung der Dictyocaulose der Einhufer.

Dtsch. tierärztl. Wschr., 80, 145-148

EPE, C.; ISING-VOLLMER, S.; STOYE, M. (1993):

Ergebnisse parasitologischer Kotuntersuchungen von Equiden, Hunden, Katzen und Igeln der Jahre 1984-1991.

Dtsch. tierärztl. Wschr., 100, 426-428

EWERT, K.M.; DI PIETRO, J.A.; DANNER, C.S.; LAWRENCE, L.M. (1991):

Ivermectin treatment of horses: Effect on proportion of faecal-fouled areas in pastures.

Vet. Rec., 129, 140-141

EYSKER, M.; JANSEN, J.; WEMMENHOVE, R.; MIRCK, M.H. (1983):

Alternate grazing of horses and sheep as control for gastro-intestinal helminthiasis in horses.

Vet. Parasitol., 13, 273-280

EYSKER, M.; JANSEN, J.; MIRCK, M.H. (1984):

Inhibited development of cyathostominae in the horse in the early third stage.

Res. vet. Sci., 37, 355-356

EYSKER, M.; MIRCK, M.H. (1986):

The distribution of inhibited early third stage cyathostominae larvae in the large intestine of the horse.

Z. Parasitenk., 72, 815-820

EYSKER, M.; JANSEN, J.; MIRCK, M.H. (1986a):

Control of strongylosis in horses and sheep and some other aspects of the epidemiology of strongylidae infections.

Vet. Parasitol., 19, 103-105

EYSKER, M.; JANSEN, J.; KOOYMAN, F.N.J.; MIRCK, M.H.; WENSING, T. (1986b):

Comparison of two control systems for cyathostome infections in the horse and further aspects of the epidemiology of these infections.

Vet. Parasitol., 22, 105-112

EYSKER, M.; BOERSEMA, J.H., KOOYMAN, F.N.J.; BERGHEN, P. (1988):

Possible resistance of small strongyles from female ponies in the Netherlands against albendazole.

Amer. J. vet. Res., 49, 995-999

EYSKER, M.; BOERSEMA, J.H.; KOOYMAN, F.N.J. (1989):

Emergence from inhibited development of cyathostominae larvae in ponies following failure to remove them by repeated treatments with benzimidazole compounds.

Vet. Parasitol., 34, 87-93

EYSKER, M.; PANDY, V.S. (1990):

Observations on some parasites of the horses in Zimbabwe.

Zimbabwe vet. J., 21, 21-24

EYSKER, M.; VERCRUYSSE, J. (1990):

Epidemiologie, chemotherapie, anthelmintica-resistentie en preventie van strongylidae-infecties bij het paard.

T. Diergeneesk., 115, 891-907

EYSKER, M.; BOERSEMA, J.H.; KOOYMAN, F.N.J. (1990):

Seasonally inhibited development of cyathostominae nematodes in Shetland ponies in the Netherlands.

Vet. Parasitol., 36, 259-264

EYSKER, M.; BOERSEMA, J.H.; KOOYMAN, F.N.J. (1991):

Effect of early season ivermectin and pyrantel treatments on strongylid infections in young Shetland ponies in the Netherlands.

Vet. Parasitol., 38, 33-39

EYSKER, M.; BOERSEMA, J.H.; KOOYMAN, F.N.J. (1992):

The effect of ivermectin treatment against inhibited early third stage, late third stage and fourth stage larvae and adult stages of the cyathostomes in Shetland ponies and spontaneous expulsion of these helminths.

Vet. Parasitol., 42, 295-302

EYSKER, M.; BOERSEMA, J.H.; GRINWIS, G.C.M.; KOOYMAN, F.N.J.; POOT, J. (1997):

Controlled dose confirmation study of a 2 % moxidectin equine gel against equine internal parasites in The Netherlands.

Vet. Parasitol., 70, 165-173

FERNANDEZ, A.S.; LARSEN, M.; NANSEN, P.; GRONVOLD, J.; HENRIKSEN, S.A.;

WOLSTRUP, J. (1997):

*Effect of the nematode-trapping fungus *Duddingtonia flagrans* on the free-living stages of horse parasitic nematodes: a plot study.*

Vet. Parasitol., 73, 257-266

FINCHER, G.T.; WANG, G.T. (1992):

Injectable moxidectin for cattle: Effects on two species of dung burying beetles.

Southw. Ent., 17, 303-306

FOGERTY, U.; DEL PIERO, F.; PURNELL, R.E.; MOSURSKI, K.R. (1994):

*Incidence of *Anoplocephala perfoliata* in horses examined at an Irish abattoir.*

Vet. Rec., 134, 515-518

FORSTNER, M.J.; HASSLINGER, M.-A. (1974):

Untersuchungen über die Wirkung des Anthelminthikums FENBENDAZOL an Pferden.

Berl. Münch. tierärztl. Wschr., 87, 325-327

FRAHM, J. (1982):

Gastroskopische Untersuchungen bei Pferden zur Feststellung der Wirksamkeit von Anthelminthika gegen Gasterophiluslarven.

Vet. med. Diss., Hannover

FRAHM, J. (1983)

Gasterophilose (Magendasselbefall) beim Pferd.

Tierärztl. Prax., 11, 201-207

FRAHM, J. (1986):

Zur Wirksamkeit von Ivermectin auf den Gasterophiluslarvenbefall bei Island-Ponies.

Tierärztl. Umsch., 41, 277-281

FRANK, W. (1976):

Parasitologie.

Verlag Eugen Ulmer, Stuttgart

FRENCH, D.D.; KLEI, T.R.; TAYLOR, H.W.; CHAPMAN, M.R.; WRIGHT, F.R. (1988):

Efficacy of ivermectin in the oral paste formulation against naturally acquired adult and larval stages of Parascaris equorum in pony foals.

Amer. J. vet. Res., 49, 1000-1003

FRENCH, D.D.; KLEI, T.R.; TAYLOR, H.W.; CHAPMAN, M.R. (1989):

Efficacy of ivermectin in oral drench and paste formulation against migrating larvae of experimentally inoculated Parascaris equorum.

Amer. J. vet. Res., 50, 1071-1073

FRENCH, D.D.; KLEI, T.R.; TAYLOR, H.W.; CHAPMAN, M.R.; MONAHAN, C.; AGUILAR, R. (1992):

Efficacy of moxidectin gel against internal parasites of ponies.

Proc. 37th Ann. Meet. Amer. Ass. Vet. Parasit. (2.-4.8., Boston / Massachusetts), 51

FRENCH, D.D.; CHAPMAN, M.R.; KLEI, T.R. (1994):

Effects of treatment with ivermectin for five years on the prevalence of Anoplocephala perfoliata in three Louisiana pony herds.

Vet. Rec., 135, 63-65

GAUDERON, H.; KIPFER, H.; HOFER, B.; Pfister, K. (1988):

Efficacité du pamoate de pyrantel (Strongid®) dans le traitement des anoplocéphales du cheval.

Schweiz. Arch. Tierheilk., 128, 321-328

GAWOR, J.J. (1995):

The prevalence and abundance of internal parasites in working horses autopsied in Poland.

Vet. Parasitol. 58, 99-108

GENCHI, C.; SOLARI BASANO, F.; NOGARA, B. (1995):

Efficacy and persistency of Moxidectin 2 % Equine gel against small strongyles (cyathostominae) in naturally infected horses.

Abstr. 15th Int. Conf. WAAVP (30.8.-2.9., Yokohama / Japan), 101

GEORGI, J.R.; RENDANO, V.T.; KING, J.M.; BIANCHI, D.G.; THEODORIDES, V.J. (1980):

Equine verminous arteritis: Efficiency and speed of larvicidal activity as influenced by dosages of albendazole.

Cornell. Vet., 70, 147-152

GERHARDS, H.; ROSENBRUCH, M. (1984):

Intermittierendes Hinken beim Pferd - Diskussion ätiologischer und therapeutischer Aspekte anhand eines Fallberichtes.

Prakt. Tierarzt, 65, 645-653

GIANGASPERO, A.; LIA, R.; VOVLAS, N.; OTRANTA, D. (1996):

Thelazia lacrymalis in native horses: First report in Italy.

Parassitologia, 38, 427

GIBSON, T.E. (1953):

The effect of repeated anthelmintic treatment with phenothiazine on the faecal egg counts of housed horses, with some observations on the life cycle of Trichonema spp. in the horse.

J. Helminth., 27, 29-40

GILES, C.J.; URQUHART, K.A.; LONGSTAFFE, J.A. (1985):

Larval cyathostomiasis (immature trichonema induced enteropathy): A report of 15 clinical cases.

Equine vet. J., 17, 196-201

GODBER, L.M.; DERKSEN, F.J.; WILLIAMS, J.F.; MAHMOUD, B. (1995):

Ivermectin toxicosis in a neonatal foal.

Aust. vet. J., 75, 191-192

GOETZ, T.E. (1984):

Dictyocaulus arnfieldi as a possible cause of chronic cough in 14 horses.

Equine Pract., 6, 33-38

GOUDIE, A.; EVANS, N.A.; GRATTON, K.A.F.; BISHOP, B.F.; GIBSON, S.P.; HOLDOM, K.S.;

KAYE, B.; WICKS, S.R.; LEWIS, D.; WEATHERLEY, A.; BRUCE, C.I.; HERBERT, A.;

SEYMOUR, D.J. (1993):

Doramectin - a potent novel endectocide.

Vet. Parasitol., 48, 5-15

GREATOREX, J.C. (1975):

*Diarrhoea in horses associated with ulceration of the colon and caecum resulting from *S. vulgaris* migration.*

Vet. Rec., 97, 221-225

GRELCK, H.; HÖRCHNER, F.; WÖHRL, H.E. (1977):

Entwicklungsfähigkeit und Überlebensdauer von Larven der Pferdestrongylidenarten.

Prakt. Tierarzt, 58, 265-268

HABEDANK, B.; ILCHMANN, G.; SCHEIN, E. (1999):

Untersuchungen zum Vorkommen von Gasterophilus-Larven (Diptera: Gasterophilidae) bei

Schlachtpferden in Nordostdeutschland.

Prakt. Tierarzt, 80, 6, 488-497

HARDER, A.; ANDREWS, P.; THOMAS, H. (1987):

Praziquantel - mode of action.

Biochem. Soc. trans., 15, 68-70

HASSLINGER, M.-A. (1963):

Untersuchung an Schlachtpferden über Vorkommen und Sitz verschiedener kleiner Strongylidenarten.

Vet. med. Diss., FU Berlin

HASSLINGER, M.-A. (1981a):

Untersuchungen über den Einfluß verschiedener Temperaturen auf Eier und Larven von Pferdestrongyliden unter Laboratoriumsbedingungen sowie das Verhalten dieser exogenen Stadien auf der Weide.

Berl. Münch. tierärztl. Wschr., 94, 1-5

HASSLINGER, M.-A. (1981b):

Über das Wanderverhalten und Vorkommen von Larven der Pferdestrongyliden sowie deren Bedeutung für das Weidemanagement.

Berl. Münch. tierärztl. Wschr., 94, 329-333

HASSLINGER, M.-A. (1984):

Control of endoparasites in horses.

Equine Pract., 6, 23-29

HASSLINGER, M.-A. (1985):

The quantitative portion of different species of small equine strongyles and their reduced sensitivity against benzimidazole.

Proc. 11th Conf. WAAVP (5.-9.8., Rio de Janeiro / Brazil), 191

HASSLINGER, M.-A. (1986):

Biologische und epizootologische Aspekte zu Parasitenbefall und -bekämpfung beim Pferd.

Prakt. Tierarzt, 67, 779-780, 789-799

HASSLINGER, M.-A. (1988):

Strongyloidosis.

In: *Handlexikon der tierärztlichen Praxis I-V*, Medical Book Company, Bjaeverskov / DK,
Lief. 184, 815d-817

HASSLINGER, M.-A. (1989a):

Magenwürmer.

In: *Handlexikon der tierärztlichen Praxis I-V*, Medical Book Company, Bjaeverskov / DK,
Lief. 185, 563a-563b

HASSLINGER, M.-A. (1989b):

Speziesabhängige Aspekte zur Epizootiologie von Oxyureninfektionen beim Pferd und Möglichkeiten der Bekämpfung.

Pferdeheilk., 5, 15-22

HASSLINGER, M.-A. (1989c):

Dictyocaulus arnfieldi in equines: Present situation and future aspects.

J. Egypt. vet. med. Ass., 49, 445-455

HASSLINGER, M.-A. (1991):

Askaridiasis.

In: *Handlexikon der tierärztlichen Praxis*, I-V, Gustav Fischer Verlag, Stuttgart / New York,
Lief. 187, 50x-50i

HASSLINGER, M.-A. (1995):

Endo- und Ektoparasiten des Pferdes.

In: *Handbuch Pferd*, 5. Aufl., BLV Verlagsgesellschaft mbH München / Wien / Zürich,
804-824

HASSLINGER, M.-A.; JONAS, D. (1975):

Control of Gasterophilus intestinalis (De Geer, 1776) with Dichlorvos.

Brit. vet. J., 131, 89-93

HASSLINGER, M.-A.; MÜLLER, R. (1978):

Rintal ®, ein neues Anthelminthikum zur Bekämpfung der Pferdestrongyliden.

Vet.-med. Nachr., H.2., 186-191

HASSLINGER, M.-A.; BARTH, D. (1982):

Untersuchungen zur Wirksamkeit von IVERMECTIN gegen Endoparasiten des Pferdes.

Dtsch. tierärztl. Wschr., 89, 62-65

HASSLINGER, M.-A.; BITTNER, G. (1984):

Zur Saisondynamik der Larven von Pferdestrongyliden und deren Beziehung zum Infektionsrisiko auf der Weide.

Zbl. Vet. Med. B., 31, 25-31

HASSLINGER, M.-A.; TAUSEND, S. (1989):

Zur Bedeutung und Bekämpfung des Bandwurmbefallen beim Pferd.

Prakt. Tierarzt, 70, 26-31

HEARN, F.P.D.; HEARN, E.E. (1995):

A simple diagnostic technique to better determine the prevalence of tapeworms.

J. Equine vet. Sci., 15, 96-98

HEDJAZI, M.; MIRZAYANS, A. (1978):

Parafilariose équine dans la région de Téhéran (Iran). Aspect clinique et traitement.

Rev. méd. Vét., 129, 1685-1686, 1689-1691

HEIL, H.G. (1983):

Zur intestinalen Parasitenfauna und Dictyocaulus arnfieldi-Infektion der Esel in Hessen sowie zur Saisondynamik der Befallsextensität und -intensität.

Vet. med. Diss., Gießen

HERD, R.P. (1986a):

Epidemiology and control of equine strongylosis at Newmarket.

Equine vet. J., 18, 447-452

HERD, R.P. (1986b):

Epidemiology and control of equine parasites in northern temperate regions.

Vet. Clin. North Amer. / Equine Pract., 2, 337-355

HERD, R.P. (1990):

Equine parasite control - solutions to anthelmintic associated problems.

Equine vet. Educ., 2, 86-91

HERD, R.P. (1992):

Choosing the optimal equine anthelmintic.

Vet. Med., 87, 231-239

HERD, R.P. (1993):

Control strategies for ruminant and equine parasites to counter resistance, encystment, and ecotoxicity in the USA.

Vet. Parasitol., 48, 327-336

HERD, R.P. (1995a):

Equine parasite control: Keeping up with evolution.

Vet. Med., 90, 477-480

HERD, R.P. (1995b):

A 10-point plan for equine worm control.

Vet. Med., 90, 481-485

HERD, R.P.; DONHAM, J.C. (1981):

Efficacy of ivermectin against cutaneous Draschia and Habronema infection (summer sores) in horses.

Amer. J. vet. Res., 42, 1953-1955

HERD, R.P.; DONHAM, J.C. (1983):

Efficacy of ivermectin against Onchocerca cervicalis microfilarial dermatitis in horses.

Amer. J. vet. Res., 44, 1102-1105

HERD, R.P.; KENT, J.E. (1986):

Serum protein changes in ponies on different parasite control programmes.

Equine vet. J., 18, 453-457

HERD, R.P.; GABEL, A.A. (1990):

Reduced efficacy of anthelmintics in young compared with adult horses.

Equine vet. J., 22, 164-169

HERD, R.P.; JERNIGAN, A.D.; STINNER, B.R.; PURRINGTON, F.F. (1990):

Ecotoxicity of anthelmintics.

Proc. 35th Ann. Meet. Amer. Ass. Vet. Parasitol. (21.-24.7., San Antonio), 37, 37

HERD, R.P.; STINNER, B.R.; PURRINGTON, F.F. (1993):

Dung dispersal and grazing area following treatment of horses with a single dose of ivermectin.

Vet. Parasitol., 48, 229-240

HERD, R.P.; MAJEWSKI, G.A. (1994):

Comparison of daily and monthly pyrantel treatment in yearling thoroughbreds and the protective effect of strategic medication of mares on their foals.

Vet. Parasitol., 55, 93-104

HERD, R.P.; COLES, G.C. (1995):

Slowing the spread of anthelmintic resistant nematodes of horses in the United Kingdom.

Vet. Rec., 136, 481-485

HÖGLUND, J.; NILSSON, O.; LJUNGSTRÖM, B.-L.; HELLANDER, J.; OSTERMANN LIND, E.; UGGLA, A. (1998):

Epidemiology of Anoplocephala perfoliata infection in foals on a stud farm in south-western Sweden.

Vet. Parasitol., 75, 71-79

HOPFER, S.M.; VAN KRUININGEN, H.J.; DANIELS, W.H. (1984):

The elimination of equine strongyles and hematological and pathological consequences following larvical doses of thiabendazole.

Vet. Parasitol., 14, 21-32

JACH, T.; ALLMELING, G. (1990):

Obturationsileus des Jejunums bei einer Traberstute aufgrund eines submukösen Hämatoms, hervorgerufen durch Massenbefall mit Anoplocephala perfoliata.
Pferdeheilk., 6, 89-92

JACOBS, D.E.; HUTCHINSON, M.J.; PARKER, L.; GIBBONS, L.M. (1995):

Equine cyathostome infection: Suppression of faecal egg output with moxidectin.
Vet. Rec., 137, 545

JONAS, D.; HASSSLINGER, M.-A.; WOLPERT, E. (1972):

Neue Aspekte zur Endoparasitenbekämpfung beim Pferd.
Prakt. Tierarzt, 53, 427-430

JONAS, D.; HASSSLINGER, M.-A. (1973):

Erfolgsaussichten einer planmäßigen Gasterophilus-Bekämpfung bei Pferden.
Dtsch. tierärztl. Wschr., 80, 369-372

JONES, R.S.; LAWRENCE, T.L.J.; VEEVERS, A.; CLEAVE, N.; HALL, J. (1989):

Accuracy of prediction of the liveweight of horses from body measurements.
Vet. Rec., 125, 549-553

JONES, R.M.; LOGAN, N.B.; WEATHERLEY, A.J.; LITTLE, A.S.; SMOTHERS, C.D. (1993):

Activity of doramectin against nematode endoparasites of cattle.
Vet. Parasitol., 49, 27-37

JORDAN, H.E.; SLUSHER, S.H. (1990):

No apparent resistance to oxicabendazole in horses during 3-year field study.
Top. vet. Med., 1, 35-37

KARNS, P.A.; LUTHER, D.G. (1984):

A survey of adverse effects associated with ivermectin use in Louisiana horses.
J. Amer. vet. med. Ass., 185, 782-783

KELLY, J.D.; BAIN, S.A. (1975):

Critical test evaluation of micronized mebendazole against A. perfoliata in the horse.
N. Z. vet. J., 23, 229-232

KELLY, J.C.; FOGARTY, U.M.G. (1993):

Outbreak of larval cyathostomiasis on a thoroughbred stud farm.
Irish vet. J., 46, 133-136

KENT, J.E.; BLACKMOORE, D.J. (1985):

Turbidimetric measurement of IgG(T) in the serum of healthy thoroughbreds and ponies.
Equine vet. J., 17, 119-124

KESTER, W.O. (1975):

Strongylus vulgaris - the horse killer.

Mod. vet. Pract., 56, 569-572

KIERMEYER, I. (1990):

Natürliche Fasciola hepatica-Infektionen beim Pferd mit Berücksichtigung der Chemotherapie unter enzymatischer und serologischer Kontrolle.

Vet. med. Diss., München

KLEI, T.R. (1986a):

Development of vaccines against equine helminths.

Parasitol. Today, 2, 80-81

KLEI, T.R. (1986b):

Immunity and potential of vaccination.

Vet. Clin. North Amer. / Equine Pract., 2, 395-402

KLEI, T.R.; TORBERT, B.J., CHAPMAN, M.R.; OCHOA, R. (1982):

*Irradiated larval vaccination of ponies against *Strongylus vulgaris*.*

J. Parasitol., 68, 561-569

KLEI, T.R.; TURK, M.A.M.; TORBERT, B.J.; CHAPMAN, M.R. (1983):

*Ivermectin in parenteral and oral paste formulations: Efficacy against gastrointestinal parasites, migrating larvae of *Strongylus vulgaris*, and post-treatment responses of infected ponies.*

Proc. MSD Symposium XXII World Vet. Congr. (19.-31.8., Perth / Australia), 271-285

KLEI, T.R.; TORBERT, B.J.; CHAPMAN, M.R.; TURK, M.A.M. (1984):

*Efficacy of ivermectin in injectable and oral paste formulations against eight-week-old *Strongylus vulgaris* larvae in ponies.*

Amer. J. vet. Res., 45, 183-185

KLEI, T.R.; FRENCH, D.D.; CHAPMAN, M.R.; McCLURE, J.R.; DENNIS, V.A.; TAYLOR, H.W.; HUTCHINSON, G.W. (1989):

*Protection of yearling ponies against *Strongylus vulgaris* by foalhood vaccination.*

Equine vet. J. (Suppl.), 7, 2-7

KLEI, T.R.; TURK, M.A.M.; McCLURE, J.R.; HOLMES, R.A.; DENNIS, V.A.; CHAPMAN, M.R. (1990):

*Effects of repeated *Strongylus vulgaris* inoculations and concurrent ivermectin treatments on mesenteric arterial lesions in pony foals.*

Amer. J. vet. Res., 51, 654-660

- KLEI, T.R.; CHAPMAN, M.R.; FRENCH, D.D.; MONAHAN, C.; TAYLOR, H.W. (1992):
Protective resistance against strongyle infections in pony foals maintained on Strongid C supplemented feed.
Proc. 37th Ann. Meet. Amer. Ass. Vet. Parasitol., (2.-4.8., Boston, Massachusetts), 48
- KLEI, T.R.; CHAPMAN, M.R.; FRENCH, D.D.; TAYLOR, H.W. (1993):
Evaluation of ivermectin at an elevated dose against encysted equine cyathostome larvae.
Vet. Parasitol., 47, 99-106
- KÖHLER, M.; HIEPE, T. (1986):
Wirksamkeit von Ivermectin auf den Strongyloides westeri-Befall beim Fohlen.
Angew. Parasitol., 27, 181-186
- KÖHLER, M.; BUCHWALDER, R.; HIEPE, T. (1989):
Untersuchungen zur Wirksamkeit von Ivermectin im Vergleich zu Mebendazol auf Nematodenbefall beim Pferd unter Gestütsbedingungen.
Mh. Vet.-Med., 44, 383-386
- KOPF, N. (1990):
Protrahierte Magenruptur infolge eines Dünndarmileus und intramurales Hämatom des Jejunums in Verbindung mit Zestodenbefall bei einem Pferd.
Pferdeheilk., 6, 169-171
- KOZLOV, D.P. (1985):
Über die Verteilung von Strongylidenlarven auf der Weide. (In Russisch).
Parazitologiya, 19, 220-225
- LARSEN, N.L. (1990):
Bændelormeinfektion hos hest.
Dansk Veterinærtidsskrift, 73, 404
- LARSEN, M.; NANSEN, P.; HENRIKSEN, S.A.; WOLSTRUP, J.; GRØNVOLD, J.; ZORN, A.; WEDØ, E. (1995):
*Predacious activity of the nematode-trapping fungus *Duddingtonia flagrans* against cyathostome larvae in faeces after passage through the gastrointestinal tract of horses.*
Vet. Parasitol., 60, 315-320
- LICHTENFELS, J.R. (1975):
Helminths of domestic equids.
Proc. helminth. Soc. Wash., 42 (Special issue), 1-92

LICHTENFELS, J.R.; KHARCHENKO, V.A.; KRECEK, R.C.; GIBBONS, L.M. (1998):

An annotated checklist by genus and species of 93 species level names for 51 recognized species of small strongyles (Nematoda: Strongylidea: Cyathostominea) of horses, asses and zebras of the world.

Vet. Parasitol., 79, 65-79

LLOYD, E.T.; SOULSBY, E.J.L. (1978):

Survey for infection with Onchocerca cervicalis in horses in eastern United States.

Amer. J. vet. Res., 39, 1962-1963

LORENZ, H. (1996):

Gasterophilose beim Pferd - ein Beitrag zur Intravitaldiagnostik und medikamentösen Bekämpfung.

Vet. med. Diss., Berlin

LOUW, J.P.; MEYER, S.; SCHRÖDER, J. (1980):

A critical efficacy test cambendazole in equids - The use of the geometric means to assess efficacy.

J. S. Afr. vet. med. Ass., 51, 259-261

LOVE, S. (1992):

The role of equine strongyles in the pathogenesis of colic and current options for prophylaxis.

Equine vet. J., 13 (Suppl.), 5-9

LOVE, S.; DUNCAN, J.L. (1992):

The development of naturally acquired cyathostome infection in ponies.

Vet. Parasitol., 44, 127-142

LOVE, S.; MAIR, T.S.; HILLYER, M.H. (1992a):

Chronic diarrhoea in adult horses: A review of 51 referred cases.

Vet. Rec., 130, 217-219

LOVE, S.; ESCALA, J.; DUNCAN, J.L.; McLEAN, J.M. (1992b):

Studies on the pathogenic effects of experimental cyathostome infections in ponies.

In: W. PLOWRIGHT, P.O. ROSSDALE, J.F. WADE, *Equine infectious diseases*, 6th Ed., R and W Publishers, Newmarket/ UK, 149-155

LOVE, S.; DUNCAN, J.L.; PARRY, J.M.; GRIMSHAW, W.T.R. (1995):

Efficacy of oral ivermectin paste against mucosal stages of cyathostomes.

Vet. Rec., 136, 18-19

LUDWIG, K.G.; CRAIG, T.M.; BOWEN, J.M.; ANSARI, M.M.; LEY, W.B. (1983):

*Efficacy of ivermectin in controlling *Strongyloides westeri* infections in foals.*

Amer. J. vet. Res., 44, 314-316

LYONS, E.T. (1994):

Vertical transmission of nematodes: Emphasis on Uncinaria lucasi in northern fur seals and Strongyloides westeri in equids.

J. helminth. Soc. Wash., 61, 169-178

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1973):

On the life cycle of Strongyloides westeri in the equine.

J. Parasitol., 59, 780-787

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1974):

Critical tests of three salts of pyrantel against internal parasites of the horse.

Amer. J. vet. Res., 35, 1515-1522

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1975):

Field tests of three salts of pyrantel against internal parasites of the horse.

Amer. J. vet. Res., 36, 161-166

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1976a):

Studies on the development and chemotherapy of larvae of Parascaris equorum (Nematoda: Askarididae) in experimentally and naturally infected horses.

J. Parasitol., 62, 453-459

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1976b):

Critical tests of anthelmintic activity of a paste formulation of thiabendazole in horses.

Amer. J. vet. Res., 37, 701-702

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1977):

Critical tests of oxfendazole against internal parasites of horses.

Amer. J. vet. Res., 38, 2049-2053

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1980a):

Antiparasitic activity of parbendazole in critical tests in horses.

Amer. J. vet. Res., 41, 123-124

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1980b):

Antiparasitic activity of ivermectin in critical tests in equids.

Amer. J. vet. Res., 41, 2069-2072

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1980c):

*Experimental infections of Thelazia lacrymalis: Maturation of third-stage larvae from face flies (*Musca autumnalis*) in eyes of ponies.*

J. Parasitol., 66, 181-182

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1981a):

*Prevalence of microfilariae (*Onchocerca* spp.) in skin of Kentucky horses at necropsy.*

J. Amer. vet. med. Ass., 179, 899-900

LYONS, E.T.; DRUDGE, J.H.; SWERCZEK, T.W.; CROWE, M.W.; TOLLIVER, S.C. (1981b):

Prevalence of strongylus vulgaris and Parascaris equorum in Kentucky thoroughbreds at necropsy.

J. Amer. vet. med. Ass., 179, 818-819

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1982):

Ivermectin: Activity against larval Strongylus vulgaris and adult Trichostrongylus axei in experimental infections in ponies.

Amer. J. vet. Res., 43, 1449-1450

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1983a):

Controlled tests with fenbendazole in equids: Special interest on activity of multiple doses against natural infections of migrating stages of strongyles.

Amer. J. vet. Res., 44, 1058-1063

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; SWERCZEK, T.W.; CROWE, M.W. (1983b):

Parasites in Kentucky thoroughbreds at necropsy: Emphasis on stomach worms and tapeworms.

Amer. J. vet. Res., 44, 839-844

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C.; SWERCZEK, T.W.; CROWE, M.W. (1984):

Prevalence of Anoplocephala perfoliata and lesions of Draschia megastoma in thoroughbreds in Kentucky at necropsy.

Amer. J. vet. Res., 45, 996-999

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1985):

Ivermectin: Treating for naturally occurring infections of lungworms and stomach worms in equids.

Vet. Med. / Small Anim. Clin., 80, 58-64

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; SWERCZEK, T.W.; CROWE, M.W. (1986a):

*Eyeworms (*Thelazia lacrymalis*) in one-to-four-year-old thoroughbreds at necropsy in Kentucky (1984 to 1985).*

Amer. J. vet. Res., 47, 315-316

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1986b):

Fenbendazole in equids: Further controlled tests with emphasis on activity of multiple doses against naturally occurring infections of migratory large strongyles.

Amer. J. vet. Res., 47, 317-321

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C.; SWERCZEK, T.W. (1986c):

Pyrantel pamoate: Evaluation its activity against equine tapeworms.

Vet. Med., 81, 280-285

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1986d):

Prevalence of Dictyocaulus arnfieldi (nematoda) in equids and clinical problems with Strongylus vulgaris (nematoda) mainly in donkeys on a farm in central Kentucky.
Proc. helminth. Soc. Wash., 53, 251-253

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; SWERCZEK, T.W. (1988):

Oxfendazole: activity of a two-dose treatment regime on natural infection of internal parasites of equids, with emphasis on migrating large strongyles in controlled tests in 1986 and 1987.

Amer. J. vet. Res., 49, 1923-1927

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C.; SWERCZEK, T.W.; COLLINS, S.S. (1989):

Determination of the efficacy of pyrantel pamoate at the therapeutic dose rate against the tapeworm Anoplocephala perfoliata in equids using a modification of the critical test method.
Vet. Parasitol., 31, 13-18

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C. (1990):

Prevalence of some internal parasites found (1971-1989) in horses born on a farm in central Kentucky.

J. Equine vet. Sci., 10, 99-107

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; GRANSTORM, D.E.; STAMPER, S. (1992a):

Activity of praziquantel against Anoplocephala perfoliata (cestoda) in horses.

J. helminth. Soc. Wash., 59, 1-4

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; GRANSTORM, D.E.; COLLINS, S.S.; STAMPER, S. (1992b):

Critical and controlled tests of activity of moxidectin (CL 301, 423) against natural infections of internal parasites of equids.

Vet. Parasitol., 41, 255-284

LYONS, E.T.; TOLLIVER, S.C.; DRUDGE, J.H.; GRANSTORM, D.E.; COLLINS, S.S. (1993):

Natural infections of Strongyloides westeri: Prevalence in horse foals on several farms in central Kentucky in 1992.

Vet. Parasitol., 50, 101-107

LYONS, E.T.; DRUDGE, J.H.; TOLLIVER, S.C.; SWERCZEK, T.W.; STAMPER, S.;

GRANSTORM, D.E. (1994a):

Control of cambendazole-resistant small strongyles (Population S) with oxicabendazole in a pony band: An 8 year field test (1984-1992).

Vet. Parasitol., 52, 271-277

LYONS, E.T.; SWERCZEK, T.W.; TOLLIVER, S.C.; DRUDGE, J.H.; STAMPER, S.; GRANSTORM, D.E.; HOLLAND, R.E. (1994b):

A study of natural infections of encysted small strongyles in a horse herd in Kentucky.
Vet. Med., 89, 1146-1155

LYONS, E.T.; TOLLIVER, S.C.; ENNIS, L.E. (1998):

*Efficacy of praziquantel (0,25 mg/kg) on the cecal tapeworm (*Anoplocephala perfoliata*) in horses.*
Vet. Parasitol., 78, 287-289

MAIR, T.S. (1993):

Recurrent diarrhoea in aged ponies associated with larval cyathostomiasis.
Equine vet. J., 25, 161-163

MAIR, T.S. (1994):

Outbreak of larval cyathostomiasis among a group of yearling and two-year-old horses.
Vet. Rec., 135, 598-600

MAIR, T.S.; DE WESTERLAKEN, L.V.; CRIPPS, P.J.; LOVE, S. (1990):

Diarrhoea in adult horses: A survey of clinical cases and assessment of some prognostic indices.
Vet. Rec., 126, 479-481

MAIR, T.S.; CRIPPS, P.J.; RICKETTS, S.W. (1993):

Diagnostic and prognostic value of serum protein electrophoresis in horses with chronic diarrhoea.
Equine vet. J., 25 (4), 324-326

MANCEBO, O.A.; VERDI, J.H.; BULMAN, G.M. (1997):

*Comparative efficacy of moxidectin 2 % equine oral gel and ivermectin 2 % equine oral paste against *Onchocerca cervicalis* (Railliet and Henry, 1910) microfilariae in horses with naturally acquired infections in Formosa (Argentina).*
Vet. Parasitol., 73, 243-248

MANNERS, H. (1989):

Efficacy of anthelmintics in horses.
Vet. Rec., 125, 584-585

MARTIN, R.J. (1996):

*An electrophysiological preparation of *Ascaris suum* pharyngeal muscle reveals a glutamate-gated chloride channel sensitive to the avermectin analogue, milbemycin D.*
Parasitol., 112, 247-252

MARTIN, R.J.; PENNINGTON, A.J. (1989):

A patch-clamp study of effects of dihydroavermectin on Ascaris muscle.

Brit. J. Pharmac., 98, 747-756

MATTHEWS, A.G.; MORRIS, J.R. (1995):

Cyathostomiasis in horses.

Vet. Rec., 136, 52

MAYHEW, I.G.; LICHTENFELS, J.R.; GREINER, E.C.; MCKAY, R.J.; ENLOE, C.W. (1982):

Migration of a spiruroid nematode through the brain of a horse.

J. Amer. vet. med. Ass., 180, 1306-1311

MAYHEW, I.G.; BREWER, B.D.; REINHARD, M.K.; GREINER, E.C. (1984):

*Verminous (*Strongylus vulgaris*) myelitis in a donkey.*

Cornell Vet., 74, 30-37

McCRAW, B.M.; SLOCOMBE, J.O.D. (1974):

*Early development of and pathology associated with *Strongylus edentatus*.*

Canad. J. comp. Med., 38, 124-138

McCRAW, B.M.; SLOCOMBE, J.O.D. (1978):

Strongylus edentatus: Development and lesions from ten weeks postinfection to patency.

Canad. J. comp. Med., 42, 340-356

McCRAW, B.M.; SLOCOMBE, J.O.D. (1985):

Strongylus equinus: Development and pathological effects in the equine host.

Canad. J. comp. Med., 49, 372-383

MEANA, A.; LUZON, M.; CORCHERO, J.; GÓMEZ-BAUTISTA, M. (1998):

*Reliability of coprological diagnosis of *Anoplocephala perfoliata* infection.*

Vet. Parasitol., 74, 79-83

MEZGER, S (1991):

Laborversuche zum Nachweis kleiner Strongyliden mit Hilfe des Enzyme-Linked-

Immunosorbent-Assay (ELISA).

Vet. med. Diss., München

MFITILODZE, M.W.; HUTCHINSON, G.W. (1985):

The site distribution of adult strongyle parasites in the large intestines of horses in tropical Australia.

Int. J. Parasitol., 15, 313-319

MIRCK, M.H. (1977):

Cyathostominose: Een vorm van ernstige strongylidose.

T. Diergeneesk., 102, 932-934

MIRCK, M.H. (1978):

Parasitologisch faecesonderzoek bij paarden en ponies.

T. Diergeneesk., 103, 991-997

MONAHAN, C.M.; TAYLOR, H.W.; CHAPMAN, M.R.; KLEI, T.R. (1994):

*Experimental immunization of ponies with *Strongylus vulgaris* radiation-attenuated larvae or crude soluble somatic extracts from larval or adult stages.*

J. Parasitol., 80, 911-923

MONAHAN, C.M.; CHAPMAN; M.R.; FRENCH, D.D.; KLEI, T.R. (1995a):

*Efficacy of moxidectin oral gel against *Onchocerca cervicalis* microfilariae.*

J. Parasitol., 81, 117-118

MONAHAN, C.M.; CHAPMAN, M.R.; FRENCH, D.D.; TAYLOR, H.W.; KLEI, T.R. (1995b):

Dose titration of moxidectin oral gel against gastrointestinal parasites of ponies.

Vet. Parasitol., 59, 241-248

MONAHAN, C.M.; CHAPMAN, M.R.; TAYLOR, H.W.; FRENCH, D.D.; KLEI, T.R. (1995c):

*Dose titration of moxidectin oral gel against migrating *Strongylus vulgaris* and *Parascaris equorum* larvae in pony foals.*

Vet. Parasitol., 60, 103-110

MONAHAN, C.M.; CHAPMAN, M.R.; TAYLOR, H.W.; FRENCH, D.D.; KLEI, T.R. (1996):

Comparison of moxidectin oral gel and ivermectin oral paste against a spectrum of internal parasites of ponies with special attention to encysted cyathostome larvae.

Vet. Parasitol., 63, 225-235

MONAHAN, C.M.; CHAPMAN, M.R.; TAYLOR, H.W.; FRENCH, D.D.; KLEI, T.R. (1997):

Foals raised on pasture with or without daily pyrantel tartrate feed additive: comparison of parasite burdens and host responses following experimental challenge with large and small strongyle larvae.

Vet. Parasitol., 73, 277-289

MONAHAN, C.M.; CHAPMAN, M.R.; TAYLOR, H.W.; FRENCH, D.D.; KLEI, T.R. (1998):

Experimental cyathostome challenge of ponies maintained with or without benefit of daily pyrantel tartrate feed additive: comparison of parasite burdens, immunity and colonic pathology.

Vet. Parasitol., 74, 229-241

MURPHY, D.; EDWARDS, S.E.; RUSSELL, T.; SAMMIN, D.; LOVE, S. (1996):

Diagnosis, management and outcome of 18 cases of chronic diarrhoea in adult horses.

Irish vet. J., 49, 216-220

MURPHY, D.; LOVE, S. (1997):

The pathogenic effects of experimental cyathostome infections in ponies.

Vet. Parasitol., 70, 99-110

MUYLLE, E.; OYAERT, W.; ROGIERS, M. (1979):

Endoskopische Untersuchung des Pferdemagens bei der Wirksamkeitsprüfung einer

Mebendazol / Trichlorfon-Paste gegen Gasterophilus intestinalis-Larven.

Vlaams Diergeneesk. T., 48, 279-282

NILSSON, O.; LJUNGSTRÖM, B.-L.; HÖGLUND, J.; LUNDQUIST, H.; UGGLA, A. (1995):

Anoplocephala perfoliata in horses in Sweden: Prevalence, infection levels and intestinal lesions.

Acta. vet. scand., 36, 319-328

OGBOURNE, C.P. (1973):

Survival on herbage plots of infective larvae of strongylid nematodes on the horse.

J. Helminth., 47, 9-16

OGBOURNE, C.P. (1975a):

Studies on the epidemiology of Strongylus vulgaris infection of the horse.

Int. J. Parasitol., 5, 423-426

OGBOURNE, C.P. (1975b):

Epidemiological studies on horses infected with nematodes of the family trichonematidae (Witenberg 1925).

Int. J. Parasitol., 5, 667-672

OGBOURNE, C.P. (1976):

The prevalence, relative abundance and site distribution of nematodes of the subfamily cyathostominae in horse killed in Britain.

J. Helminth., 50, 203-214

OGBOURNE, C.P. (1978):

Pathogenesis of cyathostome (trichonema) infections of the horse. A review.

Misc. publ. 5, Commonwealth Inst. Helminth., Glasgow, 26

OGBOURNE, C.P.; DUNCAN, J.L. (1985):

Strongylus vulgaris in the horse: Its biology and veterinary importance.

Misc. publ. 9, Commonwealth, Inst. Parasit., Glasgow, 26

OWEN, R.H.; JAGGER, D.W.; QUAN-TAYLOR, R. (1988):

Prevalence of Anoplocephala perfoliata in horses and ponies in Clwyd, Powys and adjacent English marches.

Vet. Rec., 123, 562-563

OWEN, R.H.; JAGGER, D.W.; QUAN-TAYLOR, R. (1989):

Caecal intussusceptions in horses and the significance of Anoplocephala perfoliata.
Vet. Rec., 124, 34-37

PANDEY, V.S.; OUHELLI, H.; ELKHALFANE, A. (1981):

Epidemiological observations on stomach worms of horses in Morocco.
J. Helminth., 55, 155-160

PANKAVICH, J.A.; BERGER, H.; SIMKINS, K.L. (1992):

Efficacy of moxidectin, nemadectin and ivermectin against an ivermectin-resistant strain of Haemonchus contortus in sheep.

Vet. Rec., 130, 241-243

PATTON, S.; McCACKEN, M.D. (1981):

The occurrence and effect of Thelazia in horses.
Equine Pract., 3, 53-57

PEARSON, G.R.; DAVIES, L.W.; WHITE, A.L.; O'BRIAN, J.K. (1993):

Pathological lesions associated with Anoplocephala perfoliata at the ileo-caecal junction of horses.

Vet. Rec., 132, 179-182

PECHEUR, M. (1988):

Anthelmintische Wirkung von Febantel + Trichlorfon gegen gastrointestinale Parasiten beim Pferd.

Vet.-med. Nachr., 59, S, 133-138

PEITGEN, U. (1993):

Vergleichende Untersuchungen zum Einfluß von Ivermectin, Pyrantelbonat und Cambendazol auf Strongylidenbefall bei Pferden unter Berücksichtigung von Behandlungs-intervall und Weidekontamination.

Vet. med. Diss., München

PFISTER, K.; HOFER, B.; CAUDERON, H.; VON TSCHARNER, C.; TSCHUDI, P. (1988):

Pathologische und pathophysiologische Befunde bei Aufzuchtfohlen mit massivem Befall durch kleine Strongyliden.

Abstr., 13. Tagung DGP e.V. (22.-25.3. Neuchatel / Schweiz), 58

PIETROBELLINI, M.; SCHIFF, D.; BATELLI, G.; GIORDANI, A. (1995):

Indagine sulle parassitosi intestinali degli equini in provincia di Udine.
Arch. vet. Ital., 46, 136

PÖLLEIN, W. (1989):

Natürliche Strongyliden-Infektionen bei Gestütspferden unter besonderer Berücksichtigung der Möglichkeiten und Grenzen serologischer Untersuchungen.

Vet. med. Diss., München

POMROY, W.E.; WHELAN, N.C. (1993):

*Efficacy of moxidectin against an ivermectin-resistant strain of *Ostertagia circumcinata* in young sheep.*

Vet. Rec., 132, 416

PRICHARD, R.K.; HALL, C.A.; KELLY, J.D.; MARTIN, I.C.A.; DONALD, A.D. (1980):

The problem of anthelmintic resistance in nematodes.

Aust. vet. J., 56, 239-251

PRINCIPATO, M. (1989):

*Observations on the occurrence of five species of *Gasterophilus* larvae in free-ranging horses in Umbria, Central Italy.*

Vet. Parasitol., 31, 173-177

PROUDMAN, C.J.; EDWARDS, G.B. (1992):

Validation of a centrifugation/flotation technique for the diagnosis of equine cestodiasis.

Vet. Rec., 131, 71-72

PROUDMAN, C.J.; EDWARDS, G.B. (1993):

Are tapeworms associated with equine colic? A case control study.

Equine vet. J., 25, 224-226

PROUDMAN, C.J.; SWAN, J.D.; TREES, A.J. (1995):

*Efficacy of pyrantel embonate and praziquantel against the equine tapeworm *Anoplocephala mamillana*.*

Vet. Rec., 137, 45-46

PROUDMAN, C.J.; TREES, A.J. (1996):

*Use of excretory/secretory antigens for the serodiagnosis of *Anoplocephala perfoliata* cestodosis.*

Vet. Parasitol., 61, 239-247

RAHMAN, S.A.; WADDELL, A.H. (1980):

Diet and feeding behavior of strongyles in the subfamily Cyathostominae Nicoll, 1972 (Nematoda: Strongylidea) from the horse.

Mys. J. Agric. Sci., 14, 91-102

RANJAN, S.; TRUDEAU, C.; PRICHARD, R. K.; DAIGNEAULT, J.; REW, R. S. (1997):

Nematode reinfection following treatment of cattle with doramectin and ivermectin.

Vet. Parasitol., 72, 25-31

REILLY, G.A.C.; CASSIDY, J.P.; TAYLOR, S.M. (1993):

Two fatal cases of diarrhoea in horses associated with larvae of the small strongyles.

Vet. Rec., 132, 267-268

REINEMEYER, C.R.; SMITH, S.A.; GABEL, A.A.; HERD, R.P. (1984):

The prevalence and intensity of internal parasites of horses in the U.S.A. .

Vet. Parasitol., 15, 75-83

REINEMEYER, C.R.; HERD, R.P. (1986):

Anatomic distribution of encysted cyathostome larvae in the horse.

Amer. J. vet. Res., 47, 510-513

REINEMEYER, C.R.; AGUILAR, R. (1992):

Dose titration study of moxidectin gel as an oral equine anthelmintic.

Proc. 37th Ann. Meet. Amer. Ass. Vet. Parasitol. (2.-4.8., Boston / Massachusetts), 72

REINEMEYER, C.R.; TINEO, A.A. (1993):

Comparison of the efficacies of moxidectin gel and ivermectin paste against nematode parasites of horses.

Proc. 38th Ann. Meet. Amer. Ass. Vet. Parasitol. (17.-20.7., Minneapolis / Minnesota), 35

RENDANO, V.T.; GEORGI, J.R.; WHITE, K.K.; SACK, W.O.; KING, J.M.; BIANCHI, D.G.; THEODORIDES, V.J. (1979):

Equine verminous arteritis. An arteriographic evaluation of the larvicidal activity of albendazole.

Equine vet. J., 11, 225-231

RIBBECK, R.; HEIDE, H.; SCHICHT, W.; HIEPE, T. (1983):

Beiträge zur Parasitenfauna der DDR. 7. Mitteilung. Untersuchungen zum Vorkommen von Gasterophilus-Larven (Diptera: Gasterophilidae) bei Pferden.

Angew. Parasitol. 25, 78-83

RIBBECK, R.; SCHUSSER, G.F.; HAUPT, W. (1997):

Zum Vorkommen der larvalen Cyathostominose beim Pferd.

Tierärztl. Umsch., 52, 5, 254-263

RICHTER, W.; DIETZ, O.; RIBBECK, R.; HIEPE, T. (1979):

Vorkommen und Behandlung von Sommerwunden bei Sportpferden.

Mh. Vet.-Med., 34, 663-666

ROLFE, P.F.; DAWSON, K.L. (1994):

The efficacy of a combination anthelmintic against oxicabendazole resistant small strongyles, large strongyles and ascarids.

Aust. vet. J., 71, 304-306

RUBINSTEIN, D.I.; HOHMANN, M.E. (1989):

Parasites and social behavior of Island feral horses.

Oikos, 55, 312-320

SAUNDERS, L.Z.; SHONE, D.K.; PHILIP, J.R.; BIRKHEAD, H.A. (1974):

The effect of methyl-5(6) butyl-2-benzimidazole carbamate (parbendazole) on the reproduction in sheep and other animals. Malformations in newborn lambs.

Cornell Vet., 64, 7-10

SCHAPER, R.; LIEBISCH, A. (1991):

Einfluß eines systemisch wirkenden Antiparasitikums (Ivermectin) auf die Dungfauna und den Dungabbau der Rinder bei Weidehaltung.

Tierärztl. Umsch., 46, 12-18

SCHATZMANN, U.; PFISTER, K.; SCHMITT, K. (1986):

Erfahrungen mit Ivermectin in einem größeren Pferdebestand.

Prakt. Tierarzt, 67, 649-651

SCHILLINGER, D. (1986):

Sommerwunden: Wunden die im Winter heilen.

Vet., 1, (5/6), 14-15

SCHILLINGER, D.; HASSLINGER, M.-A. (1994):

Benzimidazole resistance in small strongyles of horses - Occurrence in Germany and strategies for avoiding resistance.

Rev. méd. Vét., 145, 119-124

SCHLICHTING, C.K. (1981):

*Vorkommen, Bedeutung und Bekämpfung von Infektionen mit *Strongyloides westeri* Ihle 1917 (Strongyloididae) bei Fohlen.*

Vet. med. Diss., Hannover

SCHLICHTING, C.K.; STOYE, M. (1985):

*Zur Wirksamkeit von Ivermectin auf *Strongyloides westeri* bei Fohlen.*

Prakt. Tierarzt, 66, 128-130

SCHMIDT, G.M.; KREHBIEL, J.D.; COLEY, S.C.; LEID, W. (1982):

Equine Onchocerciasis: lesions in the nuchal ligament of midwestern U.S. horses.

Vet. Pathol., 19, 16-22

SCHUSTER, R. (1991):

Morphometrische Analyse einer Anoplocephala perfoliata-Population.

Angew. Parasitol., 32, 105-111

SEIBERT, B.P.; NEWCOMB, K.M.; MICHAEL, B.F. (1986):

Critical test and safety evaluations of an oral paste preparation of mebendazole and trichlorfon in horses.

Amer. J. vet. Res., 47, 1347-1350

SHARIR, B.; PIPANO, E.; MARKOVICS, A.; DANIELI, Y. (1987):

Field studies on gastrointestinal infestation in Israel horses.

Israel, J. vet. med., 43, 223-227

SHOOP, W.L. (1993):

Ivermectin resistance.

Parasitol. Today, 9, 154-159

SHOOP, W.L.; HAINES, H.W.; MICHAEL, B.F.; EARY, C.H. (1993):

Mutual resistance to avermectins and milbemycins: oral activity of ivermectin and moxidectin against ivermectin-resistant and susceptible nematodes.

Vet. Rec., 133, 445-447

SHOOP, W.L.; MROZIK, H.; FISHER, M.H. (1995):

Structure and activity of avermectins and milbemycins in animal health.

Vet. Parasitol., 59, 139-156

SLOCOMBE, J.O.D. (1979):

Prevalence and treatment of tapeworms in horses.

Canad. vet. J., 20, 136-140

SLOCOMBE, J.O.D. (1985):

Pathogenesis of helminths in equids.

Vet. Parasitol., 18, 139-153

SLOCOMBE, J.O.D.; SMART, J. (1975):

Evaluation of pyrantel pamoate against strongyles in horses.

Canad. vet. J., 16, 310-312

SLOCOMBE, J.O.D.; McCRAW, B.M.; PENNOCK, P.W.; VASEY, J. (1982):

*Effectiveness of ivermectin against later 4th-stage *Strongylus vulgaris* in ponies.*

Amer. J. vet. Res., 43, 1525-1529

SLOCOMBE, J.O.D.; McCRAW, B.M.; PENNOCK, P.W.; BAIRD, J.D. (1983):

*Effectiveness of fenbendazole against later 4th-stage *Strongylus vulgaris* in ponies.*

Amer. J. vet. Res., 44, 2285-2289

SLOCOMBE, J.O.D.; COTE, J.F. (1984):

Effectiveness of ivermectin paste for removal of nematodes in the horse.

Canad. vet. J., 25, 386-388

SLOCOMBE, J.O.D.; McCRAW, B.M. (1984):

*Evaluation of ivermectin against later 4th-stage *Strongylus vulgaris* in ponies at two and five weeks after treatment.*

Canad. J. comp. Med., 48, 343-348

SLOCOMBE, J.O.D.; McCRAW, B.M.; PENNOCK, P.W.; DUCHARME, N.G.;

BAIRD, J.D. (1986):

*Effectiveness of oxfendazole against early later 4th-stage *Strongylus vulgaris* in ponies.*

Canad. J. vet. Res., 47, 495-500

SLOCOMBE, J.O.D.; McCRAW, B.M.; PENNOCK, P.W.; DUCHARME, N.G.;

BAIRD, J.D. (1987):

Strongylus vulgaris in the tunica media of arteries of ponies and treatment with ivermectin.

Canad. J. vet. Res., 51, 232-235

SLOCOMBE, J.O.D.; COTE, J.F.; McMILLAN, I. (1989):

Effectiveness of oxibendazole against benzimidazole-resistant strongyles in horses.

Canad. vet. J., 30, 663-665

SMITH, H.J. (1976):

Strongyle infections in ponies. II. Reinfection of treated animals.

Canad. J. comp. Med., 40, 334-340

SMITH, H.J. (1978):

Experimental trichonema infections in mature ponies.

Vet. Parasitol., 4, 265-273

SOLARI BASANO, F.; GENCHI, C. (1996):

Efficacy of 2 % Moxidectin Equine gel given at different dose rates against nematodes in naturally infected horses.

Parassitologia, 38, 242

SOMMER, C.; GRØNVOLD, J.; HOLTER, P.; NANSEN, P. (1993):

*Effects of ivermectin on two afro tropical dung beetles, *Ontophagus gazella* and *Diastellopalpus quinquedens* (Coleoptera: Scarabaeidae).*

Vet. Parasitol., 48, 171-180

SOULSBY, E.J.L. (1965):

Textbook of Veterinary Clinical Parasitology Vol. I. Helminths

Oxford, Blackwell Scientific Publications

SREEKUMARAN, P.; RATHORE, U.S.; DHINSA, H.S.; SINGH, R.P. (1987):

Evaluation of morantel citrate („Banminth“ Pfister) against helminthic infection in equines.
J. Remount vet. Corps, 26, 115-120

SRIHAKIM, S.; SWERCZEK, T.W. (1978):

Pathologic changes and pathogenesis of Parascaris equorum infection in parasite-free pony foals.
Amer. J. vet. Res., 39, 1155-1160

STANDARD, A.A.; CELLO, R.M. (1975):

Onchocerca cervicalis infection in horses from the western United States.
Amer. J. vet. Res., 36, 1029-1031

STOYE, M. (1978):

Planmäßige Parasitenbekämpfung.
6. Arbeitstagung der Fachgruppe Pferdekrankheiten der DVG, (28.-30.10., Berlin / BRD),
61-70

STOYE, M. (1986):

Vorkommen, Bedeutung und Bekämpfung von Strongyloides westeri beim Pferd.
Vortragsmanuskript Ivomec®-P Symposion, Parasitosen des Pferdes (14.5., Frankfurt/M.)

STOYE, M. (1991):

Die Endoparasiten des Pferdes.
Tierärztl. Umsch., 46, 429-430

STRONG, L. (1993):

Overview: The impact of avermectins on pastureland ecology.
Vet. Parasitol., 48, 3-17

STRONG, L.; WALL, R. (1994):

Effects of ivermectin and moxidectin on the insects of cattle dung.
Bull. ent. Res., 84, 403-409

STRONG, L.; WALL, R.; WOOLFORD, A.; DJEDDOUR, D. (1996):

The effect of faecally excreted ivermectin and fenbendazole on the insect colonisation of cattle dung following the oral administration of sustained-release boluses.
Vet. Parasitol., 62, 253-266

TAUSEND, S. (1989):

Feldversuche über Zestodeninfektionen bei Pferden und ihre Kontrolle.
Vet. med. Diss., München

TAYLOR, M.A.; HUNT, K.R. (1989):

Anthelmintic drug resistance in the U.K. .
Vet. Rec., 25, 143-147

TAYLOR, S.M.; KENNY, J. (1995):

Comparison of moxidectin with ivermectin and pyrantel embonate for reduction of faecal egg counts in horses.

Vet. Rec., 137, 516-518

TOLOSA, J.; VÁZQUEZ, M.; CHIARETTA, A.; SBAFFO, A.; SÁNCHEZ, J.; GUARDA, I.; FERRARIS, G.; ESPINOSA, D.; MUÑOZ COBENÁS, M.; BULMAN, G.M.; AMBRUSTOLO, R.R. (1994):

Efficacia comparativa de moxidectin gel e ivermectina pasta contra Gasterophilus spp. en equinos.

Proc. VII. Congreso Argentino de Ciencias Veterinarias (8.-11.11., Buenos Aires / Argentina), 218

TOLLIVER, S.C.; LYONS, E.T.; DRUDGE, J.H. (1987):

Prevalence of internal parasites in horses in critical tests of activity of parasiticides over a 28-year period (1956-1983) in Kentucky.

Vet. Parasitol., 23, 273-284

TOLLIVER, S.C.; LYONS, E.T.; DRUDGE, J.H.; STAMPER, S.; GRANSTORM, D.E. (1993):

Critical tests of thiabendazole, oxbendazole and oxfendazole for drug resistance of population-B equine small strongyles (1989 and 1990).

Amer. J. vet. Res., 54, 908-913

TOUTAIN, P.L.; UPSON, D.W.; TERHUNE, T.N.; McKENZIE, M.E. (1997):

Comparative pharmacokinetics of doramectin and ivermectin in cattle.

Vet. Parasitol., 72, 3-8

TRAEDER, W. (1994):

Das pharmakologische Verhalten von Doramectin, einem neuen makrocyclischen Laktonderivat aus der Gruppe der Avermectine.

Tierärztl. Umsch., 49, 465-469

TURNER, M.J.; SCHAEFFER, J.M. (1989):

Mode of action of ivermectin.

In: W.C. CAMPELL, Ivermectin and Abamectin. Springer Verlag, New York, 73-78

UHLINGER, C. (1990):

Effects of three anthelmintic schedules on the incidence of colic in horses.

Equine vet. J., 22, 251-254

UHLINGER, C.; JOHNSTONE, C. (1984):

Failure to reestablish benzimidazole susceptible populations of small strongyles after prolonged treatment with non-benzimidazole drugs.

J. Equine vet. Sci., 4, 7-9

UHLINGER, C.; KRISTULA, M. (1992):

Effects of alternation of drug classes on the development of oxicabendazole resistance in a herd of horses.

J. Amer. vet. med. Ass., 201, 51-55

ULLRICH, D. (1987):

Verbreitung benzimidazol-resistenter Strongyliden in Nordrhein-Westfalen.

Vet. med. Diss., Hannover

VALDEZ, R.A.; DI PIETRO, J.A.; PAUL, A.J.; LOCK, T.F.; HUNGERFORD, L.L.;

TODD, K.S. (1995):

Controlled efficacy study of the bioequivalence of Strongid® C and generic pyrantel tartrate in horses.

Vet. Parasitol., 60, 83-102

VAN LOON, G.; DEPREZ, P.; MUYLLE, E.; SUSTRONCK, B. (1995):

Larval cyathostomiasis as a cause of death in two regularly dewormed horses.

J. Vet. Med. A., 42, 301-306

VASEY, J.R. (1981):

Equine cutaneous habronemiasis.

Comp. Cont. Educ. Pract. Vet., 3, 290-298

VERCRUYSSE, J.; EYSKER, M.; DEMEULENAERE, D.; SMETS, K.; DORNY, P. (1998):

Persistence of the efficacy of a moxidectin gel on the establishment of cyathostominae in horses.

Vet. Rec., 143, 307-309

VERBERNE, L.R.M.; MIRCK, M.H. (1975):

Over de toxiciteit van parbendazole (Helmatac® 30) voor paarden en pony's.

T. Diergeneesk., 100, 1143-1153

WALDE, I. (1986):

Differentialdiagnostische und therapeutische Aspekte bei der „Mondblindheit“ des Pferdes.

Pferdeheilk., 2, 67-78

WEBSTER, J.H.; BAIRD, J.D.; GUNAWAN, M.; MARTIN, I.C.A.; KELLY, J.D. (1981):

Resistance to benzimidazole anthelmintics in equine strongyles. II. Evidence of sideroresistance, and susceptibility of benzimidazole-resistant strongyles to non-benzimidazole compounds.

Aust. vet. J., 57, 172-181

WEILAND, G.; HASSLINGER, M.-A.; MEZGER, S.; PÖLLEIN, W. (1991):

Möglichkeiten und Grenzen der Immundiagnostik des Strongylidenbefalles beim Pferd.

Berl. Münch. tierärztl. Wschr., 104, 149-153

WEJDA, E. (1961):

Dilatatio et stenosis oesophagi bei einem Fohlen (hervorgerufen durch wandernde Gasterophiluslarven).

Mh. Vet.-Med., 16, 317-318

WELBERS, N. (1981):

Eine orientierende Felduntersuchung in Norddeutschland zur Epizootologie der Strongylideninfektion des Pferdes.

Vet. med. Diss., Hannover

WESCOTT, R.B. (1987):

Anthelmintics for horses.

Int. J. Parasitol., 17, 503-510

WESCOTT, R.B.; JEN, L.W.; HELLIER, L.E.; SCHAEFFER, D.C.; TRAUB, J.L. (1985):

Treating benzimidazole-resistant small strongyles (cyathostominae) in horses.

Vet. Med., 80, 60-64

WETZEL, R. (1951):

Verbesserte McMaster-Kammer zum Auszählen von Wurmeiern.

Tierärztl. Umschau, 6, 209-210

WICKS, S.R.; KAYE, B.; WEATHERLEY, A.J.; LEWIS, D.; DAVISON, E.; GIBSON, S.P.; SMITH, D.G. (1993):

Effect of formulation on the pharmacokinetics and efficacy of doramectin.

Vet. Parasitol., 49, 17-26

WILCOX, B.P. (1985):

The eye and ear.

In: K.V.F. JUPP, P.C. KENNEDY and N. PALMER, *Pathology of domestic animals*, Vol. 1, Academic Press, Orlando, 340-406

WILLIAMSSON, R.M.C.; GASSER, R.B.; MIDDLETON, D.; BEVERIDGE, I. (1997):

The distribution of Anoplocephala perfoliata in the intestine of the horse and associated pathological changes.

Vet. Parasitol., 73, 225-241

XIAO, L.; HERD, R.P.; MAJEWSKI, G.A. (1994):

Comparative efficacy of moxidectin and ivermectin against hypobiotic and encysted cyathostomes and other equine parasites.

Vet. Parasitol., 53, 83-90

YAGER, J.A.; SCOTT, D.W. (1985):

The skin and appendages.

In: K.V.F. JUPP, P.C. KENNEDY an N. PALMER, *Pathology of domestic animals*, Vol. 1,
Academic Press, Orlando, 407-549

YOSHIHARA, T.; OIKAWA, M.; KATAYAMA, Y.; KANEKO, M. (1993):

The prevalence of internal parasites in race horses in Japan: A necropsy survey.

Proc. 14th Int. Conf. WAAVP (8.-13.8., Cambridge / UK), 151

YOSHIHARA, T.; TOGUCHI, M.; KOMAZAWA, H.; OHWA, Y. (1995):

Efficacy of bithionol paste against Anoplocephala perfoliata in naturally infected horses.

Abstr. 15th Int. Conf. WAAVP (30.8.-2.9., Yokohama / Japan), 129

ZEEUW, G. (1997):

Vergleichende Untersuchungen über Effekt und Wirkungsdauer von Moxidectin, Ivermectin und Pyrantelbonat beim Pferd unter Gestütsbedingungen.

Vet. med. Diss., München

ZEEUW, G.; HASSLINGER, M.-A. (1997):

Vergleichende Untersuchungen zu Eizahlreduktion und Behandlungsintervall bei Endoparasiten des Pferdes.

Prakt. Tierarzt, 78, 10, 857-864

ZULALIAN, J.; STOUT, S.J.; DaCUNHA, A.R.; GARCES, T.; MILLER, P. (1994):

Absorption, tissue distribution, excretion and metabolism of moxidectin in cattle.

J. Agric. Food Chem., 42, 381-387

