

---

## 8 Literaturverzeichnis

- Alton, E.W., Stern, M., Farley, R., Jaffe, A., Chadwick, S.L., Phillips, J., Davies, J., Smith, S.N., Browning, J., Davies, M.G., Hodson, M.E., Durham, S.R., Li, D., Jeffery, P.K., Scallan, M., Balfour, R., Eastman, S.J., Cheng, S.H., Smith, A.E., Meeker, D. & Geddes, D.M. (1999). Cationic lipid-mediated CFTR gene transfer to the lungs and nose of patients with cystic fibrosis: a double-blind placebo-controlled trial. *Lancet*, **353**, 947-54.
- Anwer, K., Shi, M., French, M.F., Muller, S.R., Chen, W., Liu, Q., Proctor, B.L., Wang, J., Mumper, R.J., Singhal, A., Rolland, A.P. & Alila, H.W. (1998). Systemic effect of human growth hormone after intramuscular injection of a single dose of a muscle-specific gene medicine. *Hum Gene Ther*, **9**, 659-70.
- Behr, J. (1997). The proton sponge: A trick to enter cells the viruses did not exploit. *Chimia*, **51**, 34-36.
- Blessing, T., Kursa, M., Holzhauser, R., Kircheis, R. & Wagner, E. (2001). Different strategies for formation of pegylated EGF-conjugated PEI/DNA complexes for targeted gene delivery. *Bioconjug Chem*, **12**, 529-37.
- Boletta, A., Benigni, A., Lutz, J., Remuzzi, G., Soria, M.R. & Monaco, L. (1997). Nonviral gene delivery to the rat kidney with polyethylenimine. *Hum Gene Ther*, **8**, 1243-51.
- Boussif, O., Lezoualc'h, F., Zanta, M.A., Mergny, M.D., Scherman, D., Demeneix, B. & Behr, J.P. (1995). A versatile vector for gene and oligonucleotide transfer into cells in culture and in vivo: polyethylenimine. *Proc Natl Acad Sci U S A*, **92**, 7297-301.
- Dragonzi, A., Boletta, A., Biffi, A., Muggia, A., Sersale, G., Cheng, S.H., Bordignon, C., Assael, B.M. & Conese, M. (1999). Comparison between cationic polymers and lipids in mediating systemic gene delivery to the lungs. *Gene Ther*, **6**, 1995-2004.
- Dragonzi, A., Dina, G., Villa, A., Calori, G., Biffi, A., Bordignon, C., Assael, B.M. & Conese, M. (2000). Biodistribution and transgene expression with nonviral cationic vector/DNA complexes in the lungs. *Gene Ther*, **7**, 1753-60.
- Branden, L.J., Mohamed, A.J. & Smith, C.I. (1999). A peptide nucleic acid-nuclear localization signal fusion that mediates nuclear transport of DNA. *Nat Biotechnol*, **17**, 784-7.
- Capecchi, M.R. (1980). High efficiency transformation by direct microinjection of DNA into cultured mammalian cells. *Cell*, **22**, 479-88.
- Caplen, N.J., Alton, E.W., Middleton, P.G., Dorin, J.R., Stevenson, B.J., Gao, X., Durham, S.R., Jeffery, P.K., Hodson, M.E., Coutelle, C. & et al. (1995).

- 
- Liposome-mediated CFTR gene transfer to the nasal epithelium of patients with cystic fibrosis. *Nat Med*, **1**, 39-46.
- Chan, C.K. & Jans, D.A. (1999). Enhancement of polylysine-mediated transferrinfection by nuclear localization sequences: polylysine does not function as a nuclear localization sequence. *Hum Gene Ther*, **10**, 1695-702.
- Chemin, I., Moradpour, D., Wieland, S., Offensperger, W.B., Walter, E., Behr, J.P. & Blum, H.E. (1998). Liver-directed gene transfer: a linear polyethylenimine derivative mediates highly efficient DNA delivery to primary hepatocytes in vitro and in vivo. *J Viral Hepat*, **5**, 369-75.
- Ciolina, C., Byk, G., Blanche, F., Thuillier, V., Scherman, D. & Wils, P. (1999). Coupling of nuclear localization signals to plasmid DNA and specific interaction of the conjugates with importin alpha. *Bioconjug Chem*, **10**, 49-55.
- Coleman, M.E., DeMayo, F., Yin, K.C., Lee, H.M., Geske, R., Montgomery, C. & Schwartz, R.J. (1995). Myogenic vector expression of insulin-like growth factor I stimulates muscle cell differentiation and myofiber hypertrophy in transgenic mice. *J Biol Chem*, **270**, 12109-16.
- Colin, M., Moritz, S., Schneider, H., Capeau, J., Coutelle, C. & Brahimi-Horn, M.C. (2000). Haemoglobin interferes with the ex vivo luciferase luminescence assay: consequence for detection of luciferase reporter gene expression in vivo. *Gene Ther*, **7**, 1333-6.
- Coll, J.L., Chollet, P., Brambilla, E., Desplanques, D., Behr, J.P. & Favrot, M. (1999). In vivo delivery to tumors of DNA complexed with linear polyethylenimine. *Hum Gene Ther*, **10**, 1659-66.
- Cotten, M., Langle-Rouault, F., Kirlappos, H., Wagner, E., Mechtler, K., Zenke, M., Beug, H. & Birnstiel, M.L. (1990). Transferrin-polycation-mediated introduction of DNA into human leukemic cells: stimulation by agents that affect the survival of transfected DNA or modulate transferrin receptor levels. *Proc Natl Acad Sci U S A*, **87**, 4033-7.
- Dash, P.R., Read, M.L., Barrett, L.B., Wolfert, M.A. & Seymour, L.W. (1999). Factors affecting blood clearance and in vivo distribution of polyelectrolyte complexes for gene delivery. *Gene Ther*, **6**, 643-50.
- de Lima, M.C., Simoes, S., Pires, P., Gaspar, R., Slepushkin, V. & Duzgunes, N. (1999). Gene delivery mediated by cationic liposomes: from biophysical aspects to enhancement of transfection. *Mol Membr Biol*, **16**, 103-9.
- Densmore, C.L., Orson, F.M., Xu, B., Kinsey, B.M., Waldrep, J.C., Hua, P., Bhogal, B. & Knight, V. (2000). Aerosol delivery of robust polyethyleneimine-DNA complexes for gene therapy and genetic immunization. *Mol Ther*, **1**, 180-8.
- Eguchi, A., Akuta, T., Okuyama, H., Senda, T., Yokoi, H., Inokuchi, H., Fujita, S., Hayakawa, T., Takeda, K., Hasegawa, M. & Nakanishi, M. (2001). Protein

- 
- transduction domain of HIV-1 Tat protein promotes efficient delivery of DNA into mammalian cells. *J Biol Chem*, **276**, 26204-10.
- Erbacher, P., Bettinger, T., Belguise-Valladier, P., Zou, S., Coll, J.L., Behr, J.P. & Remy, J.S. (1999a). Transfection and physical properties of various saccharide, poly(ethylene glycol), and antibody-derivatized polyethylenimines (PEI). *J Gene Med*, **1**, 210-22.
- Erbacher, P., Remy, J.S. & Behr, J.P. (1999b). Gene transfer with synthetic virus-like particles via the integrin-mediated endocytosis pathway. *Gene Ther*, **6**, 138-45.
- Ernst, N., Ulrichskotter, S., Schmalix, W.A., Radler, J., Galneder, R., Mayer, E., Gersting, S., Plank, C., Reinhardt, D. & Rosenecker, J. (1999). Interaction of liposomal and polycationic transfection complexes with pulmonary surfactant. *J Gene Med*, **1**, 331-40.
- Fawell, S., Seery, J., Daikh, Y., Moore, C., Chen, L.L., Pepinsky, B. & Barsoum, J. (1994). Tat-mediated delivery of heterologous proteins into cells. *Proc Natl Acad Sci U S A*, **91**, 664-8.
- Felgner, P.L., Gadek, T.R., Holm, M., Roman, R., Chan, H.W., Wenz, M., Northrop, J.P., Ringold, G.M. & Danielsen, M. (1987). Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure. *Proc Natl Acad Sci U S A*, **84**, 7413-7.
- Ferrari, S., Moro, E., Pettenazzo, A., Behr, J.P., Zucchello, F. & Scarpa, M. (1997). ExGen 500 is an efficient vector for gene delivery to lung epithelial cells in vitro and in vivo. *Gene Ther*, **4**, 1100-6.
- Ferrari, S., Pettenazzo, A., Garbati, N., Zucchello, F., Behr, J.P. & Scarpa, M. (1999). Polyethylenimine shows properties of interest for cystic fibrosis gene therapy. *Biochim Biophys Acta*, **1447**, 219-25.
- Finsinger, D., Remy, J.S., Erbacher, P., Koch, C. & Plank, C. (2000). Protective copolymers for nonviral gene vectors: synthesis, vector characterization and application in gene delivery. *Gene Therapy*, **7**, 1183-1192.
- Frankel, A.D. & Pabo, C.O. (1988). Cellular uptake of the tat protein from human immunodeficiency virus. *Cell*, **55**, 1189-93.
- Futaki, S., Suzuki, T., Ohashi, W., Yagami, T., Tanaka, S., Ueda, K. & Sugiura, Y. (2001). Arginine-rich peptides. An abundant source of membrane-permeable peptides having potential as carriers for intracellular protein delivery. *J Biol Chem*, **276**, 5836-40.
- Gautam, A., Densmore, C.L. & Waldrep, J.C. (2000). Inhibition of experimental lung metastasis by aerosol delivery of PEI-p53 complexes. *Mol Ther*, **2**, 318-23.
- Gessner, A (2001) Untersuchungen zur Proteinadsorption auf kolloidalen Modellpartikeln und Arzneistoffträgern zur parenteralen Anwendung. Dissertation. Freie Universität Berlin.

- 
- Gill, D.R., Southern, K.W., Mofford, K.A., Seddon, T., Huang, L., Sorgi, F., Thomson, A., MacVinish, L.J., Ratcliff, R., Bilton, D., Lane, D.J., Littlewood, J.M., Webb, A.K., Middleton, P.G., Colledge, W.H., Cuthbert, A.W., Evans, M.J., Higgins, C.F. & Hyde, S.C. (1997). A placebo-controlled study of liposome-mediated gene transfer to the nasal epithelium of patients with cystic fibrosis. *Gene Ther*, **4**, 199-209.
- Godbey, W.T., Wu, K.K. & Mikos, A.G. (1999). Tracking the intracellular path of poly(ethylenimine)/DNA complexes for gene delivery. *Proc Natl Acad Sci U S A*, **96**, 5177-81.
- Gorlich, D. & Kutay, U. (1999). Transport between the cell nucleus and the cytoplasm. *Annu Rev Cell Dev Biol*, **15**, 607-60.
- Gossen, M., Freundlieb, S., Bender, G., Muller, G., Hillen, W. & Bujard, H. (1995). Transcriptional activation by tetracyclines in mammalian cells. *Science*, **268**, 1766-9.
- Gottschalk, S., Sparrow, J.T., Hauer, J., Mims, M.P., Leland, F.E., Woo, S.L. & Smith, L.C. (1996). A novel DNA-peptide complex for efficient gene transfer and expression in mammalian cells. *Gene Ther*, **3**, 48-57.
- Goula, D., Benoist, C., Mantero, S., Merlo, G., Levi, G. & Demeneix, B.A. (1998a). Polyethylenimine-based intravenous delivery of transgenes to mouse lung. *Gene Ther*, **5**, 1291-5.
- Goula, D., Remy, J.S., Erbacher, P., Wasowicz, M., Levi, G., Abdallah, B. & Demeneix, B.A. (1998b). Size, diffusibility and transfection performance of linear PEI/DNA complexes in the mouse central nervous system. *Gene Ther*, **5**, 712-7.
- Hagstrom, J.E., Ludtke, J.J., Bassik, M.C., Sebestyen, M.G., Adam, S.A. & Wolff, J.A. (1997). Nuclear import of DNA in digitonin-permeabilized cells. *J Cell Sci*, **110**, 2323-31.
- Harris, C.E., Agarwal, S., Hu, P., Wagner, E. & Curiel, D.T. (1993). Receptor-mediated gene transfer to airway epithelial cells in primary culture. *Am J Respir Cell Mol Biol*, **9**, 441-7.
- Hirata, T., Bitterman, P.B., Mornex, J.F. & Crystal, R.G. (1986). Expression of the transferrin receptor gene during the process of mononuclear phagocyte maturation. *J Immunol*, **136**, 1339-45.
- Huang, M.T. & Gorman, C.M. (1990). Intervening sequences increase efficiency of RNA 3' processing and accumulation of cytoplasmic RNA. *Nucleic Acids Res*, **18**, 937-47.
- James, M.B. & Giorgio, T.D. (2000). Nuclear-associated plasmid, but not cell-associated plasmid, is correlated with transgene expression in cultured mammalian cells. *Mol Ther*, **1**, 339-46.

- 
- Jans, D.A., Chan, C.K. & Huebner, S. (1998). Signals mediating nuclear targeting and their regulation: application in drug delivery. *Med Res Rev*, **18**, 189-223.
- Katkin, J.P., Husser, R.C., Langston, C. & Welty, S.E. (1997). Exogenous surfactant enhances the delivery of recombinant adenoviral vectors to the lung. *Hum Gene Ther*, **8**, 171-6.
- Keminer, O. & Peters, R. (1999). Permeability of single nuclear pores. *Biophys J*, **77**, 217-28.
- Kichler, A., Leborgne, C., Coeytaux, E. & Danos, O. (2001). Polyethylenimine-mediated gene delivery: a mechanistic study. *J Gene Med*, **3**, 135-44.
- Kircheis, R., Schuller, S., Brunner, S., Ogris, M., Heider, K.H., Zauner, W. & Wagner, E. (1999). Polycation-based DNA complexes for tumor-targeted gene delivery in vivo. *J Gene Med*, **1**, 111-20.
- Kircheis, R., Wightman, L., Schreiber, A., Robitz, B., Rossler, V., Kursa, M. & Wagner, E. (2001). Polyethylenimine/DNA complexes shielded by transferrin target gene expression to tumors after systemic application. *Gene Therapy*, **8**, 28-40.
- Kozak, M. (1991). Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J Biol Chem*, **266**, 19867-70.
- Kozak, M. (1992). Regulation of translation in eukaryotic systems. *Annu Rev Cell Biol*, **8**, 197-225.
- Kren, B.T., Parashar, B., Bandyopadhyay, P., Chowdhury, N.R., Chowdhury, J.R. & Steer, C.J. (1999). Correction of the UDP-glucuronosyltransferase gene defect in the gunn rat model of crigler-najjar syndrome type I with a chimeric oligonucleotide. *Proc Natl Acad Sci U S A*, **96**, 10349-54.
- Lee, E.R., Marshall, J., Siegel, C.S., Jiang, C., Yew, N.S., Nichols, M.R., Nietupski, J.B., Ziegler, R.J., Lane, M.B., Wang, K.X., Wan, N.C., Scheule, R.K., Harris, D.J., Smith, A.E. & Cheng, S.H. (1996). Detailed analysis of structures and formulations of cationic lipids for efficient gene transfer to the lung. *Hum Gene Ther*, **7**, 1701-17.
- Lee, K.D., Nir, S. & Papahadjopoulos, D. (1993). Quantitative analysis of liposome-cell interactions in vitro: rate constants of binding and endocytosis with suspension and adherent J774 cells and human monocytes. *Biochemistry*, **32**, 889-99.
- Lewin, M., Carlesso, N., Tung, C.H., Tang, X.W., Cory, D., Scadden, D.T. & Weissleder, R. (2000). Tat peptide-derivatized magnetic nanoparticles allow in vivo tracking and recovery of progenitor cells. *Nat Biotechnol*, **18**, 410-4.
- Li, S., Tan, Y., Viroonchatapan, E., Pitt, B.R. & Huang, L. (2000). Targeted gene delivery to pulmonary endothelium by anti-PECAM antibody. *Am J Physiol Lung Cell Mol Physiol*, **278**, L504-11.

- 
- Liu, F., Song, Y. & Liu, D. (1999). Hydrodynamics-based transfection in animals by systemic administration of plasmid DNA. *Gene Ther*, **6**, 1258-66.
- Mann, D.A. & Frankel, A.D. (1991). Endocytosis and targeting of exogenous HIV-1 Tat protein. *Embo J*, **10**, 1733-9.
- McGowan, S.E., Parenti, C.M., Hoidal, J.R. & Niewoehner, D.E. (1984). Ascorbic acid content and accumulation by alveolar macrophages from cigarette smokers and nonsmokers. *J Lab Clin Med*, **104**, 127-34.
- Mi, Z., Mai, J., Lu, X. & Robbins, P.D. (2000). Characterization of a class of cationic peptides able to facilitate efficient protein transduction in vitro and in vivo. *Mol Ther*, **2**, 339-47.
- Mislick, K.A. & Baldeschwieler, J.D. (1996). Evidence for the role of proteoglycans in cation-mediated gene transfer. *Proc Natl Acad Sci U S A*, **93**, 12349-54.
- Morris, M.C., Chaloin, L., Mery, J., Heitz, F. & Divita, G. (1999). A novel potent strategy for gene delivery using a single peptide vector as a carrier. *Nucleic Acids Res*, **27**, 3510-7.
- Mounkes, L.C., Zhong, W., Cipres-Palacin, G., Heath, T.D. & Debs, R.J. (1998). Proteoglycans mediate cationic liposome-DNA complex-based gene delivery in vitro and in vivo. *J Biol Chem*, **273**, 26164-70.
- Nagahara, H., Vocero-Akbani, A.M., Snyder, E.L., Ho, A., Latham, D.G., Lissy, N.A., Becker-Hapak, M., Ezhevsky, S.A. & Dowdy, S.F. (1998). Transduction of full-length TAT fusion proteins into mammalian cells: TAT-p27Kip1 induces cell migration. *Nat Med*, **4**, 1449-52.
- Neves, C., Escriou, V., Byk, G., Scherman, D. & Wils, P. (1999). Intracellular fate and nuclear targeting of plasmid DNA. *Cell Biol Toxicol*, **15**, 193-202.
- Ogris, M., Brunner, S., Schuller, S., Kircheis, R. & Wagner, E. (1999). PEGylated DNA/transferrin-PEI complexes: reduced interaction with blood components, extended circulation in blood and potential for systemic gene delivery. *Gene Ther*, **6**, 595-605.
- Ogris, M., Steinlein, P., Carotta, S., Brunner, S. & Wagner, E. (2001). DNA/Polyethylenimine transfection particles: Influence of ligands, polymer size, and PEGylation on internalization and gene expression. *AAPS PharmSci* (<http://www.pharmsci.org>), **3**.
- Ogris, M., Steinlein, P., Kursa, M., Mechtler, K., Kircheis, R. & Wagner, E. (1998). The size of DNA/transferrin-PEI complexes is an important factor for gene expression in cultured cells. *Gene Ther*, **5**, 1425-33.
- Plank, C., Mechtler, K., Szoka, F.C., Jr. & Wagner, E. (1996). Activation of the complement system by synthetic DNA complexes: a potential barrier for intravenous gene delivery. *Hum Gene Ther*, **7**, 1437-46.

- 
- Plank, C., Oberhauser, B., Mechtler, K., Koch, C. & Wagner, E. (1994). The influence of endosome-disruptive peptides on gene transfer using synthetic virus-like gene transfer systems. *J Biol Chem*, **269**, 12918-24.
- Plank, C., Tang, M.X., Wolfe, A.R. & Szoka, F.C., Jr. (1999). Branched cationic peptides for gene delivery: role of type and number of cationic residues in formation and in vitro activity of DNA polyplexes. *Hum Gene Ther*, **10**, 319-32.
- Pollard, H., Remy, J.S., Loussouarn, G., Demolombe, S., Behr, J.P. & Escande, D. (1998). Polyethylenimine but not cationic lipids promotes transgene delivery to the nucleus in mammalian cells. *J Biol Chem*, **273**, 7507-11.
- Porteous, D.J., Dorin, J.R., McLachlan, G., Davidson-Smith, H., Davidson, H., Stevenson, B.J., Carothers, A.D., Wallace, W.A., Moralee, S., Hoenes, C., Kallmeyer, G., Michaelis, U., Naujoks, K., Ho, L.P., Samways, J.M., Imrie, M., Greening, A.P. & Innes, J.A. (1997). Evidence for safety and efficacy of DOTAP cationic liposome mediated CFTR gene transfer to the nasal epithelium of patients with cystic fibrosis. *Gene Ther*, **4**, 210-8.
- Pouton, C.W., Lucas, P., Thomas, B.J., Uduehi, A.N., Milroy, D.A. & Moss, S.H. (1998). Polycation-DNA complexes for gene delivery: a comparison of the biopharmaceutical properties of cationic polypeptides and cationic lipids. *J Control Release*, **53**, 289-99.
- Qin, L., Ding, Y., Pahud, D.R., Chang, E., Imperiale, M.J. & Bromberg, J.S. (1997). Promoter attenuation in gene therapy: interferon-gamma and tumor necrosis factor-alpha inhibit transgene expression. *Hum Gene Ther*, **8**, 2019-29.
- Raczka, E., Kukowska-Latallo, J.F., Rymaszewski, M., Chen, C. & Baker, J.R., Jr. (1998). The effect of synthetic surfactant Exosurf on gene transfer in mouse lung in vivo. *Gene Ther*, **5**, 1333-9.
- Remy, J.S., Abdallah, B., Zanta, M.A., Boussif, O., Behr, J.P. & Demeneix, B. (1998). GENE TRANSFER WITH LIPOSERMINES AND POLYETHYLENIMINES [Review]. *Advanced Drug Delivery Reviews*, **30**, 85-95.
- Rols, M.P., Delteil, C., Golzio, M., Dumond, P., Cros, S. & Teissie, J. (1998). In vivo electrically mediated protein and gene transfer in murine melanoma. *Nat Biotechnol*, **16**, 168-71.
- Rudolph, C., Lausier, J., Naundorf, S., Muller, R.H. & Rosenecker, J. (2000). In vivo gene delivery to the lung using polyethylenimine and fractured polyamidoamine dendrimers. *J Gene Med*, **2**, 269-78.
- Sanders, N.N., Van Rompaey, E., De Smedt, S.C. & Demeester, J. (2001). Structural alterations of gene complexes by cystic fibrosis sputum. *Am J Respir Crit Care Med*, **164**, 486-93.

- 
- Schaffer, D.V., Fidelman, N.A., Dan, N. & Lauffenburger, D.A. (2000). Vector unpacking as a potential barrier for receptor-mediated polyplex gene delivery. *Biotechnol Bioeng*, **67**, 598-606.
- Schwarze, S.R., Ho, A., Vocero-Akbani, A. & Dowdy, S.F. (1999). In vivo protein transduction: delivery of a biologically active protein into the mouse. *Science*, **285**, 1569-72.
- Sebestyen, M.G., Lutkue, J.J., Bassik, M.C., Zhang, G., Budker, V., Lukhtanov, E.A., Hagstrom, J.E. & Wolff, J.A. (1998). DNA vector chemistry: the covalent attachment of signal peptides to plasmid DNA. *Nat Biotechnol*, **16**, 80-5.
- Singh, D., Bisland, S.K., Kawamura, K. & Gariepy, J. (1999). Peptide-based intracellular shuttle able to facilitate gene transfer in mammalian cells. *Bioconjug Chem*, **10**, 745-54.
- Slepushkin, V.A., Simoes, S., Dazin, P., Newman, M.S., Guo, L.S., Pedrosa de Lima, M.C. & Duzgunes, N. (1997). Sterically stabilized pH-sensitive liposomes. Intracellular delivery of aqueous contents and prolonged circulation in vivo. *J Biol Chem*, **272**, 2382-8.
- Sperinde, J.J., Choi, S.J. & Szoka, F.C., Jr. (2001). Phage display selection of a peptide DNase II inhibitor that enhances gene delivery. *J Gene Med*, **3**, 101-8.
- Subramanian, A., Ranganathan, P. & Diamond, S.L. (1999). Nuclear targeting peptide scaffolds for lipofection of nondividing mammalian cells. *Nat Biotechnol*, **17**, 873-7.
- Torchilin, V.P., Rammohan, R., Weissig, V. & Levchenko, T.S. (2001). TAT peptide on the surface of liposomes affords their efficient intracellular delivery even at low temperature and in the presence of metabolic inhibitors. *Proc Natl Acad Sci USA*, **98**, 8786-91.
- Truant, R. & Cullen, B.R. (1999). The arginine-rich domains present in human immunodeficiency virus type 1 Tat and Rev function as direct importin beta-dependent nuclear localization signals. *Mol Cell Biol*, **19**, 1210-7.
- Tseng, W.C., Haselton, F.R. & Giorgio, T.D. (1997). Transfection by cationic liposomes using simultaneous single cell measurements of plasmid delivery and transgene expression. *J Biol Chem*, **272**, 25641-7.
- Ulmer, J.B., Donnelly, J.J., Parker, S.E., Rhodes, G.H., Felgner, P.L., Dwarki, V.J., Gromkowski, S.H., Deck, R.R., DeWitt, C.M., Friedman, A. & et al. (1993). Heterologous protection against influenza by injection of DNA encoding a viral protein. *Science*, **259**, 1745-9.
- Vives, E., Brodin, P. & Lebleu, B. (1997). A truncated HIV-1 Tat protein basic domain rapidly translocates through the plasma membrane and accumulates in the cell nucleus. *J Biol Chem*, **272**, 16010-7.

- 
- Wadhwa, M.S., Collard, W.T., Adami, R.C., McKenzie, D.L. & Rice, K.G. (1997). Peptide-mediated gene delivery: influence of peptide structure on gene expression. *Bioconjug Chem*, **8**, 81-8.
- Wang, Y., O'Malley, B.W. & Tsai, S.Y. (1997). Inducible system designed for future gene therapy. *Methods Mol Biol*, **63**, 401-13.
- Ward, C.M., Read, M.L. & Seymour, L.W. (2001). Systemic circulation of poly(L-lysine)/DNA vectors is influenced by polycation molecular weight and type of DNA: differential circulation in mice and rats and the implications for human gene therapy. *Blood*, **97**, 2221-9.
- Wheeler, C.J., Felgner, P.L., Tsai, Y.J., Marshall, J., Sukhu, L., Doh, S.G., Hartikka, J., Nietupski, J., Manthorpe, M., Nichols, M., Plewe, M., Liang, X., Norman, J., Smith, A. & Cheng, S.H. (1996). A novel cationic lipid greatly enhances plasmid DNA delivery and expression in mouse lung. *Proc Natl Acad Sci U S A*, **93**, 11454-9.
- Widera, A., Beloussow, K., Kim, K.J., Crandall, E. & Shen, W.C. (2001). KGF elicits retention of transferrin-receptor expression in rat alveolar epithelial cell monolayers. *Proceed. Int'l. Symp. Control. Rel. Bioact. Mater.* 28 (2001), *Controlled Release Society*, #6215.
- Wightman, L., Kircheis, R., Rossler, V., Carotta, S., Ruzicka, R., Kursa, M. & Wagner, E. (2001). Different behavior of branched and linear polyethylenimine for gene delivery in vitro and in vivo. *J Gene Med*, **3**, 362-72.
- Wolff, J.A., Malone, R.W., Williams, P., Chong, W., Acsadi, G., Jani, A. & Felgner, P.L. (1990). Direct gene transfer into mouse muscle in vivo. *Science*, **247**, 1465-8.
- Wyllie, J.C. (1977). Transferrin uptake by rabbit alveolar macrophages in vitro. *Br J Haematol*, **37**, 17-24.
- Yang, N.S., Burkholder, J., Roberts, B., Martinell, B. & McCabe, D. (1990). In vivo and in vitro gene transfer to mammalian somatic cells by particle bombardment. *Proc Natl Acad Sci U S A*, **87**, 9568-72.
- Yew, N.S., Wysokenski, D.M., Wang, K.X., Ziegler, R.J., Marshall, J., McNeilly, D., Cherry, M., Osburn, W. & Cheng, S.H. (1997). Optimization of plasmid vectors for high-level expression in lung epithelial cells. *Hum Gene Ther*, **8**, 575-84.
- Zabner, J., Fasbender, A.J., Moninger, T., Poellinger, K.A. & Welsh, M.J. (1995). Cellular and molecular barriers to gene transfer by a cationic lipid. *J Biol Chem*, **270**, 18997-9007.
- Zanta, M.A., Belguise-Valladier, P. & Behr, J.P. (1999). Gene delivery: a single nuclear localization signal peptide is sufficient to carry DNA to the cell nucleus. *Proc Natl Acad Sci U S A*, **96**, 91-6.

- 
- Zauner, W., Ogris, M. & Wagner, E. (1998). POLYLYSINE-BASED TRANSFECTION SYSTEMS UTILIZING RECEPTOR-MEDIATED DELIVERY [Review]. *Advanced Drug Delivery Reviews*, **30**, 97-113.
- Zhang, G., Budker, V. & Wolff, J.A. (1999). High levels of foreign gene expression in hepatocytes after tail vein injections of naked plasmid DNA. *Hum Gene Ther*, **10**, 1735-7.
- Zou, S.M., Erbacher, P., Remy, J.S. & Behr, J.P. (2000). Systemic linear polyethylenimine (L-PEI)-mediated gene delivery in the mouse. *J Gene Med*, **2**, 128-34.