7. References

Afzai, M.; Ahmad, M. (1982): Effect of juvenile hormone analogues on colony foundation and caste-differentiation in *Bifiditermes beesoni* (Gardner) (Isoptera: Kalotermitidae). Material u. Organismen 17: 35-49.

Ahmad, M.; Salihah, Z.; Sultana, N.; Ahmad, S. (1986): Preliminary studies on the effects of diflubenzuron (Dimilin) on termites (Isoptera). Pakistan J. Zool. 18: 403-409.

Akhtar, M. S.; Shahid, A. S. (1993): Termites as pest of agricultural crops in Pakistan. Pakistan J. Zool. 25: 187-193.

Araujo, R. L. (1970): Termites of the Neotropical Region. In: Krishna, K.; Weesner, F. M. (Eds.): Biology of Termites. Vol. 2. New York (Acad. Press) 527-576.

Chu, Hsang-Hsoun; Tchi-Dar; T. T.; Tse-Fu, C.; Mei-Wen, K. (1974): Induction of soldier differentiation in *Reticulitermes flaviceps* (Oshima) with juvenile hormone analogues. Acta Entomol. Sinica, 17: 161-165.

Dai J. D. (1980): Induction and inhibition of soldier differentiating potentials in *Reticulitermes flaviceps* (Oshima) with juvenile hormone analogues. Acta Entomol. Sinica, 23: 374-380.

De Wilde, J.; Beetsma, J. (1982): The physiology of caste development in social insects. Adv. Insect Physiol. 16: 167-246.

Doppelreiter, V. H.; Korioth, M. (1981): Entwicklulungshemmung bei den Bodentermiten *Heterotermes indicola* und *Reticulitermes flavipes.* Z. Angew. Entomol. 91: 131-137.

Dorn, S.; Frischknecht, M. L.; Martinez, V.; Zurflüh, R.; Fischer, U. (1981): A novel non-neurotoxic insecticide with a broad activity spectrum. Z. Pflanzenschutz 88: 269-275.

Deligne, J.; Quennedy, A.; Blum, M. (1981): The enemies and defense mechanisms of termites. In: Herman, H. R. (Ed.): Social Insects. Vol. 2. New York (Acad. Press) 1-76.

Doki, H.; Tsunoda, K.; Nishimioto, K. (1984): Effect of juvenile hormone analogues on caste-differentiation of the termites *Reticulitermes speratus* (Kolbe) (Isoptera: Rhinotermitidae). Material u. Organismen 19: 175-187.

Duncan, F. D. (1997): Behavioural responses to poison baits by the termite *Hodotermes mossambicus* (Hagen). Insect Sci. Applic. 17:2, 221-225.

Edward, R.; Mill, A.E. (1986): Termites in buildings. Their biology and control. Rentokil Ltd., East Grinsted.

Eggleton, P. (1999): Termites species description rates and the state of termite taxonomy. Insectes Sociaux 46: 1-5.

El Bakri, A. (1986): Foraging activity and fungicidal control of the fungus growing termite *Microtermes* (Isoptera: Macrotermitinae) in the Sudan Ph. D. Thesis, Univ. Khartoum, Sudan.

El Bakri, A.; Eldein, N.; Kambal, M. A.; Thomas, R. J.; Wood, T. G. (1989): Colony foundation and development in *Microtermes* sp. nr. *albopartitus* (Isoptera: Macrotermitinae). Sociobiol. 15: 175-180

Emerson, A. E. (1955): Geographical origins and dispersions of termite genera. Fieldiana Zool. 37: 467-521.

Evans.T.A., Lenz, M.; Glleeson, P.V.(1998): Testing assumptions of mark-recapture protocols for estimating population size using Australian mound-building subterranean termites.Ecol. Entomol. 23:139-159.

Fragalla, A. A.; Badawi, A. I.; Dabbour, A. I. (1985): Field evaluation of the effects of the juvenile hormone analogues (JHAs) and diflubenzuron (Dimilin) on termites of the genus *Microcerotermes* (Isoptera: Termitidae) in the central region of Saudi Arabia. Sociobiol. 11: 29–37.

French, J. R. J. (1974): A juvenile hormone analogue inducing caste-differentiation in the Australian termite, *Nasutitermes exitiosus* (Hill) (Isoptera: Termitidae). J. Austral. Entomol. Soc. 13: 353-355.

French, J. R. J.; Robinson, P. J.; Yazaki, Y.; Hillis, W. E. (1979): Bioassays of extracts from white cypress pine (C*allitris columellaris* F. Muell.) against subterranean termites. Holzforsch. 33 (5): 144 -148.

Forschler, B. T.;Jenkins, T.M. (1999): Evaluation of subterranean termite, biology using genetic, chemotaxonomic and morphometric markers and ecological data: a testimonial for multi- disciplinary efforts. Trends Entomol. 2: 71-80.

Forschler, B. T.; Ryder Jr., J. C. (1996): Subterranean termite, *Reticulitermes* spp. (Isoptera: Rhinotermitidae), colony response to baiting with hexaflumuron using a prototype commercial termite baiting system. J. Entomol. Sci. 31 (2): 143-151.

Grenier, S.; Grenier A. M. (1993): Fenoxycarb, a fairy new insect growth regulator: a review of its effects on insects. Ann. Appl. Biol. 122: 369-413.

Harris, W. V. (1968): Termites of Sudan. Sudan Nat. Hist. Mus. Bull. 6: 1-29.

Harris, W. V. (1971): Termites, their recognition and control. London (Longmans, Green and Co.) 186pp.

Haverty, M. I. (1977): The proportion of soldiers in termite colonies: A list and a bibliography (Isoptera). Sociobiol. 2: 199-216.

Haverty, M. I. (1979): Selection of tunneling substrates for laboratory studies with three subterranean termite species. Sociobiol. 4: 315-320.

Haverty, M. I.; Howard, R. W. (1979): Effects of insect regulators on subterranean termites: induction of differentiation, defaunation, and starvation. Ann. Entomol. Soc. Amer. 72: 503-508.

Haverty, M. I.; Howard, R. W. (1981): Production of soldiers and maintenance of soldier proportions by laborotory experimental groups of *Reticulitermes flavipes* (Kollar) and *Reticulitermes virginicus* (Banks) (Isoptera: Rhinotermitidae). Insectes Sociaux 28: 32-39.

Haverty, M. I.; Su, N. Y.; Tamashiro, M.; Yamamoto, R. (1989): Concentration-dependent presoldier induction and feeding deterrency: Potential of two growth regulators for remedial control of the Formosan subterranean termite. J. Econ. Entomol. 82: 1370-1374.

Hermann, H. R. (1979): Social insects. Vol. 1. London (Acad. Press).

Howard, R. H. (1984): Effects of methoprene on *Reticulitermes flavipes* (Kollar) (Isoptera: Rhinotermitidae). J. Georgia Entomol. Soc. 19: 291-298.

Howard, R. H.; Haverty, M. I. (1978): Defaunation, mortality and soldier differentiation: concentration effects of methoprene in a termite. Sociobiol. 3: 73-78.

Howard, R. H.; Haverty, M. I. (1979): Termites and juvenile hormone analogues: a review of methodology and observed effects. Sociobiol. 4: 269-278.

Howard, R. H.; Haverty, M. I. (1979): Comprison of feeding substrates for evaluating effects of insect growth regulators on subterranean termites. J. Georgia. Entomol. Soc. 14: 3-7.

Hrdy, I. (1972): Der Einfluss von zwei Juvenilhormonalogen auf die Differenzierung der Soldaten bei *Reticulitermes lucifugus santonensis (*Feyt.) (Isoptera: Rhinotermitidae). Z. Angew. Entomol. 72: 129-143.

Hrdy, I. (1976): The influence of juvenile hormone analogues on caste determination in termites. Lüscher, M. (Ed.): Phase and Caste Determination in Insects: Endocrine Aspects. Oxford (Pergamon Press) 71.

Hrdy, I.; Kreicek, J. (1972): Development of superfluous soldiers induced by juvenile hormone analogues in the termite, *Reticulitermes lucifugus santonensis.* Insectes Sociaux 19: 105-109.

Hrdy, I.; Krecek, J.; Zuskova, Z. (1979): Juvenile hormone analogues: effect on the soldier caste differentiation in termites (Isoptera). Vestn. Cesk. Spol. Zool. 43: 260-269.

Hunter, E.; Vincent, J. F. (1974): The effect of novel insecticide on insect cuticle. Experimenta 30: 1432-1433.

Johnson, B. A.; Lamb, R. W.; Wood, T. G. (1981): Termite damage and crop loss studies in Nigeria. A survey of damage to groundnuts. Trop. Pest Managem. 27 (3): 325-342.

Jones, S. C. (1984): Evaluation of two insect growth regulators for the bait–block method of subterranean termites (Isoptera: Rhinotermitidae) control. J. Econ. Entomol. 77: 1086-1091.

Jones, S. C. (1987): Effects of methoprene on *Coptotermes formoanus* (Isoptera: Rhinotermitidae). International Research Group on Wood Preservation, Doc. IRG/WP/1322.

Jones, S. C. (1987): Extended exposure of two insect growth regulators on *Coptotermes formoanus* (Shiraki). In: Tamashiro, M.; Su, N. Y. (Eds.): Biology and Control of the Formosan subterranean Termites. University of Hawaii, Honolulu, Research Extension Ser. 083, 58-61.

Jones, S. C. (1989): Field evaluation of fenoxycarb as a bait toxicant for subterranean termite control. Sociobiol. 15: 33-41.

Jones, S. C. (1990): Effect of population density on tunneling by Formosan subterranean termites and east subterranean termites (Isoptera: Rhinotermitidae) through treated soil. J. Econ. Entomol. 83 (3): 875-878.

Kambhampati, S.; Eggleton (2000): Taxonomy and phylogeny of termites. In: Abe et al. (Eds.): Termites: Evoluation, Sociality, Symbioses, Ecology. Amsterdam (Kluwer Acad.) 1-23.

Ker, R. F. (1977): Investigation of locust cuticle using the insecticide Diflubenzuron. J. Insect Physiol.

Krishna, K.; Weesner, F. M. (1969): Biology of termites. Vol. 1. New York (Acad. Press).

Krishna, K.; Weesner, F. M. (1970): Biology of termites. Vol. 2. New York (Acad. Press).

Lee, K. E.; Wood, T. G. (1971): Termites and soils. New York (Acad. Press) 251pp.

Lelis, A. T.; Everaerts, C. (1993): Effects of juvenile hormone analogues upon soldier differentiation in the termites (Rhinotermitidae, Hetero-termitinae). J. Morphol. 217: 239-261.

Lenz, M. (1976): The dependence of hormone effects in termites caste determination on external factors. In: Lüscher, M. (Ed.): Phase and Caste Determination in Insects: Endocrine Aspects. New York (Pergamon Press Inc.) 73-89. .

Lenz, M.; Gleeson, P. V.; Miller, L. R.; Abbey, H. M. (1996): How predictive are laboratory experiments for assessing the effects of termites. A comparison of laboratory and field data from Australian mound building species of termite. International Research Group on Wood Preservation, Doc. IRG/WP/96-10143, 11pp.

Lobry de Bruyn, L. A.; Conacher, A. J. (1990): The role of termites and ants in soil modification: a review.Aust. J. of soil Res.28: 55-93.

Logan, J.W.M.; Cowie,R.H.; Wood T. G.(1990): Termite (Isoptera) control in Agriculture and forestry by non-chemical methods: a review: Bull. Entomol.Res.80: 309-330.

Logan, J.W.M.;,Rajagopal, D. ;Wightman,J.A. ; Pearce, M.J.(1992) : Control of termites and other soil pests of groundnuts with special reference to controlled release formulations of non- persistent insecticides in India and Sudan. Bull.Entomol. Res. 82:57-66.

Lüscher, M. (1960): Hormonal control of caste differentiation in termites. Ann. New York Acad. Sci. 89: 549-563.

Lüscher, M. (1969): Die Bedeutung des Juvenilhormons für die Differenzierung der Soldaten bei der Termite *Kalotermes flavicollis*. In: Proc. 6th Internat. Congr. IUSSI, Bern, 268-273.

Lüscher, M. (1976): Evidence for an endocrine control of caste determination in higher termites. In: Lüscher, M. (Ed.): Phase and Caste Determination in Insects: Endocrine Aspects. New York (Pergamon Press Inc.) 91-103.

Mando, A. (1997): Effect of termites and mulch on the physical rehabilitation of structurally crusted soils in the sahel. Land Degradn. and Dev. 8: 269-278.

Marks, E. P.; Sowa, B. A. (1974): An in vitro model system for the production of insect cuticle. In: Mechanisms of Pesticide Action. ACS Symp. Ser. 144-155.

Masner, P.; Dorn, S.; Vogel, W. K.; Graf, O.; Günthart (1981): Types of response of insects to a new IRG and to proven standards. In: Sehnal Zabza, F. A.; Menn, J. J.; Cymborowski (Eds.): Regulation of Insect Development and Behaviour. Wroclaw (Wroclaw Techn. Univ. Press) 809-811.

Masner, P.; Angest, M.; Dorn, S. (1987): Fenoxycarb, an insect regulator with juvenile hormone activity: A candidate for *Heliothis virescens* (F.) control on cotton. Pesticide Sci. 18: 89-94.

Meyer, D.; Lüscher, M. (1973): Juvenile hormone activity in the haemolymph and the anal secretion of the queen of *Macrotermes subhyalinus* (Rambur) (Isoptera, Termitidae)In: Proc. 7th Internat. Congr. IUSSI, London, 268-273.

Mill, A. E. (1992): Termites as agricultural pests in Amazonia, Brazil. Outlook Agric. 21 (1): 41-46.

Mohamed, M. A. (1991): Foraging activity and control of the fungus growing termite *Microtermes* (Isoptera: Macrotermitinae) in the Sudan. M. Sc. Thesis, Univ. Khartoum, Sudan.

Noirot, C. (1969): Formation of castes in the higher termites. In: Krishna, K.; Weesner, F. M. (Eds.): Biology of Termites. Vol. 1. New York (Acad. Press) 311-350.

Noirot, C. (1974): Polymorphismus bei höheren Termiten. In: Schmidt, G. H. (Ed.): Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich. New York (Acad. Press) 73-25.

Noirot, C. (1985): Pathways of caste development in the lower termites. In: Watson, J. A. L.; Okot-Kotber, B. M.; Noirot, C. (Eds.): Caste Differentiation in Social Insects. Oxford (Pergamon Press) 41-57.

Oberlander, H.; Silhacek, D. L.; Eli, S.; Ishaaya, I. (1997): Current status and future perspectives of the use of insect growth regulators for the control of stored product insects. J. Stored Prod. 33: 1-6.

Ogino, K.; Hirono, Y.; Matsumoto, T. (1993): Juvenile hormone analogue, S-31183, causes a high level induction of presoldiers differentiation in the Japanese damp wood termite. J. Zool. Sci. 10: 361-366.

Okot-Kotber, B. M. (1980a): The influence of juvenile hormone analogues (ZR 515, methoprene) on soldiers differentiation in *Macrotermes michaelseni* (Isoptera; Macrotermitinae). Physiol. Entomol. 5: 407-416.

Okot-Kotber, B. M. (1980b): Competence of *Macrotermes michaelseni* (Isoptera; Macrotermitinae) larvae to differentiate into soldiers under the influence of juvenile hormone analogues (ZR 515, methoprene). J. Insect Physiol. 26: 655-659.

Okot-Kotber, B. M. (1985): Mechanism of caste-differentiation in a higher termites, *Macrotermes michaelseni* (Isoptera; Macrotermitinae). In:Watson, J. A. L.; Okot-Kotber, B. M.; Noirot, C. (Eds.): Caste Determination in Social Insects. Oxford (Pergamon Press) 267-306.

Pallaske, M.; Metzner, W.; Wegen, H. W. (1993): Mittel zum Schutz technischer Materialien, insbesondere Holz und Holzwerkstoffe. Patentschrift DE 43 03 012 A1.

Pallaske, M. (1997): Insect growth regulators: modes of action and mode of action-dependent peculiarities. International Research Group on Wood Preservation, Doc. IRG/WP/97-301155.

Pearce, M. J.; Waite, B.S. (1994): A list of termite genera (Isoptera) with comments on taxonomic changes and regional distribution. Sociobiol. 23: 247-263.

Peppuy, A.; Robert, A.; Delbecque, J. P.; Leca, J. L.; Roulland, C.; Bordereau, C. (1998) : Efficacy of Hexaflumuron against the fungus growing termite *Pseudacanthotermes spiniger* (Sjöstedt) (Isoptera, Macrotermitinae) Pestic. Sci. 54: 22 - 26.

Post, L. C.; De Jong, B. G.; Vincent, W. R. (1974): 1-(2,6-Disubstituated benoyl)-3-phenylurea insecticides: inhibitors of chitin synthesis. Pesticide Biochem. Physiol. 4: 473-483.

Sands, W. A. (1981): The social life of termite. Central Assoc. Beekeepers, Essex, 1-8.

Slama, K.; Wimmer, Z.; Romanuk, M.; Sorm, F. (1974): Insect hormones and bioanalogues. Wien (Springer-Verl.).

Snyder, T. E. (1949): Catalog of the termites (Isoptera) of the world. Smithson. Misc. Collections 112: 1-490.

Springhetti, A. (1974): The influence of farnesenic acid ethyl ester on the differentiation of *Kalotermes flavicolis* (Fabr). (Isoptera) soldiers. Experientia, 30: 1197-1198.

Staal, G. B. (1975): Insect growth regulators with juvenile hormone activity. Ann. Rev. Entomol. 20: 417-460.

Stuart, A. M. (1969): Social behavior and communication. In: Krishna, K.; Weesner, F. W. (Eds.): Biology of Termites. Vol. 1. New York (Acad. Press).

Su, N.-Y. (1990): Measuring termiticides. Pest Control Mag. 58: 24, 30, 34, 35.

Su, N.-Y. (1991): Evaluation bait toxicants for suppression of subterranean termite populations. Sociobiol. 19: 211-220.

Su, N.-Y. (1994): Field evaluation of a hexaflumuron bait for population suppression of subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 87: 389-397.

Su, N.-Y.; Scheffrahn, R. H. (1989): Comparative effect of an insect growth regulator, S-31183, against the Formosan subterranean termites and east subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 82: 1125-1129.

Su, N.-Y.; Scheffrahn, R. H. (1990): Potential of insect growth regulators as termiticides: a review. Sociobiol. 17: 313-328.

Su, N.-Y.; Scheffrahn, R. H. (1991): Laboratory evaluation of two slow-acting toxicants against Formosan and east subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 84: 170-175.

Su, N.-Y.; Scheffrahn, R. H. (1993): Laboratory evaluation of two chitin synethesis inhibitors, hexaflumuron and diflubenzuron, as bait toxicants against Formosan and east subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 86: 1453-1457.

Su, N.-Y.; Scheffrahn, R. H. (1996): Comparative effect of two chitin synthesis inhibitors, hexaflumuron and lufenuron, in a bait matrix against subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 89: 1156-1160.

Su, N.-Y.; Ban, P. M.; Scheffrahn, R. H. (1991): Suppression of foraging populations of Formosan subterranean termites (Isoptera: Rhinotermitidae) by field applications of a slow-acting toxicant bait. J. Econ. Entomol. 84: 1525-1531.

Su, N.-Y.; Ban, P. M.; Scheffrahn, R. H. (1996): An aboveground station for monitoring structure-infesting populations of Formosan subterranean termites (Isoptera: Rhinotermitidae). Sociobiol. 27: 39-45.

Su, N.-Y.; Ban, P. M.; Scheffrahn, R. H. (1997): Remedial baiting with hexaflumuron in aboveground stationsto control structure-infesting populations of the Formosan subterranean termite (Isoptera: Rhinotermitidae). J. Econ. Entomol. 90: 809-817.

Su, N.-Y.; Tamashiro, M.; Haverty, M. I. (1985): Effect of three insect growth regulators, feeding substrates, and colony origin on survival and presoldiers production of the Formosan subterranean termites (Isoptera: Rhinotermitidae). J. Econ. Entomol. 78: 1259-1263.

Su, N.-Y.; Tamashiro, M.; Yates, J. R.; Haverty, M. I. (1982): Effect of behavior on the evolution of insecticides for the prevention of or remedial control of the Formosan subterranean termite. J. Econ. Entomol. 75: 188-193.

Su, N.-Y.; Thomas, E. M.; Ban, P. M.; Scheffrahn, R. H. (1995): A monitoring/ baiting station to detect and eliminate foraging population of subterranean termites (Isoptera: Rhinotermitidae) near structures. J. Econ. Entomol. 88: 932-936.

Tsunoda, K.; Doki, H.; Nishimoto, K. (1986): Effect of developmental stages of workers and nymph of *Reticulitermes speratus* (Kolbe) (Isoptera: Rhinotermitidae) on caste differenation induced by JHA treatment. Material u. Organismen 21: 47-61.

Tsunoda, K.; Matsuoka, H.; Yoshimura, T. (1998): Colony elimination of *Reticulitermes speratus* by bait application and the effect on foraging territory. J. Econ. Entomol. 91(6): 1383-1386.

Unistat (1995): Statstical package, Unistat Ltd., London-Version 4.5.

Valcke, A.; Pallaske, M. (1995): Flurox a new breakthrough in insect control for wood preservation. International Research Group on Wood Preservation, Doc. IRG/WP/95-30079.

Varma, R. V. (1977): The influence of juvenile hormone analogue, farnesyl methyl ether on caste development in termites *Postelectrotermes nayari* (Roonwal and Verma, 1971). Indian J. Exper. Biol. 15: 564-565.

Verloop, A.; Ferrell, C. D. (1977): Pesticide chemistry in the 20th Century. ACS Symp. Ser. 237-271.

Waller, D. A.; La Fage, J. P. (1987): Nutritional ecology of termites. P. 487-532. In: Slansky, F.; Rodriguez, J. G. (Eds.): Nutritional Ecology of Insects, Mites and Spiders. New York (Wiley and Sons).

Wanyoni, K. (1974): The influence of juvenile hormone analogues (ZR 512, Zoecon) on caste development in *Zootermopsis nevadensis* (Hagen) (Isoptera). Insectes Sociaux 21: 35-44.

Wanyoni, K.; Lüscher, M. (1973): The action of juvenile hormone analogues on caste development in *Zootermopsis* (Isoptera). In: Proc. 7th Internat. Congr. IUSSI, London, 392-395.

Wightman, J. A.; Wightman, A. S. (1994): An insect, agronomic and sociological survey of groundnut fields in southern Africa. Agric., Ecosystems and Environm. 51: 311-331.

Wigglesworth, V. B. (1970): Insect hormones. San Francisco (W. H. Freeman).

Wood, T. G.; Johnson, R. A. (1983): Modification of soils in Nigerian savanna by soil feeding *Cubitermes* (Isoptera,Termitidae). Soil Biol. and Biochem. 15: 575-579.

Wood, T. G.; Johnson, R. A. (1986): The biology, physiology and ecology of termites.In Vinson, S.B.(ed.). Economic Impact and Control of Social Insects.Praeger Publishers, New York, pp. 1-68.

Wood, T. G.; Smith, R. W.; Johnson, R. A.; Komolafe, P. O. (1980a): Termite damage and crop loss studies in Nigeria. Pre-harvest loves to yams due to termites, and other soil pests. Trop Pest Managem. 26: 355-370.

Wood, T. G.; Johnson, R. A.; Ohiagu, C. E. (1980b): Termite damage and crop loss studies in Nigeria, a review of termite (Isoptera) damage to maize and estimation of damage, loss in yield and termite (*Microtermes*) abundance at Mokwa, Nigeria. Trop Pest Managem. 26 (3): 241-253.

Wood, T. G. (1996): The agricultural importance of termites in the tropics. Agric. Zool. Rev. 7: 117-155.