

5. Literatur

- Aalfs, J.D. und R.E. Kingston. 2000. What does 'chromatin remodeling' mean? *Trends Biochem Sci.* 25:548-55.
- Albright, S.R. und R. Tjian. 2000. TAFs revisited: more data reveal new twists and confirm old ideas. *Gene.* 242:1-13.
- Balbirnie, M., R. Grothe und D.S. Eisenberg. 2001. An amyloid-forming peptide from the yeast prion Sup35 reveals a dehydrated beta-sheet structure for amyloid. *Proc Natl Acad Sci U S A.* 98:2375-80.
- Bates, G.P., L. Mangiarini, A. Mahal und S.W. Davies. 1997. Transgenic models of Huntington's disease. *Hum Mol Genet.* 6:1633-7.
- Becher, M.W., J.A. Kotzuk, A.H. Sharp, S.W. Davies, G.P. Bates, D.L. Price und C.A. Ross. 1998. Intranuclear neuronal inclusions in Huntington's disease and dentatorubral and pallidolusian atrophy: correlation between the density of inclusions and IT15 CAG triplet repeat length. *Neurobiol Dis.* 4:387-97.
- Beighton, P. und M.R. Hayden. 1981. Huntington's chorea. *S Afr Med J.* 59:250.
- Biere, A.L., S.J. Wood, J. Wypych, S. Steavenson, Y. Jiang, D. Anafi, F.W. Jacobsen, M.A. Jarosinski, G.M. Wu, J.C. Louis, F. Martin, L.O. Narhi und M. Citron. 2000. Parkinson's disease-associated alpha-synuclein is more fibrillogenic than beta- and gamma-synuclein and cannot cross-seed its homologs. *J Biol Chem.* 275:34574-9.
- Boutell, J.M., P. Thomas, J.W. Neal, V.J. Weston, J. Duce, P.S. Harper und A.L. Jones. 1999b. Aberrant interactions of transcriptional repressor proteins with the Huntington's disease gene product, huntingtin. *Hum Mol Genet.* 8:1647-55.
- Bruijn, L.I., M.K. Houseweart, S. Kato, K.L. Anderson, S.D. Anderson, E. Ohama, A.G. Reaume, R.W. Scott und D.W. Cleveland. 1998. Aggregation and motor neuron toxicity of an ALS-linked SOD1 mutant independent from wild-type SOD1. *Science.* 281:1851-4.
- Burke, J.R., J.J. Enghild, M.E. Martin, Y.S. Jou, R.M. Myers, A.D. Roses, J.M. Vance und W.J. Strittmatter. 1996. Huntingtin and DRPLA proteins selectively interact with the enzyme GAPDH. *Nat Med.* 2:347-50.
- Burley, S.K. 1996. The TATA box binding protein. *Curr Opin Struct Biol.* 6:69-75.

- Bussow, K., D. Cahill, W. Nietfeld, D. Bancroft, E. Scherzinger, H. Lehrach und G. Walter. 1998. A method for global protein expression and antibody screening on high- density filters of an arrayed cDNA library. *Nucleic Acids Res.* 26:5007-8.
- Caputo, C.B., P.E. Fraser, I.E. Sobel und D.A. Kirschner. 1992. Amyloid-like properties of a synthetic peptide corresponding to the carboxy terminus of beta-amyloid protein precursor. *Arch Biochem Biophys.* 292:199-205.
- Carmichael, J., J. Chatellier, A. Woolfson, C. Milstein, A.R. Fersht und D.C. Rubinsztein. 2000. Bacterial and yeast chaperones reduce both aggregate formation and cell death in mammalian cell models of Huntington's disease. *Proc Natl Acad Sci U S A.* 97:9701-5.
- Cha, J.H. 2000. Transcriptional dysregulation in Huntington's disease. *Trends Neurosci.* 23:387-92.
- Cha, J.H., C.M. Kosinski, J.A. Kerner, S.A. Alsdorf, L. Mangiarini, S.W. Davies, J.B. Penney, G.P. Bates und A.B. Young. 1998. Altered brain neurotransmitter receptors in transgenic mice expressing a portion of an abnormal human Huntington disease gene. *Proc Natl Acad Sci U S A.* 95:6480-5.
- Chen, D., C.S. Hinkley, R.W. Henry und S. Huang. 2002. TBP dynamics in living human cells: constitutive association of TBP with mitotic chromosomes. *Mol Biol Cell.* 13:276-84.
- Chen, S., V. Berthelie, W. Yang und R. Wetzel. 2001. Polyglutamine aggregation behavior in vitro supports a recruitment mechanism of cytotoxicity. *J Mol Biol.* 311:173-82.
- Chien, P. und J.S. Weissman. 2001. Conformational diversity in a yeast prion dictates its seeding specificity. *Nature.* 410:223-7.
- Ciechanover, A. und A.L. Schwartz. 1989. How are substrates recognized by the ubiquitin-mediated proteolytic system? *Trends Biochem Sci.* 14:483-8.
- Clarke, G., R.A. Collins, B.R. Leavitt, D.F. Andrews, M.R. Hayden, C.J. Lumsden und R.R. McInnes. 2000. A one-hit model of cell death in inherited neuronal degenerations. *Nature.* 406:195-9.
- Conway, K.A., J.D. Harper und P.T. Lansbury. 1998. Accelerated *in vitro* fibril formation by a mutant α -synuclein linked to early-onset Parkinson disease. *Nature Medicine.* 4:1318-20.

- Cooper, J.K., G. Schilling, M.F. Peters, W.J. Herring, A.H. Sharp, Z. Kaminsky, J. Masone, F.A. Khan, M. Delaney, D.R. Borchelt, V.L. Dawson, T.M. Dawson und C.A. Ross. 1998. Truncated N-terminal fragments of Huntingtin with expanded glutamine repeats form nuclear and cytoplasmic aggregates in cell culture. *Hum Mol Genet.* 7:783-90.
- David, G., N. Abbas, G. Stevanin, A. Durr, G. Yvert, G. Cancel, C. Weber, G. Imbert, F. Saudou, E. Antoniou, H. Drabkin, R. Gemmill, P. Giunti, A. Benomar, N. Wood, M. Ruberg, Y. Agid, J.L. Mandel und A. Brice. 1997. Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. *Nat Genet.* 17:65-70.
- Davies, S.W., M. Turmaine, B.A. Cozens, M. DiFiglia, A.H. Sharp, C.A. Ross, E. Scherzinger, E.E. Wanker, L. Mangiarini und G.P. Bates. 1997. Formation of neuronal intranuclear inclusions underlies the neurological dysfunction in mice transgenic for the HD mutation. *Cell.* 90:537-48.
- DiFiglia, M. 1997. Clinical Genetics, II. Huntington's disease: from the gene to pathophysiology. *Am J Psychiatry.* 154:1046.
- DiFiglia, M., E. Sapp, K.O. Chase, S.W. Davies, G.P. Bates, J.P. Vonsattel und N. Aronin. 1997. Aggregation of Huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science.* 277:1990-3.
- Dragatsis, I., P. Dietrich und S. Zeitlin. 2000. Expression of the Huntingtin-associated protein 1 gene in the developing and adult mouse. *Neurosci Lett.* 282:37-40.
- Duyao, M.P., A.B. Auerbach, A. Ryan, F. Persichetti, G.T. Barnes, S.M. McNeil, P. Ge, J.P. Vonsattel, J.F. Gusella, A.L. Joyner, et al. 1995. Inactivation of the mouse Huntington's disease gene homolog Hdh. *Science.* 269:407-10.
- Faber, P.W., J.R. Alter, M.E. MacDonald und A.C. Hart. 1999. Polyglutamine-mediated dysfunction and apoptotic death of a *Caenorhabditis elegans* sensory neuron. *Proc Natl Acad Sci U S A.* 96:179-84.
- Fink, A.L. 1998. Protein aggregation: folding aggregates, inclusion bodies and amyloid. *Fold Des.* 3:R9-23.
- Furtado, S. und O. Suchowersky. 1995. Huntington's disease: recent advances in diagnosis and management. *Can J Neurol Sci.* 22:5-12.
- Fusco, F.R., Q. Chen, W.J. Lamoreaux, G. Figueredo-Cardenas, Y. Jiao, J.A. Coffman, D.J. Surmeier, M.G. Honig, L.R. Carlock und A. Reiner. 1999. Cellular

- localization of Huntingtin in striatal and cortical neurons in rats: lack of correlation with neuronal vulnerability in Huntington's disease. *J Neurosci.* 19:1189-202.
- Garcia-Mata, R., Z. Bebok, E.J. Sorscher und E.S. Sztul. 1999. Characterization and dynamics of aggresome formation by a cytosolic GFP- chimera. *J Cell Biol.* 146:1239-54.
- Gill, S.C. und P.H. von Hippel. 1989. Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem.* 182:319-26.
- Glenner, G.G. 1980. Amyloid Deposits and Amyloidosis - the Beta-Fibrilloses 1. *New England Journal of Medicine.* 302:1283-92.
- Gourfinkel-An, I., G. Cancel, C. Duyckaerts, B. Faucheux, J.J. Hauw, Y. Trotter, A. Brice, Y. Agid und E.C. Hirsch. 1998. Neuronal distribution of intranuclear inclusions in Huntington's disease with adult onset. *Neuroreport.* 9:1823-6.
- Green, H. 1993. Human genetic diseases due to codon reiteration: relationship to an evolutionary mechanism. *Cell.* 74:955-6.
- Gusella, J.F. und M.E. MacDonald. 1995. Huntington's disease: CAG genetics expands neurobiology. *Curr Opin Neurobiol.* 5:656-62.
- Gusella, J.F. und M.E. MacDonald. 2000. Molecular genetics: unmasking polyglutamine triggers in neurodegenerative disease. *Nat Rev Neurosci.* 1:109-15.
- Gutekunst, C.-A., S.-H. Li, H. Yi, J.S. Mulroy, S. Kuemmerle, R. Jones, D. Rye, R.J. Ferrante, S.M. Hersch und X.-J. Li. 1999. Nuclear and neuropil aggregates in Huntington's disease: relationship to neuropathology. *J. Neuroscience.* 19:2522-34.
- Hackam, A.S., R. Singaraja, C.L. Wellington, M. Metzler, K. McCutcheon, T. Zhang, M. Kalchman und M.R. Hayden. 1998. The influence of Huntingtin protein size on nuclear localization and cellular toxicity. *J Cell Biol.* 141:1097-105.
- Hackam, A.S., R. Singaraja, T. Zhang, L. Gan und M.R. Hayden. 1999. In vitro evidence for both the nucleus and cytoplasmic subcellular sites of pathogenesis in Huntington's disease. *Hum Mol Genet.* 8:25-33.
- Harper, J.D. und P.T. Lansbury, Jr. 1997. Models of amyloid seeding in Alzheimer's disease and scrapie: mechanistic truths and physiological consequences of the time-dependent solubility of amyloid proteins. *Annu Rev Biochem.* 66:385-407.

- Harper, P.S. 1991. Huntington's disease. W.B. Saunders Co, Ltd, London.
- Hattula, K. und J. Peranen. 2000. FIP-2, a coiled-coil protein, links Huntingtin to Rab8 and modulates cellular morphogenesis. *Curr Biol.* 10:1603-6.
- HDCRG. 1993. A novel gene containing a trinucleotide repeat that is unstable on Huntington's disease chromosomes. *Cell.* 72:971-83.
- Hernandez, N. 1993. TBP, a universal eukaryotic transcription factor? *Genes Dev.* 7:1291-308.
- Hilditch-Maguire, P., F. Trettel, L.A. Passani, A. Auerbach, F. Persichetti und M.E. MacDonald. 2000. Huntingtin: an iron-regulated protein essential for normal nuclear and perinuclear organelles. *Human Molecular Genetics.* 9:2789-97.
- Hodgson, J.G., N. Agopyan, C.A. Gutekunst, B.R. Leavitt, F. LePiane, R. Singaraja, D.J. Smith, N. Bissada, K. McCutcheon, J. Nasir, L. Jamot, X.J. Li, M.E. Stevens, E. Rosemond, J.C. Roder, A.G. Phillips, E.M. Rubin, S.M. Hersch und M.R. Hayden. 1999. A YAC mouse model for Huntington's disease with full-length mutant Huntingtin, cytoplasmic toxicity, and selective striatal neurodegeneration. *Neuron.* 23:181-92.
- Holbert, S., I. Denghien, T. Kiechle, A. Rosenblatt, C. Wellington, M.R. Hayden, R.L. Margolis, C.A. Ross, J. Dausset, R.J. Ferrante und C. Neri. 2001. The Gln-Ala repeat transcriptional activator CA150 interacts with Huntingtin: neuropathologic and genetic evidence for a role in Huntington's disease pathogenesis. *Proc Natl Acad Sci U S A.* 98:1811-6.
- Holmberg, M., C. Duyckaerts, A. Durr, G. Cancel, I. Gourfinkel-An, P. Damier, B. Faucheux, Y. Trotter, E.C. Hirsch, Y. Agid und A. Brice. 1998. Spinocerebellar ataxia type 7 (SCA7): a neurodegenerative disorder with neuronal intranuclear inclusions. *Hum Mol Genet.* 7:913-8.
- Hope, J., L.J. Reekie, N. Hunter, G. Multhaup, K. Beyreuther, H. White, A.C. Scott, M.J. Stack, M. Dawson und G.A. Wells. 1988. Fibrils from brains of cows with new cattle disease contain scrapie-associated protein. *Nature.* 336:390-2.
- Huang, C.C., P.W. Faber, F. Persichetti, V. Mittal, J.-P. Vonsattel, M.E. MacDonald und J.F. Gusella. 1998. Amyloid Formation by Mutant Huntingtin: Threshold, Progressivity and Recruitment of Normal Polyglutamine Proteins. *Somat. Cell Mol. Genet.* 24:217-33.

- Igarashi, S., R. Koide, T. Shimohata, M. Yamada, Y. Hayashi, H. Takano, H. Date, M. Oyake, T. Sato, A. Sato, S. Egawa, T. Ikeuchi, H. Tanaka, R. Nakano, K. Tanaka, I. Hozumi, T. Inuzuka, H. Takahashi und S. Tsuji. 1998. Suppression of aggregate formation and apoptosis by transglutaminase inhibitors in cells expressing truncated DRPLA protein with an expanded polyglutamine stretch. *Nat Genet.* 18:111-7.
- Igarashi, S., Y. Tanno, O. Onodera, M. Yamazaki, S. Sato, A. Ishikawa, N. Miyatani, M. Nagashima, Y. Ishikawa, K. Sahashi, et al. 1992. Strong correlation between the number of CAG repeats in androgen receptor genes and the clinical onset of features of spinal and bulbar muscular atrophy. *Neurology.* 42:2300-2.
- Igarashi, S. und S. Tsuji. 1998. [The molecular mechanisms of the instability of the CAG repeat]. *Nippon Rinsho.* 56:1064-73.
- Jackson, G.R., I. Salecker, X. Dong, X. Yao, N. Arnheim, P.W. Faber, M.E. MacDonald und S.L. Zipursky. 1998. Polyglutamine-expanded human Huntingtin transgenes induce degeneration of Drosophila photoreceptor neurons. *Neuron.* 21:633-42.
- Jarrett, J.T. und P.T. Lansbury Jr. 1993. Seeding "One-dimensional crystallization" of amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell.* 73:1055-8.
- Jarrett, J.T. und P.T. Lansbury, Jr. 1993. Seeding "one-dimensional crystallization" of amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell.* 73:1055-8.
- Jeitner, T.M., M.B. Bogdanov, W.R. Matson, Y. Daikhin, M. Yudkoff, J.E. Folk, L. Steinman, S.E. Browne, M.F. Beal, J.P. Blass und A.J. Cooper. 2001. N(epsilon)-(gamma-L-glutamyl)-L-lysine (GGEL) is increased in cerebrospinal fluid of patients with Huntington's disease. *J Neurochem.* 79:1109-12.
- Johnston, J.A., C.L. Ward und R.R. Kopito. 1998. Aggresomes: a cellular response to misfolded proteins. *J Cell Biol.* 143:1883-98.
- Jones, A.L. 1999. The localization and interactions of Huntingtin. *Philos Trans R Soc Lond B Biol Sci.* 354:1021-7.
- Kabore, A.F., W.J. Wang, S.J. Russo und M.F. Beers. 2001. Biosynthesis of surfactant protein C: characterization of aggresome formation by EGFP chimeras containing propeptide mutants lacking conserved cysteine residues. *J Cell Sci.* 114:293-302.

- Kahlem, P., C. Terre, H. Green und P. Djian. 1996. Peptides containing glutamine repeats as substrates for transglutaminase-catalyzed cross-linking: relevance to diseases of the nervous system. *PNAS*. 93:14580-5.
- Kao, C.C., P.M. Lieberman, M.C. Schmidt, Q. Zhou, R. Pei und A.J. Berk. 1990. Cloning of a transcriptionally active human TATA binding factor. *Science*. 248:1646-50.
- Karpuj, M.V., M.W. Becher, J.E. Springer, D. Chabas, S. Youssef, R. Pedotti, D. Mitchell und L. Steinman. 2002. Prolonged survival and decreased abnormal movements in transgenic model of Huntington disease, with administration of the transglutaminase inhibitor cystamine. *Nat Med*. 8:143-9.
- Karpuj, M.V., H. Garren, H. Slunt, D.L. Price, J. Gusella, M.W. Becher und L. Steinman. 1999. Transglutaminase aggregates Huntingtin into nonamyloidogenic polymers, and its enzymatic activity increases in Huntington's disease brain nuclei. *Proc Natl Acad Sci USA*. 96:7388-93.
- Kawaguchi, Y., T. Okamoto, M. Taniwaki, M. Aizawa, M. Inoue, S. Katayama, H. Kawakami, S. Nakamura, M. Nishimura, I. Akiguchi et al. 1994. CAG expansions in a novel gene for Machado-Joseph disease at chromosome 14q32.1 [see comments]. *Nat Genet*. 8:221-8.
- Kazantsev, A., E. Preisinger, A. Dranovsky, D. Goldgaber und D. Housman. 1999a. Insoluble detergent-resistant aggregates form between pathological and nonpathological lengths of polyglutamine in mammalian cells. *Proc Natl Acad Sci U S A*. 96:11404-9.
- Kazantsev, A., E. Preisinger, A. Dranovsky, D. Goldgaber und D. Housman. 1999b. Insoluble detergent-resistant aggregates form between pathological and nonpathological lengths of polyglutamine in mammalian cells. *Proc Natl Acad Sci U S A*. 96:11404-9.
- Kazantsev, A., H.A. Walker, N. Slepko, J.E. Bear, E. Preisinger, J.S. Steffan, Y.Z. Zhu, F.B. Gertler, D.E. Housman, J.L. Marsh und L.M. Thompson. 2002. A bivalent Huntingtin binding peptide suppresses polyglutamine aggregation and pathogenesis in *Drosophila*. *Nat Genet*. 30:367-76.
- Kazemi-Esfarjani, P. und S. Benzer. 2000. Genetic suppression of polyglutamine toxicity in *Drosophila*. *Science*. 287:1837-40.

- Kegel, K.B., M. Kim, E. Sapp, C. McIntyre, J.G. Castano, N. Aronin und M. DiFiglia. 2000. Huntingtin expression stimulates endosomal-lysosomal activity, endosome tubulation, and autophagy. *J Neurosci.* 20:7268-78.
- Kegel, K.B., A.R. Meloni, Y. Yi, Y.J. Kim, E. Doyle, B.G. Cuiffo, E. Sapp, Y. Wang, Z.H. Qin, J.D. Chen, J.R. Nevins, N. Aronin und M. DiFiglia. 2002. Huntingtin is present in the nucleus, interacts with the transcriptional corepressor C-terminal binding protein, and represses transcription. *J Biol Chem.* 277:7466-76.
- Kim, M., H.S. Lee, G. LaForet, C. McIntyre, E.J. Martin, P. Chang, T.W. Kim, M. Williams, P.H. Reddy, D. Tagle, F.M. Boyce, L. Won, A. Heller, N. Aronin und M. DiFiglia. 1999. Mutant Huntingtin expression in clonal striatal cells: dissociation of inclusion formation and neuronal survival by caspase inhibition. *J Neurosci.* 19:964-73.
- Kim, N.S., T. Yamaguchi, S. Sekine, M. Saeki, S. Iwamuro und S. Kato. 1998. Cloning of human polyubiquitin cDNAs and a ubiquitin-binding assay involving its in vitro translation product. *J Biochem (Tokyo).* 124:35-9.
- Koide, R., T. Ikeuchi, O. Onodera, H. Tanaka, S. Igarashi, K. Endo, H. Takahashi, R. Kondo, A. Ishikawa, T. Hayashi et al. 1994. Unstable expansion of CAG repeat in hereditary dentatorubral- pallidoluysian atrophy (DRPLA). *Nat Genet.* 6:9-13.
- Koide, R., S. Kobayashi, T. Shimohata, T. Ikeuchi, M. Maruyama, M. Saito, M. Yamada, H. Takahashi und S. Tsuji. 1999. A neurological disease caused by an expanded CAG trinucleotide repeat in the TATA-binding protein gene: a new polyglutamine disease? *Hum Mol Genet.* 8:2047-53.
- Komuro, A., M. Saeki und S. Kato. 1999a. Association of two nuclear proteins, Npw38 and NpwBP, via the interaction between the WW domain and a novel proline-rich motif containing glycine and arginine. *J Biol Chem.* 274:36513-9.
- Komuro, A., M. Saeki und S. Kato. 1999b. Npw38, a novel nuclear protein possessing a WW domain capable of activating basal transcription. *Nucleic Acids Res.* 27:1957-65.
- Koo, E.H., P.T. Lansbury und J.W. Kelly. 1999. Amyloid diseases: Abnormal protein aggregation in neurodegeneration. *Proceedings of the National Academy of Sciences of the United States of America.* 96:9989-90.

- Kopito, R.R. 2000. Aggresomes, inclusion bodies and protein aggregation. *Trends Cell Biol.* 10:524-30.
- Krobitsch, S. und S. Lindquist. 2000. Aggregation of Huntingtin in yeast varies with the length of the polyglutamine expansion and the expression of chaperone proteins. *Proc Natl Acad Sci U S A.* 97:1589-94.
- Kuemmerle, S., C.A. Gutekunst, A.M. Klein, X.J. Li, S.H. Li, M.F. Beal, S.M. Hersch und R.J. Ferrante. 1999. Huntington aggregates may not predict neuronal death in Huntington's disease. *Ann Neurol.* 46:842-9.
- Kyte, J. und R.F. Doolittle. 1982. A simple method for displaying the hydrophobic character of a protein. *J Mol Biol.* 157:105-32.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227:680-5.
- Leavitt, B.R., J.A. Guttman, J.G. Hodgson, G.H. Kimel, R. Singaraja, A.W. Vogl und M.R. Hayden. 2000. Wild-Type Huntingtin Reduces the Cellular Toxicity of Mutant Huntingtin In Vivo. *Am J Hum Genet.* 68.
- Lee, T.I., J.J. Wyrick, S.S. Koh, E.G. Jennings, E.L. Gadbois und R.A. Young. 1998. Interplay of positive and negative regulators in transcription initiation by RNA polymerase II holoenzyme. *Mol Cell Biol.* 18:4455-62.
- Li, H., S.H. Li, H. Johnston, P.F. Shelbourne und X.J. Li. 2000. Amino-terminal fragments of mutant Huntingtin show selective accumulation in striatal neurons and synaptic toxicity. *Nature Genet.* 25:385-9.
- Li, M., S. Miwa, Y. Kobayashi, D.E. Merry, M. Yamamoto, F. Tanaka, M. Doyu, Y. Hashizume, K.H. Fischbeck und G. Sobue. 1998. Nuclear inclusions of the androgen receptor protein in spinal and bulbar muscular atrophy. *Ann Neurol.* 44:249-54.
- Lunkes, A. und J.L. Mandel. 1997. Polyglutamines, nuclear inclusions and neurodegeneration [news]. *Nat Med.* 3:1201-2.
- Lunkes, A. und J.L. Mandel. 1998. A cellular model that recapitulates major pathogenic steps of Huntington's disease. *Hum Mol Genet.* 7:1355-61.
- Luthi-Carter, R., A. Strand, N.L. Peters, S.M. Solano, Z.R. Hollingsworth, A.S. Menon, A.S. Frey, B.S. Spektor, E.B. Penney, G. Schilling, C.A. Ross, D.R. Borchelt, S.J. Tapscott, A.B. Young, J.H. Cha und J.M. Olson. 2000. Decreased expression of

- striatal signaling genes in a mouse model of Huntington's disease. *Hum Mol Genet.* 9:1259-71.
- MacDonald, M.E. und J.F. Gusella. 1996. Huntington's disease: translating a CAG repeat into a pathogenic mechanism. *Curr Opin Neurobiol.* 6:638-43.
- Mangiarini, L., K. Sathasivam, M. Seller, B. Cozens, A. Harper, C. Hetherington, M. Lawton, Y. Trottier, H. Lehrach, S.W. Davies und G.P. Bates. 1996. Exon 1 of the Huntington's disease gene containing a highly expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice. *Cell.* 87:493-506.
- Maniatis, T., E.F. Fritsch und J. Sambrook. 1982. *Molecular cloning: A laboratory manual.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Marsh, J.L., H. Walker, H. Theisen, Y.Z. Zhu, T. Fielder, J. Purcell und L.M. Thompson. 2000. Expanded polyglutamine peptides alone are intrinsically cytotoxic and cause neurodegeneration in *Drosophila*. *Hum Mol Genet.* 9:13-25.
- Martindale, D., A. Hackam, A. Wieczorek, L. Ellerby, C. Wellington, K. McCutcheon, R. Singaraja, P. Kazemi-Esfarjani, R. Devon, S.U. Kim, D.E. Bredesen, F. Tufaro und M.R. Hayden. 1998. Length of Huntingtin and its polyglutamine tract influences localization und frequency of intracellular aggregates. *Nat Genet.* 18:150-4.
- Matilla, A., B.T. Koshy, C.J. Cummings, T. Isobe, H.T. Orr und H.Y. Zoghbi. 1997. The cerebellar leucine-rich acidic nuclear protein interacts with ataxin-1 [published erratum appears in *Nature* 1998 Feb 19;391(6669):818]. *Nature.* 389:974-8.
- McEvelly, R.J., M.O. de Diaz, M.D. Schonemann, F. Hooshmand und M.G. Rosenfeld. 2002. Transcriptional regulation of cortical neuron migration by POU domain factors. *Science.* 295:1528-32.
- Meriin, A., K. Mabuchi, V. Gabai, J. Yaglom, A. Kazantsev und M. Sherman. 2001. Intracellular aggregation of polypeptides with expanded polyglutamine domain is stimulated by stress-activated kinase MEKK1. *Journal of Cell Biology.* 153:851-64.
- Merz, P.A., R.A. Somerville, H.M. Wisniewski und K. Iqbal. 1981. Abnormal fibrils from scrapie-infected brain. *Acta Neuropathol.* 54:63-74.

- Merz, P.A., H.M. Wisniewski, R.A. Somerville, S.A. Bobin, C.L. Masters und K. Iqbal. 1983. Ultrastructural morphology of amyloid fibrils from neuritic and amyloid plaques. *Acta Neuropathol.* 60:113-24.
- Metzler, M., C.D. Helgason, I. Dragatsis, T. Zhang, L. Gan, N. Pineault, S.O. Zeitlin, R.K. Humphries und M.R. Hayden. 2000. Huntingtin is required for normal hematopoiesis. *Hum Mol Genet.* 9:387-94.
- Minton, A.P. 2001. The influence of macromolecular crowding and macromolecular confinement on biochemical reactions in physiological media. *J Biol Chem.* 276:10577-80.
- Muchowski, P.J., K. Ning, C. D'Souza-Schorey und S. Fields. 2002. Requirement of an intact microtubule cytoskeleton for aggregation and inclusion body formation by a mutant Huntingtin fragment. *Proc Natl Acad Sci U S A.* 99:727-32.
- Muchowski, P.J., G. Schaffar, A. Sittler, E.E. Wanker, M.K. Hayer-Hartl und F.U. Hartl. 2000. Hsp70 and hsp40 chaperones can inhibit self-assembly of polyglutamine proteins into amyloid-like fibrils. *Proc Natl Acad Sci U S A.* 97:7841-6.
- Nagafuchi, S., H. Yanagisawa, K. Sato, T. Shirayama, E. Ohsaki, M. Bundo, T. Takeda, K. Tadokoro, I. Kondo, N. Murayama, et al. 1994. Dentatorubral and pallidoluysian atrophy expansion of an unstable CAG trinucleotide on chromosome 12p. *Nat Genet.* 6:14-8.
- Nakamura, K., S.Y. Jeong, T. Uchihara, M. Anno, K. Nagashima, T. Nagashima, S. Ikeda, S. Tsuji und I. Kanazawa. 2001. SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. *Hum Mol Genet.* 10:1441-8.
- Narain, Y., A. Wytttenbach, J. Rankin, R.A. Furlong und D.C. Rubinsztein. 1999. A molecular investigation of true dominance in Huntington's disease. *J Med Genet.* 36:739-46.
- Nasir, J., S.B. Floresco, J.R. O'Kusky, V.M. Diewert, J.M. Richman, J. Zeisler, A. Borowski, J.D. Marth, A.G. Phillips und M.R. Hayden. 1995. Targeted disruption of the Huntington's disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes. *Cell.* 81:811-23.
- Notterpek, L., M.C. Ryan, A.R. Tobler und E.M. Shooter. 1999. PMP22 accumulation in aggresomes: implications for CMT1A pathology. *Neurobiol Dis.* 6:450-60.

- Nokazaki, K., O. Onodera, H. Tanako und S. Tsuji . 2001. Amino acid sequences flanking polyglutamine stretches influence their potential aggregates formation. *Neuroreport* 12, 15:3357-3364
- Nucifora, F.C., M. Sasaki, M.F. Peters, H. Huang, J.K. Cooper, M. Yamada, H. Takahashi, S. Tsuji, J. Troncoso, V.L. Dawson, T.M. Dawson und C.A. Ross. 2001. Interference by Huntingtin and atrophin-1 with CBP-mediated transcription leading to cellular toxicity. *Science*. 291:2423-8.
- O'Kusky, J.R., J. Nasir, F. Cicchetti, A. Parent und M.R. Hayden. 1999. Neuronal degeneration in the basal ganglia and loss of pallido- subthalamic synapses in mice with targeted disruption of the Huntington's disease gene. *Brain Res*. 818:468-79.
- Ona, V.O., M. Li, J.P. Vonsattel, L.J. Andrews, S.Q. Khan, W.M. Chung, A.S. Frey, A.S. Menon, X.J. Li, P.E. Stieg, J. Yuan, J.B. Penney, A.B. Young, J.H. Cha und R.M. Friedlander. 1999. Inhibition of caspase-1 slows disease progression in a mouse model of Huntington's disease [see comments]. *Nature*. 399:263-7.
- Oostra, B.A. und P. Chiurazzi. 2001. The fragile X gene and its function. *Clin Genet*. 60:399-408.
- Ordway, J.M., S. Tallaksen-Greene, C.A. Gutekunst, E.M. Bernstein, J.A. Cearley, H.W. Wiener, L.S.t. Dure, R. Lindsey, S.M. Hersch, R.S. Jope, R.L. Albin und P.J. Detloff. 1997. Ectopically expressed CAG repeats cause intranuclear inclusions and a progressive late onset neurological phenotype in the mouse. *Cell*. 91:753-63.
- Paulson, H.L., M.K. Perez, Y. Tottier, J.Q. Trojanowski, S.H. Subramony, S.S. Das, P. Vig, J.-L. Mandel, K.H. Fischbeck und R.N. Pittman. 1997. Intranuclear inclusions of expanded polyglutamine protein in spinocerebellar ataxia type 3. *Neuron*. 19:333-44.
- Pawson, T. und G.D. Gish. 1992. SH2 and SH3 domains: From structure to function. *Cell*. 71:359-62.
- Perez, M.K., H.L. Paulson, S.J. Pendse, S.J. Saionz, N.M. Bonini und R.N. Pittman. 1998. Recruitment and the role of nuclear localization in polyglutamine- mediated aggregation. *J Cell Biol*. 143:1457-70.

- Perutz, M.F. 1999. Glutamine repeats and neurodegenerative diseases. *Brain Res Bull.* 50:467.
- Perutz, M.F., T. Johnson, M. Suzuki und J.T. Finch. 1994. Glutamine repeats as polar zippers: their possible role in inherited neurodegenerative diseases. *Proc Natl Acad Sci U S A.* 91:5355-8.
- Perutz, M.F., B.J. Pope, D. Owen, E.E. Wanker und E. Scherzinger. 2002. Aggregation of proteins with expanded glutamine and alanine repeats of the glutamine-rich and asparagine-rich domains of Sup35 and of the amyloid beta-peptide of amyloid plaques. *Proc Natl Acad Sci U S A.* 99:5596-600.
- Prusiner, S.B., M.R. Scott, S.J. DeArmond und F.E. Cohen. 1998. Prion protein biology. *Cell.* 93:337-48.
- Rajan, R.S., M.E. Illing, N.F. Bence und R.R. Kopito. 2001. Specificity in intracellular protein aggregation and inclusion body formation. *Proc Natl Acad Sci U S A.* 98:13060-5.
- Rigamonti, D., J.H. Bauer, C. De-Fraja, L. Conti, S. Sipione, C. Sciorati, E. Clementi, A. Hackam, M.R. Hayden, Y. Li, J.K. Cooper, C.A. Ross, S. Govoni, C. Vincenz und E. Cattaneo. 2000. Wild-type Huntingtin protects from apoptosis upstream of caspase-3. *J Neurosci.* 20:3705-13.
- Roizin, L., S. Stellar, N. Willson, J. Whittier und J.C. Liu. 1974. Electron microscope and enzyme studies in cerebral biopsies of Huntington's chorea. *Trans Am Neurol Assoc.* 99:240-3.
- Ross, C.A. 1997. Intranuclear neuronal inclusions: a common pathogenic mechanism for glutamine-repeat neurodegenerative diseases? *Neuron.* 19:1147-50.
- Ross, T.S., O.A. Bernard, R. Berger und D.G. Gilliland. 1998. Fusion of Huntingtin interacting protein 1 to platelet-derived growth factor beta receptor (PDGFbetaR) in chronic myelomonocytic leukemia with t(5;7)(q33;q11.2). *Blood.* 91:4419-26.
- Rubinsztein, D.C. 2002. Lessons from animal models of Huntington's disease. *Trends Genet.* 18:202-9.
- Rubinsztein, D.C., J. Leggo, R. Coles, E. Almqvist, V. Biancalana, J.J. Cassiman, K. Chotai, M. Connarty, D. Crauford, A. Curtis, D. Curtis, M.J. Davidson, A.M. Differ, C. Dode, A. Dodge, M. Frontali, N.G. Ranen, O.C. Stine, M. Sherr, M.H. Abbott, M.L. Franz, C.A. Graham, P.S. Harper, J.C. Hedreen, M.R. Hayden und

- et al. 1996. Phenotypic characterization of individuals with 30-40 CAG repeats in the Huntington disease (HD) gene reveals HD cases with 36 repeats and apparently normal elderly individuals with 36-39 repeats. *Am J Hum Genet.* 59:16-22.
- Sanpei, K., H. Takano, S. Igarashi, T. Sato, M. Oyake, H. Sasaki, A. Wakisaka, K. Tashiro, Y. Ishida, T. Ikeuchi, R. Koide, M. Saito, A. Sato, T. Tanaka, S. Hanyu, Y. Takiyama, M. Nishizawa, N. Shimizu, Y. Nomura, M. Segawa, K. Iwabuchi, I. Eguchi, H. Tanaka, H. Takahashi und S. Tsuji. 1996. Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT [see comments]. *Nat Genet.* 14:277-84.
- Sathasivam, K., I. Amaechi, L. Mangiarini und G. Bates. 1997. Identification of an HD patient with a (CAG)180 repeat expansion and the propagation of highly expanded CAG repeats in lambda phage. *Hum Genet.* 99:692-5.
- Satyal, S.H., E. Schmidt, K. Kitagawa, N. Sondheimer, S. Lindquist, J.M. Kramer und R.I. Morimoto. 2000. Polyglutamine aggregates alter protein folding homeostasis in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A.* 97:5750-5.
- Saudou, F., S. Finkbeiner, D. Devys und M.E. Greenberg. 1998. Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions. *Cell.* 95:55-66.
- Sawa, A. 2001. Mechanisms for neuronal cell death and dysfunction in Huntington's disease: pathological cross-talk between the nucleus and the mitochondria? *J Mol Med.* 79:375-81.
- Scherzinger, E., R. Lurz, M. Turmaine, L. Mangiarini, B. Hollenbach, R. Hasenbank, G.P. Bates, S.W. Davies, H. Lehrach und E.E. Wanker. 1997. Huntingtin-encoded polyglutamine expansions form amyloid-like protein aggregates in vitro and in vivo. *Cell.* 90:549-58.
- Scherzinger, E., A. Sittler, K. Schweiger, V. Heiser, R. Lurz, R. Hasenbank, G.P. Bates, H. Lehrach und E.E. Wanker. 1999. Self-assembly of polyglutamine-containing Huntingtin fragments into amyloid-like fibrils: implications for Huntington's disease pathology. *Proc Natl Acad Sci U S A.* 96:4604-9.
- Schilling, G., M.W. Becher, A.H. Sharp, H.A. Jinnah, K. Duan, J.A. Kotzuk, H.H. Slunt, T. Ratovitski, J.K. Cooper, N.A. Jenkins, N.G. Copeland, D.L. Price, C.A. Ross

- und D.R. Borchelt. 1999. Intranuclear inclusions and neuritic aggregates in transgenic mice expressing a mutant N-terminal fragment of Huntingtin [published erratum appears in *Hum Mol Genet* 1999 May;8(5):943]. *Hum Mol Genet.* 8:397-407.
- Schonemann, M.D., A.K. Ryan, R.J. McEvilly, S.M. O'Connell, C.A. Arias, K.A. Kalla, P. Li, P.E. Sawchenko und M.G. Rosenfeld. 1995. Development and survival of the endocrine hypothalamus and posterior pituitary gland requires the neuronal POU domain factor Brn-2. *Genes Dev.* 9:3122-35.
- Schreiber, E., K. Harshman, I. Kemler, U. Malipiero, W. Schaffner und A. Fontana. 1990. Astrocytes and glioblastoma cells express novel octamer-DNA binding proteins distinct from the ubiquitous Oct-1 and B cell type Oct-2 proteins. *Nucleic Acids Res.* 18:5495-503.
- Schreiber, E., A. Tobler, U. Malipiero, W. Schaffner und A. Fontana. 1993. cDNA cloning of human N-Oct3, a nervous-system specific POU domain transcription factor binding to the octamer DNA motif. *Nucleic Acids Res.* 21:253-8.
- Sharp, A.H., S.J. Loev, G. Schilling, S.-H. Li, X.-J. Li, J. Bao, M.V. Wagster, J.A. Kotzuk, J.P. Steiner, A. Lo, J. Hedreen, S. Sisodia, S.H. Snyder, T.M. Dawson, D.K. Ryugo und C.A. Ross. 1995. Widespread expression of Huntington's disease gene (IT15) protein product. *Neuron.* 14:1065-74.
- Sharp, A.H. und C.A. Ross. 1996. Neurobiology of Huntington's disease. *Neurobiol Dis.* 3:3-15.
- Sieradzan, K., D.M. Mann und A. Dodge. 1997. Clinical presentation and patterns of regional cerebral atrophy related to the length of trinucleotide repeat expansion in patients with adult onset Huntington's disease. *Neurosci Lett.* 225:45-8.
- Sieradzan, K.A., A.O. Mehan, L. Jones, E.E. Wanker, N. Nukina und D.M. Mann. 1999. Huntington's disease intranuclear inclusions contain truncated, ubiquitinated Huntingtin protein. *Exp Neurol.* 156:92-9.
- Sittler, A., S. Walter, N. Wedemeyer, R. Hasenbank, E. Scherzinger, H. Eickhoff, G.P. Bates, H. Lehrach und E.E. Wanker. 1998. SH3GL3 associates with the Huntingtin exon 1 protein and promotes the formation of polyglu-containing protein aggregates. *Mol Cell.* 2:427-36.

- Skinner, P.J., B.T. Koshy, C.J. Cummings, I.A. Klement, K. Helin, A. Servadio, H.Y. Zoghbi und H.T. Orr. 1997. Ataxin-1 with an expanded glutamine tract alters nuclear matrix-associated structures [published erratum appears in Nature 1998 Jan 15;391(6664):307]. *Nature*. 389:971-4.
- Smith, D.B. und K.S. Johnson. 1988. Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene*. 67:31-40.
- Snell, R.G., L.A. Doucette-Stamm, K.M. Gillespie, S.A. Taylor, L. Riba, G.P. Bates, M.R. Altherr, M.E. MacDonald, J.F. Gusella, J.J. Wasmuth, et al. 1993. The isolation of cDNAs within the Huntington disease region by hybridisation of yeast artificial chromosomes to a cDNA library. *Hum Mol Genet*. 2:305-9.
- Speed, M.A., D.I. Wang und J. King. 1996. Specific aggregation of partially folded polypeptide chains: the molecular basis of inclusion body composition. *Nat Biotechnol*. 14:1283-7.
- Spillantini, M.G., R.A. Crowther, R. Jakes, M. Hasegawa und M. Goedert. 1998. alpha-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with lewy bodies. *Proc Natl Acad Sci U S A*. 95:6469-73.
- Steffan, J.S., A. Kazantsev, O. Spasic-Boskovic, M. Greenwald, Y.Z. Zhu, H. Gohler, E.E. Wanker, G.P. Bates, D.E. Housman und L.M. Thompson. 2000. The Huntington's disease protein interacts with p53 and CREB-binding protein and represses transcription. *Proc Natl Acad Sci U S A*. 97:6763-8.
- Sugihara, T.M., E.I. Kudryavtseva, V. Kumar, J.J. Horridge und B. Andersen. 2001. The POU domain factor Skin-1a represses the keratin 14 promoter independent of DNA binding. A possible role for interactions between Skn-1a and CREB-binding protein/p300. *J Biol Chem*. 276:33036-44.
- Suhr, S.T., M.C. Senut, J.P. Whitelegge, K.E. Faull, D.B. Cuizon und F.H. Gage. 2001. Identities of sequestered proteins in aggregates from cells with induced polyglutamine expression. *Journal of Cell Biology*. 153:283-94.
- Tanzi, R.E. 1998. Of calcium, caspases, and cognitive decline. *Nat Med*. 4:1127-8.
- Thomson, J.A., K. Murphy, E. Baker, G.R. Sutherland, P.G. Parsons, R.A. Sturm und F. Thomson. 1995. The brn-2 gene regulates the melanocytic phenotype and tumorigenic potential of human melanoma cells. *Oncogene*. 11:691-700.

- Tilley, W.D., M. Marcelli, J.D. Wilson und M.J. McPhaul. 1989. Characterization and expression of a cDNA encoding the human androgen receptor. *Proc Natl Acad Sci U S A.* 86:327-31.
- Timasheff, S.N. 1993a. The control of protein stability and association by weak interactions with water: how do solvents affect these processes? *Annu Rev Biophys Biomol Struct.* 22:67-97.
- Timasheff, S.N. 1993b. Protein-Protein -Interactions.pp. 315-36.
- Trojanowski, J.Q., M. Goedert, T. Iwatsubo und V.M. Lee. 1998. Fatal attractions: abnormal protein aggregation and neuron death in Parkinson's disease and Lewy body dementia [see comments]. *Cell Death Differ.* 5:832-7.
- Trottier, Y., D. Devys, G. Imbert, F. Saudou, I. An, Y. Lutz, C. Weber, Y. Agid, E.C. Hirsch und J.L. Mandel. 1995. Cellular localization of the Huntington's disease protein and discrimination of the normal and mutated form [see comments]. *Nat Genet.* 10:104-10.
- Velier, J., M. Kim, C. Schwarz, T.W. Kim, E. Sapp, K. Chase, N. Aronin und M. DiFiglia. 1998. Wild-type and mutant Huntingtins function in vesicle trafficking in the secretory and endocytic pathways. *Exp Neurol.* 152:34-40.
- Vonsattel, J.P., R.H. Myers, T.J. Stevens, R.J. Ferrante, E.D. Bird und E.P. Richardson, Jr. 1985. Neuropathological classification of Huntington's disease. *J Neuropathol Exp Neurol.* 44:559-77.
- Waelter, S., A. Boeddrich, R. Lurz, E. Scherzinger, G. Lueder, H. Lehrach und E.E. Wanker. 2001a. Accumulation of mutant Huntingtin fragments in aggresome-like inclusion bodies as a result of insufficient protein degradation. *Mol Biol Cell.* 12:1393-407.
- Waelter, S., E. Scherzinger, R. Hasenbank, E. Nordhoff, R. Lurz, H. Goehler, C. Gauss, K. Sathasivam, G.P. Bates, H. Lehrach und E.E. Wanker. 2001b. The Huntingtin interacting protein HIP1 is a clathrin and alpha-adaptin- binding protein involved in receptor-mediated endocytosis. *Hum Mol Genet.* 10:1807-17.
- Walker. 1994. *BIO/TECHNOLOGY.* 12:601.
- Wanker, E.E. 2000a. Protein aggregation and pathogenesis of Huntington's disease: mechanisms and correlations [In Process Citation]. *Biol Chem.* 381:937-42.

- Wanker, E.E. 2000b. Protein aggregation in Huntington's and Parkinson's disease: implications for therapy [In Process Citation]. *Mol Med Today*. 6:387-91.
- Wanker, E.E., C. Rovira, E. Scherzinger, R. Hasenbank, S. Walter, D. Tait, J. Colicelli und H. Lehrach. 1997. HIP-I: a Huntingtin interacting protein isolated by the yeast two- hybrid system. *Hum Mol Genet*. 6:487-95.
- Wanker, E.E., E. Scherzinger, V. Heiser, A. Sittler, H. Eickhoff und H. Lehrach. 1999. Membrane filter assay for detection of amyloid-like polyglutamine- containing protein aggregates. *Methods Enzymol*. 309:375-86.
- Waragai, M., C.H. Lammers, S. Takeuchi, I. Imafuku, Y. Udagawa, I. Kanazawa, M. Kawabata, M.M. Mouradian und H. Okazawa. 1999. PQBP-1, a novel polyglutamine tract-binding protein, inhibits transcription activation by Brn-2 and affects cell survival. *Hum Mol Genet*. 8:977-87.
- Wellington, C.L. und M.R. Hayden. 2000. Caspases and neurodegeneration: on the cutting edge of new therapeutic approaches. *Clin Genet*. 57:1-10.
- Wellington, C.L., R. Singaraja, L. Ellerby, J. Savill, S. Roy, B. Leavitt, E. Cattaneo, A. Hackam, A. Sharp, N. Thornberry, D.W. Nicholson, D.E. Bredesen und M.R. Hayden. 2000. Inhibiting caspase cleavage of Huntingtin reduces toxicity and aggregate formation in neuronal and nonneuronal cells. *J Biol Chem*.
- Wetzel, R. 1996. For protein misassembly, it's the "I" decade. *Cell*. 86:699-702.
- Wheeler, V.C., J.K. White, C.A. Gutekunst, V. Vrbanac, M. Weaver, X.J. Li, S.H. Li, H. Yi, J.P. Vonsattel, J.F. Gusella, S. Hersch, W. Auerbach, A.L. Joyner und M.E. MacDonald. 2000. Long glutamine tracts cause nuclear localization of a novel form of Huntingtin in medium spiny striatal neurons in HdhQ92 and HdhQ111 knock- in mice. *Hum Mol Genet*. 9:503-13.
- White, J.K., W. Auerbach, M.P. Duyao, J.P. Vonsattel, J.F. Gusella, A.L. Joyner und M.E. MacDonald. 1997. Huntingtin is required for neurogenesis and is not impaired by the Huntington's disease CAG expansion. *Nat Genet*. 17:404-10.
- Wood, S.J., J. Wypych, S. Steavenson, J.-C. Louis, M. Citron und A.L. Biere. 1999. α -Synuclein fibrillogenesis is nucleation-dependent. *J. Biol. Chem*. 274:19509-15.
- Yamamoto, A., J.J. Lucas und R. Hen. 2000. Reversal of neuropathology and motor dysfunction in a conditional model of Huntington's disease. *Cell*. 101:57-66.

- Zeitlin, S., J.P. Liu, D.L. Chapman, V.E. Papaioannou und A. Efstratiadis. 1995. Increased apoptosis and early embryonic lethality in mice nullizygous for the Huntington's disease gene homologue. *Nat Genet.* 11:155-63.
- Zhuchenko, O., J. Bailey, P. Bonnen, T. Ashizawa, D.W. Stockton, C. Amos, W.B. Dobyns, S.H. Subramony, H.Y. Zoghbi und C.C. Lee. 1997. Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the alpha 1A-voltage-dependent calcium channel. *Nat Genet.* 15:62-9.
- Zuccato, C., A. Ciammola, D. Rigamonti, B.R. Leavitt, D. Goffredo, L. Conti, M.E. MacDonald, R.M. Friedlander, V. Silani, M.R. Hayden, T. Timmusk, S. Sipione und E. Cattaneo. 2001. Loss of Huntingtin-mediated BDNF gene transcription in Huntington's disease. *Science.* 293:493-8.