On a theorem by do Carmo and Dajczer

G. Haak*
Fachbereich Mathematik
TU Berlin
D-10623 Berlin

Abstract

We give a new proof of a theorem by M.P. do Carmo and M. Dajczer on helicoidal surfaces of constant mean curvature.

1 Introduction

Let G be a one-parameter group of proper Euclidean motions of $\mathbb{R}^3$ of the form

$$g_t(x, y, z) = (x \cos t + y \sin t, -x \sin t + y \cos t, z + ht), t \in \mathbb{R}.$$

I.e., G is a group of helicoidal transformations with pitch $h \in \mathbb{R}$. In the degenerate case $h = 0$, G becomes a group of pure rotations. Up to an affine change of coordinates and reparametrization, all one-parameter groups of Euclidean motions are either of this form or are groups of pure translations.

In 1982 do Carmo and Dajczer [5] investigated surfaces of constant mean curvature (CMC-surfaces) which are generated from a plane curve by the action of a helicoidal group in the same way as a rotational surface is generated by the action of a group of rotations. They proved the following theorem:

Theorem 1: *A complete immersed CMC-surface is helicoidal if and only if it is in the associated family of a Delaunay surface.*

They proved this result by introducing for each helicoidal CMC-immersion the 2-parameter family of helicoidal surfaces given by Bour's Lemma [2] and evaluating the constant curvature condition for the elements of these families. This approach on one hand gives an explicit parametrization of helicoidal CMC-immersions. On the other hand, it reaches its goal, the proof of Theorem 1, in a fairly indirect way.

Since helicoidal surfaces still spawn interest [4, 7], we want to show in this note how Theorem 1 can be obtained in a much simpler way using a more recent theorem of Smyth [8] and some results from [3]. We state Smyth's theorem in the language of [3]:

*supported by Sonderforschungsbereich 288

Theorem 2: *Let $\Phi : M \to \mathbb{R}^3$, M a Riemann surface, be a complete conformally immersed CMC-surface admitting a one-parameter group of self-isometries. Then the simply connected cover of M is the complex plane and the surface is either in the associated family of a Delaunay surface or its metric is rotationally invariant.*

Here a self-isometry is an automorphism of the simply connected cover $\mathcal{D}$ of M, which preserves the metric of the universal covering immersion $\Psi : \mathcal{D} \to \mathbb{R}^3$ given by pulling back the immersion Φ to $\mathcal{D}$. For details see [3]. Those CMC-surfaces which have a rotationally invariant metric are now commonly called Smyth surfaces.

We also introduce the notion of a space symmetry of a CMC-immersion $\Phi : M \to \mathbb{R}^3$. A space symmetry of Φ is a Euclidean motion in $\mathbb{R}^3$ which preserves the image of Φ as a set. The relation between space symmetries and self-isometries was also studied exhaustively in [3]. By [3, Lemma 2.15] the group of space symmetries of a Smyth surface is discrete.

2 The proof of Theorem 1

We can now give the proof of Theorem 1 right away:

Proof: For a given CMC-immersion $\Phi : M \to \mathbb{R}^3$ there exists (see e.g. [3, Theorem 2.2]) a conformal structure on M such that M becomes a Riemann surface and Φ becomes a conformal CMC-immersion. If Φ is also complete and in addition admits a one-parameter group of helicoidal space symmetries, then by [3, Prop. 2.12] and [3, Corollary 2.6], Φ admits also a one-parameter group of self-isometries. In particular it satisfies the assumptions of Theorem 2 above. Since a group of helicoidal Euclidean motions is never discrete, the surface cannot be a Smyth surface. It therefore has to be in the associated family of a Delaunay surface.

Conversely, by [3, Lemma 2.15] and [3, Prop. 3.4] each element of the associated family of a Delaunay surface admits a one-parameter group of space symmetries. Since the most general one-parameter group of Euclidean motions is a group of helicoidal transformations (with possibly degenerate pitch), all surfaces in the associated family of a Delaunay surface are helicoidal or rotational. □

It should also be noted that in the language of integrable systems (the metric of a conformal CMC-immersion without umbilics satisfies the integrable sinh-Gordon equation), Theorem 1 also implies that helicoidal CMC-surfaces are of finite type (see [6] and [1]). Thus for helicoidal surfaces, apart from the parametrizations given in [5] and [7], there is Bobenko's parametrization in terms of theta functions [1].

References

[1] A. Bobenko, *All constant mean curvature tori in R^3, S^3, H^3 in terms of theta-functions*, Math. Ann., 290 (1991), pp. 209–245.

[2] E. Bour, *Memoire sur le deformation de surfaces*, Journal de l'Ecole Polytechnique, XXXIX Cahier (1862), pp. 1–148.

[3] J. Dorfmeister and G. Haak, *On symmetries of constant mean curvature surfaces*, preprint 197, KITCS and SFB288, 1996.

[4] B. Konopelchenko and I. Taimanov, *Constant mean curvature surfaces via anintegrable dynamical system*, J. Phys. A, 29 (1996), pp. 1261–1265.

[5] M.P. do Carmo and M. Dajczer, *Helicoidal surfaces with constant mean curvature*, Tohoku Math. Journal, 34 (1982), pp. 425–435.

[6] U. Pinkall and I. Sterling, *On the classification of constant mean curvature tori*, Annals of Math., 130 (1989), pp. 407–451.

[7] T. Sasai, *On helicoidal surfaces with constant mean curvature*, Tokyo J. Math., 19 (1996), pp. 39–50.

[8] B. Smyth, *A generalization of a theorem of Delaunay on constant mean curvature surfaces*, in Statistical thermodynamics and differential geometry of microstructured materials, Springer, Berlin, Heidelberg, New York, 1993.