

7 Literaturverzeichnis

- [1] Aaslid RH, Markwalder TM, Nornes H: Noninvasive transcranial Doppler ultrasound recording of flow velocity in basal cerebral arteries. *J. Neurosurg.* 57 (1982), 769-774
- [2] Aaslid RH, Nornes H: Musical murmurs in human cerebral arteries after subarachnoid hemorrhage. *J Neurosurg* 59 (1984), 32-36
- [3] Aaslid RH, Huber P, Nornes H: Evaluation of cerebrovascular spasm with transcranial Doppler ultrasound. *J Neurosurg* 60 (1984), 37-41
- [4] Barber PA, Davis SM, Infeld B, Baird A, Donnan GA, Jolley D, Lichtenstein M: Spontaneous reperfusion after ischemic stroke is associated with improved outcome. *Stroke* 29 (1998), 2522-2528
- [5] Baumgartner RW, Regard M: Role of impaired CO₂-reactivity in the diagnosis of cerebral low flow infarcts. *J Neurol Neurosurg Psychiatry* 57 (1994), 814-817
- [6] Bennett SA, Tenniswood M, Chen JH, Davidson CM, Keyes MT, Fortin T: Chronic cerebral hypoperfusion elicits neuronal apoptosis and behavioural impairment. *Neuroreport* 9 (1998), 161-166
- [7] Bishop CCR, Powell S, Rutt D, Browse NL: Transcranial Doppler measurements of middle cerebral artery blood flow velocity: validation study. *Stroke* 17 (1986), 913-915
- [8] Bornstein RA, Starling RC, Myerowitz PD, Haas GJ: Neuropsychological function in patients with end-stage heart failure before and after cardiac transplantation. *Acta Neurol Scand* 91 (1995), 260-265
- [9] Bradack GB, Simon RS, Heitsieck CH: Angiographically verified transient alteration of the intracranial arteries and veins in dependence on different CO₂ tension. *Neuroradiology* 10 (1976), 257-262
- [10] Brockenbrough EC: Periorbital Doppler velocity evaluation of carotid obstruction. In: Bernstein EF (e.d.): *Non-invasive Diagnostic Techniques Vascular Disease*. Mosby, St. Louis, 1968

- [11] Bullock R, Mendelow AD, Bone I, Patterson J, Macleod WN, Allardice G: Cerebral blood flow and CO₂ reponsiveness as an indicator of collateral reserve capacity in patient with carotid arterial disease. *Br J Surg* 72 (1985), 348-351
- [12] Cacciatore F, Abete P, Ferrara N, Calabrese C, Napoli C, Maggi S, Varicchio M: Congestive heart failure and cognitive impairment in an older population. Osservatorio Geriatrico Campano Study Group. *J Am Geriatr Soc* 46 (1998), 1343-1348
- [13] Dahl A, Russell D, Nyberg-Hansen R, Rootwelt K: Effect of nitroglycerin on cerebral circulation measured by transcranial Doppler and SPECT. *Stroke* 20 (1989), 1733-1736
- [14] Dahl A, Lindegaard KF, Russell D, Nyberg-Hansen R, Rootwelt K, Sorteberg W: A comparison of transcranial Doppler and cerebral blood flow studies to assess cerebral vasoreactivity. *Stroke* 23 (1992); 15-19
- [15] Dahl A, Russell D, Rootwelt K, Nyberg-Hansen R, Kerty E: Cerebral vasoreactivity assessed with transcranial Doppler and regional cerebral blood flow measurements. *Stroke* 26 (1995), 2302-2306
- [16] Diehl RR, Henkes H, Nahser HC, Kühne d, Berlit P: Blood flow velocity and vasomotor reactivity in patients with arteriovenous malformations: a transcranial Doppler study. *Stroke* 25 (1994), 1574-1580
- [17] Diehl RR, Berlit P: Funktionelle Dopplersonographie in der Neurologie. Springer, Berlin Heidelberg, 1996, S. 32-39
- [18] Doppler C: Über das farbige Licht der Doppelsterne und einige andere Gestirne des Himmels. *Abh Kgl Böhm Ge d Wissensch*, Prag 1843, 465-482
- [19] Droste DW, Dittrich R, Hermes S, Kem ny V, Schulte-Altedorneburg G, Hansberg T, Ringelstein EB: Four-gated transcranial Doppler ultrasound in the detection of circulating microemboli. *Eur J Ultrasound* 9 (1999), 117-125
- [20] Fiermonte G, Pierelli F, Pauri F, Cosentino FII, Soccorsi R, Giacomini P: Cerebrovascular CO₂ reactivity in migraine with aura and without aura. A transcranial Doppler study. *Acta Neurol Scand* 92 (1995), 166-169

- [21] Franklin DL, Schlegel W, Rushmer RF: Blood flow measured by Doppler frequency shift of back-scattered ultrasound. *Science* 132 (1961), 564-565
- [22] Georgiadis D, Baumgartner RW, Karatschai R, Lindner A, Zerkowski HR: Further evidence of gaseous embolic material in patients with artificial heart valves. *J Thorac Cardiovasc Surg* 115 (1998), 808-810
- [23] Goebell H, Wagner J: *Innere Medizin*. de Gruyter, Berlin, New York, 1992, S. 38-43
- [24] Grimm M, Yeganehfar W, Laufer G, Madl C, Kramer I, Eisenhuber E, Simon P, Kupilik N, Schreiner W, Pacher R, Bunzel B, Wolner E, Grimm E: Cyclosporine may affect improvement of cognitive brain function after successful cardiac transplantation. *Circulation* 94 (1996), 1339-1345
- [25] Grosset DG, Straiton J, Mc Donald I, Cockburn M, Bullock R: Use of transcranial Doppler sonography to predict development of a delayed ischemic deficit after subarachnoid hemorrhage. *J Neurosurg* 78 (1993), 183-187
- [26] Grosset DG, Cowburn P, Georgiadis D, Dargie HJ, Faichney A, Lee KR: Ultrasound detection of cerebral emboli in patients with prosthetic heart valves. *J Heart Valve Dis* 3 (1994), 128-132
- [27] Grosset DG, Georgiadis D, Kelman AW, Cowburn P, Stirling S, Lee KR: Detection of microemboli by transcranial Doppler ultrasound. *Tex Heart Inst J* 23 (1996), 298-292
- [28] Hennerici M, Rautenberg W, Schwartz A: Transcranial Doppler ultrasound for the assessment of intracranial arterial flow velocity - Part II. Evaluation of intracranial artery disease. *Surg Neurol* 27 (1987), 523-532
- [29] Huber P, Handa J: Effect of contrast material, hyperkapnia, hyperventilation, hypertonic glucose and papaverine on the diameter of the cerebral arteries. *Invest. Radiol.* 2 (1967), 17-32
- [30] Karnik A, Valentin A, Winkler WB, Khaffaf N, Donath P, Slany J: Sex dependency of cerebrovascular CO₂ reactivity in normal subjects. *Stroke* 28 (1997), 2353-2356

- [31] Kastrup A, Dichgans J, Niemeier M, Schabet M: Changes of cerebrovascular CO₂ reactivity during normal aging. *Stroke* 29 (1998), 1311-1314
- [32] Keller HM, Baumgartner G, Brunner HH, Regli F: Karotisstenosen: Diagnose durch perkutane Messung mit dem Doppler-Ultraschall-Prinzip an der A. Supraorbitalis und A. frontalis medialis. K. Prenner (Hrsg.): *Die Chirurgie der supraaortischen Äste*. IV. Jahr
- [33] Keller HM, Baumgartner G: Karotisstenosen und -okklusionen. Diagnose durch perkutane Ultraschall-Doppler-Sonographie an der A. supraorbitalis und A. supratrochlearis. *Dtsch Med Wschr* 98 (1973), 1691-1698
- [34] Kleiser B, Widder B: Course of carotid artery occlusions with impaired cerebrovascular reactivity. *Stroke* 23 (1992), 171-174
- [35] Kofke WA, Brauer P, Polycare R, Panthany S, Barker D, Horton J: Middle cerebral artery blood flow velocity and stable Xenon-enhanced Computed Tomographic blood flow during balloon test occlusion of the internal carotid artery. *Stroke* 26 (1995), 1603-1606
- [36] Kontos HA, Wei EP, Nawari RM, Levasseur JE, Rosenblum WI, Patterson JL: Responses of cerebral arteries and arterioles to acute hypotension and hypertension. *Am J Physiol* 234 (1978), 371-383
- [37] Kontos HA, Wei EP, Dietrich WD, Nawari RM, Povlishock JT, Ghatak NR: Mechanism of cerebral arteriolar abnormalities after acute hypertension. *Am J Physiol* 240 (1981), 511-527
- [38] Larsen FS, Olsen KS, Hansen BA, Paulson OB, Knutsen GM: Transcranial Doppler is valid for determination of the lower limit of cerebral blood flow autoregulation. *Stroke* 25 (1994), 1985-1988
- [39] Lindegaard KF, Bakke SJ, Grolimund P, Aaslid R, Huber P, Nornes H: Assessment of intracranial hemodynamics in carotid artery disease by transcranial Doppler ultrasound. *J Neurosurg* 63 (1985), 890-898
- [40] Lindegaard KF, Grolimund P, Aaslid RF, Nornes H: Evaluation of cerebral AVM's using transcranial Doppler ultrasound. *J Neurosurg* 64 (1986), 335-344

- [41] Lindegaard KF, Bakke SJ, Aaslid RF, Nornes H: Doppler diagnosis of intracranial artery occlusive disorders. *J Neurol Neurosurg Psychiatry* 49 (1986), 510-518
- [42] Lindegaard KF, Lundar T, Wiberg J, Sjoberg D, Aaslid R Nornes H: Variations in middle cerebral artery blood flow investigated with noninvasive transcranial blood velocity measurements. *Stroke* 18 (1987), 1025-1030
- [43] Manchola IF, Salles AAF, Foo TK, Ackermann RH, Candia GT, Kjellberg RN: Arteriovenous malformation hemodynamics: a transcranial Doppler study. *Neurosurg* 33 (1993), 556-562
- [44] Markus HS, Harrison MJ: Estimation of cerebrovascular reactivity using transcranial Doppler, including the use of breath-holding as the vasodilatory stimulus. *Stroke* 23 (1992), 668-673
- [45] Markus HS, Droste DW, Brown MM: Detection of asymptomatic cerebral embolic signals with Doppler ultrasound. *Lancet* 343 (1994), 1011-1012
- [46] Markus HS, Clifton A, Buckenham T, Taylor R, Brown MM: Improvement in cerebral hemodynamics after carotid angioplasty. *Stroke* 27 (1996), 612-616
- [47] Markwalder TM, Grolimund P, Seiler W, Roth F, Aaslid R: Dependency of blood flow velocity in the middle cerebral artery on end-tidal carbondioxide partial pressure - A transcranial ultrasound Doppler study. *J Cereb Blood flow Metab* 4 (1984), 368-372
- [48] Maroon JC, Pieroni DW, Campell RL: Ophthalmosonometry, an ultrasonic method for assessing carotid blood flow. *J. Neurosurg.* 30 (1969), 238-246
- [49] Mast H, Mohr JP, Thompson JLP, Osipow A, Trocio SH, Mayer S, Young WL: Transcranial Doppler ultrasonography in cerebral arteriovenous malformations: diagnostic sensivity and association of flow velocity with spontaneous hemorrhage and focal neurological deficit. *Stroke* 26 (1995), 1024-1027
- [50] Matteis M, Troisi E, Monaldo BC, Caltagirone C, Silvestrini M: Age and sex differences in cerebral hemodynamics: a transcranial Doppler study. *Stroke* 29 (1998), 963-967

- [51] Mattle H, Grolimund P, Huber P, Sturzenegger M, Zubrugg HR: Transcranial Doppler sonographic findings in middle cerebral artery disease. *Arch Neurol* 45 (1988), 289-295
- [52] Miyazaki M, Kato K: Measurement of cerebral blood flow by ultrasonic Doppler technique. Hemodynamic correlation of internal carotid artery and vertebral artery. *Jap Circulat J* 30 (1966), 981-985
- [53] Miyazaki M: Measurement of cerebral blood flow by ultrasonic Doppler technique. *Jap Circulat J* 29 (1965), 383-386
- [54] Mortara A, Bernadi L, Pinna GD, Spadacini G, Maestri R, Dambacher M: Alterations of breathing in chronic heart failure: clinical relevance of arterial oxygen saturation instability. *Clin Sci (Colch)* 91 (1996), 72-74
- [55] Müller HR: Direktionale Doppler-Sonographie der Arteria frontalis medialis. *EEG/EMG* 2 (1971), 24-32
- [56] Müller M, Voges M, Piepgras U, Schimrigk K: Assessment of cerebral vasomotor reactivity by transcranial Doppler ultrasound and breath-holding. A comparison with azetazolamid as vasodilatory stimulus. *Stroke* 26 (1995), 96-100
- [57] Müller M, Hermes M, Bruckmann H, Schimrigk K: Transcranial Doppler ultrasound in the evaluation of collateral blood flow in patients with internal carotid artery occlusion: correlaton with cerebral angiography. *Am J Neuroradiol* 16 (1995), 195-202
- [58] Ni J, Ohta H, Matsumoto K, Watanabe H: Progressive cognitive impairment following chronic cerebral hypoperfusion induced by permanent occlusion of bilateral carotid arteries in rats. *Brain Res* 653 (1994), 231-236
- [59] Ohta H, Nishikawa H, Kimura H, Anayama H, Miyamoto M: Chronic cerebral hypoperfusion by permanent internal carotid ligation produces learning impairment without brain damage in rat. *Neuroscience* 79 (1997), 1039-1050

- [60] Pappas BA, de la Torre JC, Davidson CM, Keyes MT, Fortin T: Chronic reduction of cerebral blood flow in the adult rat: late-emerging CA1 cell loss and memory dysfunction. *Brain Res* 708 (1996), 50-58
- [61] Paulson OB, Strandgaard S, Edvinsson L: Cerebral autoregulation. *Cerebrovasc & Brain metabolism Rev* 2 (1990), 161-192
- [62] Piepgras A, Schmiedek, Leinsinger G, Haberl RL, Kirsch CM, Einhüpl KM: A simple test to assess cerebrovascular reserve capacity using transcranial Doppler ultrasonography and acetazolamide. *Stroke* 21 (1990), 1306-1311
- [63] Poulin MJ, Robins PA: Indexes of flow and cross-sectional area of the middle cerebral artery using Doppler ultrasound during hypoxia and hypercapnia in humans. *Stroke* 27 (1996), 2244-2250
- [64] Rath SA, Richter HP: Transcranial Doppler sonography as a reliable tool in craniocerebral trauma. *Unfallchirurg* 11 (1993), 569-575
- [65] Ringelstein EB, Zeumer K, Poeck K: Non-invasive diagnosis of intracranial lesions in the vertebrobasilar system. A comparison of Doppler sonographic and angiographic findings. *Stroke* 16 (1985), 848-855
- [66] Ringelstein EB, Zeumer K, Korbmacher G, Wulfinghoff F: Transkranielle Dopplersonographie der hirnersorgenden Arterien: atraumatische Diagnostik von Stenosen und Verschlüssen des Karotissiphons und der Arteria cerebri media. *Nervenarzt* 56 (1985), 296-306
- [67] Ringelstein EB, Sievers C, Ecker S, Schneider PA, Otis SM: Noninvasive assessment of CO₂-induced cerebral vasomotor response in normal individuals and patients with internal carotid artery occlusions. *Stroke* 19 (1988), 963-968
- [68] Ringelstein EB, Van Eyck S, Mertens I: Evaluation of cerebral vasomotor reactivity by various vasodilating stimuli: Comparison of CO₂ and acetazolamide. *J Cerebr Blood Flow Metab.* 12 (1992), 162-168
- [69] Rootwelt K, Dybevoold S, Nyberg-Hansen R, Russell D: Measurement of cerebral blood flow with Xe-133 inhalation and dynamic single photon

- emission computed tomography: normal values. *Scand J Clin Lab Invest Suppl* 184 (1986), 97-105
- [70] Rorick MB, Nichols FT, Adams RJ: Transcranial Doppler correlation with angiography in detection of intracranial stenosis. *Stroke* 25 (1994), 1931-1934
- [71] Satomura S: Study of the patterns in peripheral arteries by ultrasonic J acoust. *Soc. Jap.* 15 (1959), 151
- [72] Satomura S, Kaneko Z: Ultrasonic blood rheograph. *Proc. 3rd Intern. Conf. Med. Electronics IEE, London* 1961, 254-258
- [73] Schall RR, Petrucci RJ, Brozena SC, Cavarocchi NC, Jessup M: Cognitive function in patients with symptomatic dilated cardiomyopathy before and after cardiac transplantation. *J Am Coll Cardiol* 14 (1989), 1666-1672
- [74] Schmidt RF, Thews G: *Physiologie des Menschen*. 24. Auflage. Springer, Berlin Heidelberg, New York, 1990, S. 563
- [75] Sekhon LH, Morgan MK, Spence I, Weber NC: Chronic cerebral hypoperfusion and impaired neuronal function in rats. *Stroke* 25 (1994), 1022-1027
- [76] Shigoai T, Sato E, Tokitsu M, Hara M, Takeuchi K: Transcranial Doppler monitoring in severe brain damage: relationships between intracranial hemodynamics, brain dysfunction and outcome. *Neurol. Re* 12 (1990), 205-213
- [77] Siebler M, Sitzer M, Steinmetz H: Detection of intracranial emboli in patients with symptomatic extracranial carotid artery disease. *Stroke* 23 (1992), 1652-1654
- [78] Silvestrini M, Matteis M, Troisi E, Cupini LM, Bernadi G: Cerebrovascular reactivity in migraine with aura an without aura. *Headache* 36 (1996), 37-40
- [79] Sitzer M, Sohngen D, Siebler M, Specker C, Rademacher J, Janda I, Aulich A, Steinmetz H: Cerebral microembolism in patients with Sneddon's syndrome. *Arch Neurol* 52 (1995), 271-275

- [80] Sliwka U, Lingnau A, Stohlmann WD, Schmidt P, Mull M, Diehl RR, Noth J: Prevalence and time course of microembolic signals in patients with acute stroke. A prospective study. *Stroke* 28 (1997), 358-363
- [81] Sorteberg W, Langmoen IA, Lindegaard KF, Nornes H: Side-to-side differences and day-to-day variations of transcranial Doppler parameters in normal subjects. *J Ultrasound Med* 9 (1990), 403-409
- [82] Stokely EM, Sveinsdottir E, Lassen NA, Rommer P: A single photon dynamic computer assisted tomograph (DCAT) for imaging brain function in multiple cross sections. *J Comput Assist Tomograph* 4 (1980), 230-240
- [83] Sugimori H, Ibayashi S, Kenichiro F, Sadoshima S, Kuwabara Y, Fujishima M: Can transcranial Doppler really detect reduced cerebral perfusion states? *Stroke* 26 (1995), 2053-2060
- [84] Thuillez C, Annane D: Mechanisms and consequences of the vasomotor activity in cardiac insufficiency. *Arch Mal Coeur Vaiss* 84 (1991), 65-68
- [85] Wanless RB, Anand IS, Poole-Wilson PA, Harris P: An experimental model of chronic cardiac failure using adriamycin in the rabbit: central hemodynamics and regional blood flow. *Cardiovasc Res* 21 (1987), 7-13
- [86] Wardlaw JM, Is routine transcranial Doppler ultrasound monitoring useful in the management of subarachnoid hemorrhage? *J Neurosurg* 88 (1998), 272-276
- [87] Widder B: *Doppler-und Duplexsonographie der hirnersorgenden Arterien*. Springer, Berlin 1985
- [88] Widder B, Paulat K, Hackspacher J, Mayr E: Transcranial Doppler CO₂ test for the detection of hemodynamically critical artery stenoses and occlusion *Eur Arch Psychiatr Neurol Sci* 236 (1986), 162-168
- [89] Widder B: The Doppler CO₂-test to exclude patients not in need of extracranial/intracranial bypass surgery. *J Neurol Neurosurg Psychiatry* 52 (1989), 38-42
- [90] Widder B, Kleiser B, Krapf H: Course of cerebrovascular reactivity in patients with carotide artery occlusions. *Stroke* 25 (1994), 1963-1967

- [91] Zelis R, Sinoway L, Musch T, Davis D: The peripheral distribution of cardiac output in heart failure. *Z Kardiol* 77 (1988), 61-65
- [92] Zuccala G, Cattel C, Manes-Gravina E, Di Niro MG, Cocchi A, Bernabei R: Left ventricular dysfunction: a clue to cognitive impairment in older patients with heart failure. *J Neurol Neurosurg Psychiatry* 63 (1997), 509-512