

Max-Planck-Institut  
für Mathematik  
in den Naturwissenschaften  
Leipzig

Approximation of non-convex  
functionals in GBV

by

*Roberto Alicandro, Andrea Braides  
and Jayant Shah*

Preprint-Nr.: 22

1998





# Approximation of non-convex functionals in GBV

Roberto Alicandro, Andrea Braides

SISSA, via Beirut 4, 34014 Trieste, Italy

Jayant Shah

Northeastern University, Department of Mathematics

360 Huntington Avenue, Boston, MA 02115, USA

## 1 Introduction

In many problems of Computer Vision, the unknown is a pair  $(u, K)$  with  $K$  a union of (sufficiently smooth) closed curves contained in a fixed open set  $\Omega \subset \mathbf{R}^2$  and  $u : \Omega \setminus K \rightarrow \mathbf{R}$  belonging to a class of (sufficiently smooth) functions. A variational formulation of some of these problems was given by Mumford and Shah [14] introducing the functional

$$F(u, K) = \int_{\Omega \setminus K} |\nabla u|^2 dx + c_1 \mathcal{H}^1(K) + c_2 \int_{\Omega \setminus K} |u - g|^2 dx. \quad (1)$$

In this case  $g$  is interpreted as the input picture taken from a camera,  $u$  is the ‘cleaned’ image, and  $K$  is the relevant contour of the objects in the picture;  $c_1$  and  $c_2$  are contrast parameters, and  $\mathcal{H}^1(K)$  denotes the total length of  $K$ . Problems involving functionals of this form are usually called free-discontinuity problems, after a terminology introduced by De Giorgi. They have been intensively studied in recent times through weak formulations in the framework of the spaces of special functions of bounded variation (see [10], [9], [2], [5]).

The presence of the unknown surface  $K$  leads to numerical problems, and some kind of approximation of this functional is needed to obtain approximate smooth solutions. The Ambrosio and Tortorelli approach [3] provides a variational approximation of the Mumford and Shah functional (1) via elliptic functionals. The lack of convexity of the limiting functional is overcome by the introduction of an additional function variable which approaches the characteristic of the complement of the set  $K$ . The approximating functionals have the form

$$F_\varepsilon(u, v) = \int_{\Omega} v^2 |\nabla u|^2 dx + c_1 \int_{\Omega} \left( \varepsilon |\nabla v|^2 + \frac{1}{4\varepsilon} (1 - v)^2 \right) dx + c_2 \int_{\Omega} |u - g|^2 dx, \quad (2)$$

defined on functions  $u, v$  such that  $u, v \in H^1(\Omega)$  and  $0 \leq v \leq 1$ . The interaction of the terms in the second integral provide an approximate interfacial energy, as in the theory of phase transitions for Cahn - Hilliard fluids. This phenomenon had previously been described in analytic terms by Modica and Mortola [13] in the case of phase boundaries. The adaptation of the Ambrosio and Tortorelli approximation to obtain as limits more complex surface energies does not seem to follow easily from their approach.

Motivated by applications in Computer Vision and Fracture Mechanics, in this paper we study a variant of the Ambrosio Tortorelli construction by considering functionals of the form

$$G_\varepsilon(u, v) = \int_{\Omega} v^2 |\nabla u| dx + c_1 \int_{\Omega} \left( \varepsilon |\nabla v|^2 + \frac{1}{4\varepsilon} (1 - v)^2 \right) dx + c_2 \int_{\Omega} |u - g|^\gamma dx, \quad (3)$$

where  $\gamma \geq 1$ . Even though the form of these functionals is quite similar to the previous one, the domain of the limiting functional will be different. In fact, as we have  $G_\varepsilon(u, 1) \leq \int_{\Omega} |\nabla u| + c_2 |u - g|^\gamma dx$ , it is clear that the limit of these functionals will be finite if  $u \in BV(\Omega)$ . In fact we prove (Theorem 4.1 and Example 4.6) that  $G_\varepsilon$  converge to functionals related to the function-surface energy

$$G(u, K) = |Du|(\Omega \setminus K) + c_1 \int_K \frac{|u^+ - u^-|}{1 + |u^+ - u^-|} d\mathcal{H}^1 + c_2 \int_{\Omega \setminus K} |u - g|^\gamma dx, \quad (4)$$

where  $|Du|(A)$  denotes the total variation on  $A$  of the distributional derivative  $Du$ , and  $u^\pm$  are the traces of  $u$  on both sides of  $K$ . We push this approach further, constructing a variational approximation for a wide class of non-convex functionals defined on spaces of generalized functions of bounded variation.

The paper is divided as follows. In Section 2 we introduce the spaces of generalized functions of bounded variation  $GBV$  and  $GSBV$ , which are needed for a weak formulation of the functionals in (1)–(4), and the notion of  $\Gamma$ -convergence, which precises in which sense the convergence of these functionals is understood. In Section 3 we state the many preliminaries which are needed in the course of the proof. Section 4 is devoted to the statement and proof of the main result, in a slightly more general form than above. The proof of the result lies on a lower bound which is obtained by a new definition of the limit interfacial energy density, taking into account the interaction of the first two integrals of the approximating energies  $G_\varepsilon$ , and on an upper bound which is obtained by direct construction and a density result of pairs function-polyhedral surface. Section 5 contains the statement and proof of the approximation result for general isotropic functionals with convex bulk energy density and concave surface energy density defined on  $GBV$ .

## 2 Notation

We use standard notation for Sobolev and Lebesgue spaces.  $\mathcal{L}^n$  will denote the Lebesgue measure in  $\mathbf{R}^n$  and  $\mathcal{H}^k$  will denote the  $k$ -dimensional Hausdorff measure.  $\mathcal{A}(\Omega)$  and  $\mathcal{B}(\Omega)$  will be the families of open and Borel sets, respectively. If  $\mu$  is a Borel measure and  $E$  is a Borel set, then the measure  $\mu \llcorner B$  is defined as  $\mu \llcorner B(A) = \mu(A \cap B)$ . Let  $A' \subset\subset A$  be open sets. By a *cut-off function between  $A'$  and  $A$*  we mean a function  $\phi \in C_0^\infty(A)$  with  $0 \leq \phi \leq 1$  and  $\phi = 1$  on  $A'$ .

## 2.1 Generalized functions of bounded variation

Let  $u \in L^1(\Omega)$ . We say that  $u$  is a *function of bounded variation* on  $\Omega$  if its distributional derivative is a measure; i.e., there exist signed measures  $\mu_i$  such that

$$\int_{\Omega} u D_i \phi \, dx = - \int_{\Omega} \phi \, d\mu_i$$

for all  $\phi \in C_c^1(\Omega)$ . The vector measure  $\mu = (\mu_i)$  will be denoted by  $Du$ . The space of all functions of bounded variation on  $\Omega$  will be denoted by  $BV(\Omega)$ .

It can be proven that if  $u \in BV(\Omega)$  then the complement of the set of Lebesgue points  $S_u$ , that will be called the *jump set* of  $u$ , is *rectifiable*, i.e. there exists a countable family  $(\Gamma_i)$  of graphs of Lipschitz functions of  $(n-1)$  variables such that  $\mathcal{H}^{n-1}(S_u \setminus \bigcup_{i=1}^{\infty} \Gamma_i) = 0$ . Hence, a *normal*  $\nu_u$  can be defined  $\mathcal{H}^{n-1}$ -a.e. on  $S_u$ , as well as the *traces*  $u^{\pm}$  of  $u$  on both sides of  $S_u$  as

$$u^{\pm}(x) = \lim_{\rho \rightarrow 0^+} \int_{\{y \in B_{\rho}(x) : \pm \langle y-x, \nu_u(x) \rangle > 0\}} u(y) \, dy,$$

where  $\int_B u \, dy = |B|^{-1} \int_B u \, dy$ .

If  $u \in BV(\Omega)$  we define the three measures  $D^a u$ ,  $D^j u$  and  $D^c u$  as follows. By the Radon Nikodym Theorem we set  $Du = D^a u + D^s u$  where  $D^a u \ll \mathcal{L}^n$  and  $D^s u$  is the *singular part* of  $Du$  with respect to  $\mathcal{L}^n$ .  $D^a u$  is the *absolutely continuous part* of  $Du$  with respect to the Lebesgue measure,  $D^j u = Du \llcorner S_u$  is the *jump part* of  $Du$ , and  $D^c u = D^s u \llcorner (\Omega \setminus S_u)$  is the *Cantor part* of  $Du$ . We can write then

$$Du = D^a u + D^j u + D^c u.$$

It can be seen that  $D^j u = (u^+ - u^-) \nu_u \mathcal{H}^{n-1} \llcorner S_u$ , and that the Radon Nikodym derivative of  $Du$  with respect of  $\mathcal{L}^n$  is the *approximate gradient*  $\nabla u$  of  $u$ .

A function  $u \in L^1(\Omega)$  is a *special function of bounded variation* on  $\Omega$  if  $D^c u = 0$ , or, equivalently, if its distributional derivative can be written as

$$Du = \nabla u \mathcal{L}^n + (u^+ - u^-) \nu_u \mathcal{H}^{n-1} \llcorner S_u.$$

The space of special functions of bounded variation on  $\Omega$  is denoted  $SBV(\Omega)$ . We will also use the auxiliary spaces

$$SBV^p(\Omega) = \{u \in SBV(\Omega) : |\nabla u| \in L^p(\Omega), \mathcal{H}^{n-1}(S_u) < +\infty\}.$$

We define the space  $GBV(\Omega)$  of *generalized functions of bounded variation* as the space of all functions  $u \in L^1(\Omega)$  whose truncations  $u_T = (-T) \vee (u \wedge T)$  are in  $BV(\Omega)$  for any  $T > 0$ . For such functions we can define  $S_u = \bigcup_{T>0} S_{u_T}$ , and the approximate gradient and the traces  $u^{\pm}$  as the limits of the corresponding quantities defined for  $u_T$ . Moreover, we define the measure  $|D^c u| : \mathcal{B}(\Omega) \rightarrow [0, +\infty]$  as

$$|D^c u|(B) = \sup_{T>0} |D^c u_T|(B) = \lim_{T \rightarrow +\infty} |D^c u_T|(B).$$

If  $u \in BV(\Omega)$   $|D^c u|$  coincides with the usual notion of total variation of  $D^c u$ . Finally, we set

$$GSBV(\Omega) = \{u \in GBV(\Omega) : |D^c u| = 0\} = \{u \in L^1(\Omega) : u_T \in SBV(\Omega) \text{ for all } T\}.$$

For a detailed study of the properties of  $BV$ -functions we refer to [2], [11] and [12]. For an introduction to the study of free-discontinuity problems in the  $BV$  setting we refer to [2].

## 2.2 Relaxation and $\Gamma$ -convergence

Let  $(X, d)$  be a metric space. We first recall the notion of *relaxed functional*. Let  $F : X \rightarrow \mathbf{R} \cup \{+\infty\}$ . Then the relaxed functional  $\overline{F}$  of  $F$ , or *relaxation* of  $F$ , is the greatest  $d$ -lower semicontinuous functional less than or equal to  $F$ .

We say that a sequence  $F_j : X \rightarrow [-\infty, +\infty]$   $\Gamma$ -converges to  $F : X \rightarrow [-\infty, +\infty]$  (as  $j \rightarrow +\infty$ ) if for all  $u \in X$  we have

(i) (*lower limit inequality*) for every sequence  $(u_j)$  converging to  $u$

$$F(u) \leq \liminf_j F_j(u_j); \quad (5)$$

(ii) (*existence of a recovery sequence*) there exists a sequence  $(u_j)$  converging to  $u$  such that

$$F(u) \geq \limsup_j F_j(u_j), \quad (6)$$

or, equivalently by (5),

$$F(u) = \lim_j F_j(u_j). \quad (7)$$

The function  $F$  is called the  $\Gamma$ -limit of  $(F_j)$  (with respect to  $d$ ), and we write  $F = \Gamma\text{-}\lim_j F_j$ . If  $(F_\varepsilon)$  is a family of functionals indexed by  $\varepsilon > 0$  then we say that  $F_\varepsilon$   $\Gamma$ -converges to  $F$  as  $\varepsilon \rightarrow 0^+$  if  $F = \Gamma\text{-}\lim_{\varepsilon \rightarrow 0^+} F_\varepsilon$  for all  $(\varepsilon_j)$  converging to 0.

The reason for the introduction of this notion is explained by the following fundamental theorem.

**Theorem 2.1** *Let  $F = \Gamma\text{-}\lim_j F_j$ , and let a compact set  $K \subset X$  exist such that  $\inf_X F_j = \inf_K F_j$  for all  $j$ . Then*

$$\exists \min_X F = \lim_j \min_X F_j. \quad (8)$$

*Moreover, if  $(u_j)$  is a converging sequence such that  $\lim_j F_j(u_j) = \lim_j \min_X F_j$  then its limit is a minimum point for  $F$ .*

The definition of  $\Gamma$ -convergence can be given pointwise on  $X$ . It is convenient to introduce also the notion of  $\Gamma$ -lower and upper limit, as follows: let  $F_\varepsilon : X \rightarrow [-\infty, +\infty]$  and  $u \in X$ . We define

$$\Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u) = \inf \{ \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) : u_\varepsilon \rightarrow u \}; \quad (9)$$

$$\Gamma\text{-}\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u) = \inf \{ \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) : u_\varepsilon \rightarrow u \}. \quad (10)$$

If  $\Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u) = \Gamma\text{-}\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u)$  then the common value is called the  $\Gamma$ -limit of  $(F_\varepsilon)$  at  $u$ , and is denoted by  $\Gamma\text{-}\lim_{\varepsilon \rightarrow 0^+} F_\varepsilon(u)$ . Note that this definition is in accord with the previous one, and that  $F_\varepsilon$   $\Gamma$ -converges to  $F$  if and only if  $F(u) = \Gamma\text{-}\lim_{\varepsilon \rightarrow 0^+} F_\varepsilon(u)$  at all points  $u \in X$ .

We recall that:

- (i) if  $F = \Gamma\text{-}\lim_j F_j$  and  $G$  is a continuous function then  $F + G = \Gamma\text{-}\lim_j (F_j + G)$ ;
- (ii) the  $\Gamma$ -lower and upper limits define lower semicontinuous functions.

From (i) we get that in the computation of our  $\Gamma$ -limits we can drop all  $d$ -continuous terms. Remark (ii) will be used in the proofs combined with approximation arguments.

For an introduction to  $\Gamma$ -convergence we refer to [8]. For an overview of  $\Gamma$ -convergence techniques for the approximation of free-discontinuity problems see [5].

### 3 Preliminaries

In the following  $\Omega$  will denote a bounded open set in  $\mathbf{R}^n$  with Lipschitz boundary.

We denote by  $\mathcal{W}(\Omega)$  the space of all functions  $w \in SBV(\Omega)$  satisfying the following properties:

- (i)  $\mathcal{H}^{n-1}(\overline{S}_w \setminus S_w) = 0$ ;
- (ii)  $\overline{S}_w$  is the intersection of  $\Omega$  with the union of a finite number of pairwise disjoint  $(n-1)$ -dimensional simplexes;
- (iii)  $w \in W^{k,\infty}(\Omega \setminus \overline{S}_w)$  for every  $k \in \mathbf{N}$ .

The following result is due to Cortesani and Toader [7] (see also [6]).

**Theorem 3.1** (Strong approximation in  $SBV^2$ ) *Let  $u \in SBV^2(\Omega) \cap L^\infty(\Omega)$ . Then there exists a sequence  $(w_j)$  in  $\mathcal{W}(\Omega)$  such that  $w_j \rightarrow u$  strongly in  $L^1(\Omega)$ ,  $\nabla w_j \rightarrow \nabla u$  strongly in  $L^2(\Omega, \mathbf{R}^n)$ ,  $\limsup_{j \rightarrow +\infty} \|w_j\|_\infty \leq \|u\|_\infty$  and*

$$\limsup_{j \rightarrow +\infty} \int_{S_{w_j}} \phi(w_j^+, w_j^-, \nu_{w_j}) d\mathcal{H}^{n-1} \leq \int_{S_u} \phi(u^+, u^-, \nu_u) d\mathcal{H}^{n-1}$$

*for every upper semicontinuous function  $\phi : \mathbf{R} \times \mathbf{R} \times S^{n-1} \rightarrow [0, +\infty)$  such that  $\phi(a, b, \nu) = \phi(b, a, -\nu)$ , for every  $a, b \in \mathbf{R}$  and  $\nu \in S^{n-1}$ .*

The next result is a particular case of a theorem by Bouchitté, Braides and Buttazzo [4], and deals with relaxation in  $BV$  of isotropic functionals.

**Theorem 3.2** (Relaxation in  $BV$ ) *Let  $g : \mathbf{R} \rightarrow [0, +\infty]$  be a lower semicontinuous function with*

$$g(0) = 0, \quad \lim_{t \rightarrow 0^+} \frac{g(t)}{t} = 1,$$

and such that the map  $t \rightarrow g(|t|)$  is subadditive and locally bounded. Let  $F : BV(\Omega) \rightarrow [0, +\infty]$  be defined by

$$F(u) := \begin{cases} \int_{\Omega} |\nabla u| dx + \int_{S_u} g(|u^+ - u^-|) d\mathcal{H}^{n-1} & \text{if } u \in SBV^2(\Omega) \cap L^\infty(\Omega) \\ +\infty & \text{otherwise in } BV(\Omega) \end{cases}$$

Then the relaxation of  $F$  with respect to the  $L^1(\Omega)$ -topology is given on  $BV(\Omega)$  by the functional

$$\overline{F}(u) = \int_{\Omega} |\nabla u| dx + \int_{S_u} g(|u^+ - u^-|) d\mathcal{H}^{n-1} + |D^c u|(\Omega).$$

The following lemma is a commonly used tool (see [5]).

**Lemma 3.3** (Supremum of measures) *Let  $\mu : \mathcal{A}(\Omega) \rightarrow [0, +\infty)$  be an open-set superadditive function, let  $\lambda \in \mathcal{M}^+(\Omega)$ , let  $\psi_i$  be positive Borel functions such that  $\mu(A) \geq \int_A \psi_i d\lambda$  for all  $A \in \mathcal{A}(\Omega)$  and let  $\psi(x) = \sup_i \psi_i(x)$ . Then  $\mu(A) \geq \int_A \psi d\lambda$  for all  $A \in \mathcal{A}(\Omega)$ .*

We finally include a ‘slicing’ result by Ambrosio (see [1]). We introduce first some notation. Let  $\xi \in S^{n-1}$ , and let  $\Pi_\xi := \{y \in \mathbf{R}^n : \langle y, \xi \rangle = 0\}$  be the linear hyperplane orthogonal to  $\xi$ . If  $y \in \Pi_\xi$  and  $E \subset \mathbf{R}^n$  we define  $E_{\xi,y} = \{t \in \mathbf{R} : y + t\xi \in E\}$ . Moreover, if  $u : \Omega \rightarrow \mathbf{R}$  we set  $u_{\xi,y} : \Omega_{\xi,y} \rightarrow \mathbf{R}$  by  $u_{\xi,y}(t) = u(y + t\xi)$ .

**Theorem 3.4** (a) *Let  $u \in BV(\Omega)$ . Then, for all  $\xi \in S^{n-1}$  the function  $u_{\xi,y}$  belongs to  $BV(\Omega_{\xi,y})$  for  $\mathcal{H}^{n-1}$ -a.a.  $y \in \Pi_\xi$ . For such  $y$  we have*

$$u'_{\xi,y}(t) = \langle \nabla u(y + t\xi), \xi \rangle \text{ for a.a. } t \in \Omega_{\xi,y} \quad (11)$$

$$S_{u_{\xi,y}} = \{t \in \mathbf{R} : y + t\xi \in S_u\}, \quad (12)$$

$$v(t\pm) = u^\pm(y + t\xi) \quad \text{or} \quad v(t\pm) = u^\mp(y + t\xi), \quad (13)$$

according to the cases  $\langle \nu_u, \xi \rangle > 0$  or  $\langle \nu_u, \xi \rangle < 0$  (the case  $\langle \nu_u, \xi \rangle = 0$  being negligible). Moreover, we have

$$\int_{\Pi_\xi} |D^c u_{\xi,y}|(A_{\xi,y}) d\mathcal{H}^{n-1}(y) = |\langle D^c u, \xi \rangle|(A) \quad (14)$$

for all  $A \in \mathcal{A}(\Omega)$ , and for all Borel functions  $g$

$$\int_{\Pi_\xi} \sum_{t \in S_{u_{\xi,y}}} g(t) d\mathcal{H}^{n-1}(y) = \int_{S_u} g(x) |\langle \nu_u, \xi \rangle| d\mathcal{H}^{n-1}. \quad (15)$$

(b) *Conversely, if  $u \in L^1(\Omega)$  and for all  $\xi \in \{e_1, \dots, e_n\}$  and for a.e.  $y \in \Pi_\xi$   $u_{\xi,y} \in BV(\Omega_{\xi,y})$  and*

$$\int_{\Pi_\xi} |Du_{\xi,y}|(\Omega_{\xi,y}) d\mathcal{H}^{n-1}(y) < +\infty, \quad (16)$$

then  $u \in BV(\Omega)$ .



## 4 The main result

Using the space  $GBV$  defined in the previous section, it is possible to give a weak formulation for problems as in (1) and (4), which has been successfully used to obtain solutions of free-discontinuity problems (see [2]). In what follows we drop the term containing  $\int |u - g|^2 dx$ , which is of lower order, and does not affect the form of the  $\Gamma$ -limit, and we generalize the form of the functional (3).

**Theorem 4.1** *Let  $W : [0, 1] \rightarrow [0, +\infty)$  be a continuous function such that  $W(x) = 0$  if and only if  $x = 1$ , and let  $\psi : [0, 1] \rightarrow [0, 1]$  be an increasing lower semicontinuous function with  $\psi(0) = 0$ ,  $\psi(1) = 1$ , and  $\psi(t) > 0$  if  $t \neq 0$ . Let  $G_\varepsilon : L^1(\Omega) \times L^1(\Omega) \rightarrow [0, +\infty)$  be defined by*

$$G_\varepsilon(u, v) = \begin{cases} \int_{\Omega} \left( \psi(v) |\nabla u| + \frac{1}{\varepsilon} W(v) + \varepsilon |\nabla v|^2 \right) dx & \text{if } u, v \in H^1(\Omega) \\ & \text{and } 0 \leq v \leq 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

*Then there exists the  $\Gamma$ - $\lim_{\varepsilon \rightarrow 0+} G_\varepsilon(u, v) = G(u, v)$  with respect to the  $L^1(\Omega) \times L^1(\Omega)$ -convergence, where*

$$G(u, v) = \begin{cases} \int_{\Omega} |\nabla u| dx + \int_{S_u} g(|u^+ - u^-|) d\mathcal{H}^{n-1} + |D^c u|(\Omega) & \text{if } u \in GBV(\Omega) \\ & \text{and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise,} \end{cases}$$

and

$$g(z) := \min \{ \psi(x)z + 2c_W(x) : 0 \leq x \leq 1 \}, \quad (17)$$

with  $c_W(x) := 2 \int_x^1 \sqrt{W(s)} ds$ .

The proof of the theorem above will be a consequence of the propositions in the rest of the section. Before entering into the details of the proof, we define also a ‘localized version’ of our functionals as follows:

$$G_\varepsilon(u, v, A) = \begin{cases} \int_A \left( \psi(v) |\nabla u| + \frac{1}{\varepsilon} W(v) + \varepsilon |\nabla v|^2 \right) dx & \text{if } u, v \in H^1(\Omega) \\ & \text{and } 0 \leq v \leq 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

and

$$G(u, v, A) = \begin{cases} \int_A |\nabla u| dx + \int_{S_u \cap A} g(|u^+ - u^-|) d\mathcal{H}^{n-1} + |D^c u|(A) & \text{if } u \in GBV(\Omega) \text{ and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise,} \end{cases}$$

for any  $A \in \Omega$  bounded open set.

**Remark 4.2** By the assumptions on  $\psi$  and  $W$ , it can be easily proved that  $g$  satisfies the following properties

(i)  $g$  is increasing,  $g(0) = 0$  and

$$\lim_{z \rightarrow +\infty} g(z) = 2c_W(0) = 4 \int_0^1 \sqrt{W(s)} ds;$$

(ii)  $g$  is subadditive, i.e.

$$g(z_1 + z_2) \leq g(z_1) + g(z_2) \quad \forall z_1, z_2 \in \mathbf{R}^+;$$

(iii)  $g$  is Lipschitz-continuous with Lipschitz constant 1;

(iv)  $g(z) \leq z$  for all  $z \in \mathbf{R}^+$  and

$$\lim_{z \rightarrow 0^+} \frac{g(z)}{z} = 1;$$

(v) for any  $T > 0$  there exists a constant  $c_T > 0$  such that  $z \leq c_T g(z)$  for all  $z \in [0, T]$ .

**Proposition 4.3** *Let  $n = 1$ . Then  $G(u, v) \leq \Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} G_\varepsilon(u, v)$  for all  $u, v \in L^1(\Omega)$ .*

PROOF. It suffices to consider the case in which the right-hand side is finite. Let  $\varepsilon_j \rightarrow 0^+$ ,  $u_j \rightarrow u$  and  $v_j \rightarrow v$  in  $L^1(\Omega)$  be such that  $\lim_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) = \Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} G_\varepsilon(u, v)$ . Up to passing to subsequences we may suppose

$$u_j \rightarrow u, \text{ and } v_j \rightarrow v \text{ a.e.} \quad (18)$$

We have

$$\int_{\Omega} W(v_j) dx < c\varepsilon_j;$$

hence, by the continuity of  $W$ , for any  $\eta > 0$   $\mathcal{L}^1(\{x \in \Omega : W(v(x)) > \eta\}) = \lim_{j \rightarrow +\infty} \mathcal{L}^1(\{x \in \Omega : W(v_j(x)) > \eta\}) = 0$ . We conclude that  $W(v) = 0$  a.e., i.e.  $v = 1$  a.e.

By simplicity, suppose that  $\Omega = (a, b)$  (otherwise we split  $\Omega$  into its connected components). We now use a discretization argument similar to the one used in the proof of [3]. Let  $N \in \mathbf{N}$  and consider the intervals

$$I_N^k = \left( a + \frac{(k-1)}{N}(b-a), a + \frac{k}{N}(b-a) \right), \quad k \in \{1, \dots, N\}.$$

Up to passing to subsequences we may suppose that

$$\lim_{j \rightarrow +\infty} \inf_{I_N^k} v_j$$

exists for all  $N \in \mathbf{N}$  and  $k \in \{1, \dots, N\}$ . Let  $z \in (0, 1)$  be fixed and consider the set

$$J_N^z = \left\{ k \in \{1, \dots, N\} : \lim_{j \rightarrow +\infty} \inf_{I_N^k} v_j \leq z \right\}.$$

Note that for any  $(\alpha, \beta)$  interval in  $\mathbf{R}$  and for any  $w \in H^1(\alpha, \beta)$  we have, by Young's inequality,

$$\int_{\alpha}^{\beta} \left( \frac{1}{\varepsilon} W(w) + \varepsilon |w'|^2 \right) dx \geq 2 \int_{\alpha}^{\beta} \sqrt{W(w)} |w'| dx \geq 2 \left| \int_{w(\alpha)}^{w(\beta)} \sqrt{W(s)} ds \right|.$$

From this inequality we deduce, arguing as in [3], that

$$\left( 2 \int_z^1 \sqrt{W(s)} ds \right) \# J_N^z \leq \lim_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) < +\infty.$$

Then

$$\# J_N^z \leq C$$

with  $C$  independent of  $N$ . Hence, up to a subsequence, we may suppose

$$J_N^z = \{k_1^N, \dots, k_L^N\}$$

with  $L$  independent of  $N$ , and up to a further subsequence that there exist  $S = \{t_1, \dots, t_L\} \subset [a, b]$  such that

$$\lim_{N \rightarrow +\infty} \frac{k_i^N}{N} = t_i$$

for any  $i \in \{1, \dots, L\}$ . For every  $\eta > 0$  we have

$$I_N^k \subset S_{\eta} := S + [-\eta, \eta]$$

for all  $k \in J_N^z$  and for  $N$  large enough. Then

$$\begin{aligned} \liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) &\geq \liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j, \Omega \setminus S_{\eta}) \\ &\quad + \liminf_{j \rightarrow +\infty} \sum_{i=1}^L G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta)) \\ &\geq \liminf_{j \rightarrow +\infty} \psi(z) \int_{\Omega \setminus S_{\eta}} |u'_j| dt \\ &\quad + \sum_{i=1}^L \liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta)). \end{aligned} \quad (19)$$

With fixed  $i \in \{1, \dots, L\}$ , we focus our attention on the term  $G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta))$ . By definition and by (18), we have that for any  $\delta > 0$  there exist

$x_1, x_2 \in (t_i - \eta, t_i + \eta)$  such that

$$\begin{aligned} \lim_{j \rightarrow +\infty} u_j(x_1) &= u(x_1) < \operatorname{ess-}\inf_{(t_i - \eta, t_i + \eta)} u + \delta, \\ \lim_{j \rightarrow +\infty} u_j(x_2) &= u(x_2) > \operatorname{ess-}\sup_{(t_i - \eta, t_i + \eta)} u - \delta, \\ \lim_{j \rightarrow +\infty} v_j(x_1) &= \lim_{j \rightarrow +\infty} v_j(x_2) = 1. \end{aligned} \quad (20)$$

Let  $x_j^i \in [x_1, x_2]$  be such that  $v_j(x_j^i) = \inf_{[x_1, x_2]} v_j$ . Then we obtain the following estimate:

$$\begin{aligned} G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta)) &\geq G_{\varepsilon_j}(u_j, v_j, (x_1, x_2)) \\ &\geq \psi(v_j(x_j^i)) \left| \int_{x_1}^{x_2} u'_j dx \right| + 2 \int_{x_1}^{x_2} \sqrt{W(v_j)} |v'_j| dx \\ &\geq \psi(v_j(x_j^i)) |u_j(x_2) - u_j(x_1)| \\ &\quad + 2 \int_{v_j(x_j^i)}^{v_j(x_1)} \sqrt{W(s)} ds + 2 \int_{v_j(x_j^i)}^{v_j(x_2)} \sqrt{W(s)} ds \\ &\geq \inf_{t \in [0, 1]} \left\{ \psi(t) |u_j(x_2) - u_j(x_1)| \right. \\ &\quad \left. + 2 \left( \int_t^{v_j(x_1)} \sqrt{W(s)} ds + \int_t^{v_j(x_2)} \sqrt{W(s)} ds \right) \right\}. \end{aligned} \quad (21)$$

Letting  $j \rightarrow +\infty$  and taking into account (20), we get

$$\begin{aligned} &\liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta)) \\ &\geq \inf_{t \in [0, 1]} \left\{ \psi(t) \left| \operatorname{ess-}\sup_{(t_i - \eta, t_i + \eta)} u - \operatorname{ess-}\inf_{(t_i - \eta, t_i + \eta)} u - 2\delta \right| + 4 \int_t^1 \sqrt{W(s)} ds \right\}. \end{aligned}$$

Thus, by the arbitrariness of  $\delta > 0$ ,

$$\liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j, (t_i - \eta, t_i + \eta)) \geq g \left( \operatorname{ess-}\sup_{(t_i - \eta, t_i + \eta)} u - \operatorname{ess-}\inf_{(t_i - \eta, t_i + \eta)} u \right) \quad (22)$$

Now we turn back to the estimate (19). Since  $\sup_j G_{\varepsilon_j}(u_j, v_j) < +\infty$ , by (19) we get the equiboundedness of  $\int_{\Omega \setminus S_\eta} |u'_j| dt$ . Hence  $u \in BV(\Omega \setminus S_\eta)$  and, by (19) and (22),

$$\liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) \geq \psi(z) |Du|(\Omega \setminus S_\eta) + \sum_{i=1}^L g \left( \operatorname{ess-}\sup_{(t_i - \eta, t_i + \eta)} u - \operatorname{ess-}\inf_{(t_i - \eta, t_i + \eta)} u \right). \quad (23)$$

By the arbitrariness of  $\eta$ , we deduce that  $u \in BV(\Omega \setminus S)$ , i.e., since  $S$  is finite,  $u \in BV(\Omega)$ . Then, letting  $\eta \rightarrow 0$  in (23), we get

$$\begin{aligned} \liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) &\geq \psi(z)|Du|(\Omega \setminus S) + \sum_{i=1}^L g(|u^+ - u^-|(t_i)) \\ &\geq \psi(z)|Du|(\Omega \setminus S_u) + \sum_{t \in S_u} \left( g(|u^+ - u^-|(t)) \wedge \psi(z)|u^+ - u^-|(t) \right). \end{aligned} \quad (24)$$

Finally, letting  $z \rightarrow 1$  in (24) we obtain the required inequality, since  $g(t) \leq t$ .  $\square$

We recover, now, the  $n$ -dimensional analogue of the previous inequality, by using Theorem 3.4.

**Proposition 4.4** *Let  $n \in \mathbf{N}$ . Then  $G(u, v) \leq \Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} G_\varepsilon(u, v)$  for all  $u, v \in L^1(\Omega)$ .*

PROOF. In the following we will use the notation  $G' = \Gamma\text{-}\liminf_{\varepsilon \rightarrow 0^+} G_\varepsilon$ .

Let  $\xi \in S^{n-1}$  be fixed and let  $\Pi_\xi$  be the hyperplane through 0 orthogonal to  $\xi$ . For any  $u \in L^1(\Omega)$ ,  $A \in \mathcal{A}(\Omega)$ ,  $y \in \Pi_\xi$  we set

$$A_{\xi y} := \{t \in \mathbf{R} : y + t\xi \in A\}, \quad u_{\xi y}(t) := u(y + t\xi).$$

In particular, if  $u \in H^1(\Omega)$ , we get

$$u'_{\xi y}(t) := \langle \nabla u(y + t\xi), \xi \rangle.$$

For any  $u, v \in H^1(\Omega)$ ,  $0 \leq v \leq 1$ , we have, by Fubini's Theorem,

$$\begin{aligned} &G_\varepsilon(u, v, A) \\ &= \int_{\Pi_\xi} \int_{A_{\xi y}} \left( \psi(v(y + t\xi)) |\nabla u(y + t\xi)| \right. \\ &\quad \left. + \frac{1}{\varepsilon} W(v(y + t\xi)) + \varepsilon |\nabla v(y + t\xi)|^2 \right) dt d\mathcal{H}^{n-1}(y) \\ &\geq \int_{\Pi_\xi} \int_{A_{\xi y}} \left( \psi(v_{\xi y}(t)) |u'_{\xi y}| + \frac{1}{\varepsilon} W(v_{\xi y}(t)) + \varepsilon |v'_{\xi y}(t)|^2 \right) dt d\mathcal{H}^{n-1}(y) \\ &= \int_{\Pi_\xi} \mathcal{G}_\varepsilon(u_{\xi y}, v_{\xi y}, A_{\xi y}) d\mathcal{H}^{n-1}(y), \end{aligned} \quad (25)$$

where  $\mathcal{G}_\varepsilon$  is defined by

$$\mathcal{G}_\varepsilon(u, v, I) = \begin{cases} \int_I \left( \psi(v) |u'| + \frac{1}{\varepsilon} W(v) + \varepsilon |v''|^2 \right) dt & \text{if } u, v \in H^1(I) \\ & \text{and } 0 \leq v \leq 1 \\ +\infty & \text{otherwise,} \end{cases}$$

for any  $u, v \in L^1(I)$  and  $I \subset \mathbf{R}$  open and bounded.

Let  $\varepsilon_j \rightarrow 0$  and let  $u_j \rightarrow u, v_j \rightarrow v$  in  $L^1(\Omega)$  be such that

$$\liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j) < +\infty. \quad (26)$$

Then  $u_j, v_j \in H^1(\Omega)$ ,  $0 \leq v_j \leq 1$  a.e. and, as in the proof of Proposition 4.3,  $v = 1$  a.e. Moreover, by Fubini's Theorem,  $(u_j)_{\xi y} \rightarrow u_{\xi y}, (v_j)_{\xi y} \rightarrow 1$  in  $L^1(\Omega_{\xi y})$  for  $\mathcal{H}^{n-1}$ -a.a.  $y \in \Pi_\xi$ .

Thus by Proposition 4.3 and by Fatou's Lemma we get

$$\begin{aligned} & \liminf_{j \rightarrow +\infty} G_{\varepsilon_j}(u_j, v_j, A) \\ & \geq \int_{\Pi_\xi} \liminf_{j \rightarrow +\infty} \mathcal{G}_{\varepsilon_j}((u_j)_{\xi y}, (v_j)_{\xi y}, A_{\xi y}) d\mathcal{H}^{n-1}(y) \\ & \geq \int_{\Pi_\xi} \left( \int_{A_{\xi y}} |u'_{\xi y}| dt + \int_{S_{u_{\xi y}} \cap A_{\xi y}} g(|u_{\xi y}^+ - u_{\xi y}^-|) d\# + |D^c u_{\xi y}|(A_{\xi y}) \right) d\mathcal{H}^{n-1}(y). \end{aligned} \quad (27)$$

Let  $T > 0$  and set

$$u_T = (-T) \vee (u \wedge T).$$

Since  $g$  is increasing, it is clear that we decrease the last term in (27) if we substitute  $u$  by  $u_T$ . Moreover, since  $u_T \in L^\infty(\Omega)$ , with  $\|u_T\|_\infty \leq T$ , by Remark 4.2(v), we have

$$|u_T^+ - u_T^-| \leq c_T g(|u_T^+ - u_T^-|)$$

for a suitable constant  $c_T$  depending only on  $T$ . Then, by (26) and (27), we have

$$\int_{\Pi_\xi} |Du_T|(A_{\xi y}) d\mathcal{H}^{n-1}(y) < +\infty.$$

Thus, applying Theorem 3.4, we get that  $u_T \in BV(\Omega)$  and, by the arbitrariness of  $(u_j)$  and  $(v_j)$ ,

$$G'(u, 1, A) \geq \int_A |\langle \nabla u_T, \xi \rangle| dx + \int_{S_u \cap A} g(|u_T^+ - u_T^-|) |\langle \nu_u, \xi \rangle| d\mathcal{H}^{n-1} + |\langle D^c u_T, \xi \rangle|(A) \quad (28)$$

for all  $A \in \mathcal{A}(\Omega)$  and  $\xi \in S^{n-1}$ .

Consider the superadditive increasing function defined on  $\mathcal{A}(\Omega)$  by

$$\gamma(A) := G'(u, 1, A)$$

and the Radon measure

$$\lambda := \mathcal{L}^n \llcorner \Omega + g(|u_T^+ - u_T^-|) \mathcal{H}^{n-1} \llcorner S_{u_T} + |D^c u_T|.$$

Fixed a sequence  $(\xi_i)_{i \in \mathbf{N}}$ , dense in  $S^{n-1}$ , we have, by (28),

$$\gamma(A) \geq \int_A \psi_i d\lambda$$

for all  $i \in \mathbf{N}$ , where

$$\psi_i(x) = \begin{cases} |\langle \nabla u_T(x), \xi_i \rangle| & \mathcal{L}^n \text{ a.e. on } \Omega \\ |\langle \nu_u(x), \xi_i \rangle| & |D^c u_T| \text{ a.e. on } \Omega \setminus S_{u_T} \\ |\langle \nu_u(x), \xi_i \rangle| & \mathcal{H}^{n-1} \text{ a.e. on } S_{u_T}. \end{cases}$$

Hence, applying Lemma 3.3, we get

$$G'(u, 1, A) \geq \int_A |\nabla u_T| dx + \int_{S_{u_T} \cap A} g(|u_T^+ - u_T^-|) d\mathcal{H}^{n-1} + |D^c u_T|(A) \quad (29)$$

for all  $A \in \mathcal{A}(\Omega)$ . In particular

$$G'(u, 1, \Omega) \geq \int_{\Omega} |\nabla u_T| dx + \int_{S_{u_T}} g(|u_T^+ - u_T^-|) d\mathcal{H}^{n-1} + |D^c u_T|(\Omega). \quad (30)$$

Finally, by the arbitrariness of  $T > 0$ ,  $u \in GBV(\Omega)$  and the thesis follows letting  $T \rightarrow +\infty$  in (30).  $\square$

**Proposition 4.5** *We have  $\Gamma\text{-lim sup}_{\varepsilon \rightarrow 0^+} G_\varepsilon(u, v) \leq G(u, v)$  for all  $u, v \in L^1(\Omega)$ .*

PROOF. It suffices to prove the inequality for  $v = 1$  a.e. Since we will use density and relaxation arguments, we divide the proof into five steps, passing from a particular choice of  $u$  to the general one. In the following we will use the notation  $G'' = \Gamma\text{-lim sup}_{\varepsilon \rightarrow 0^+} G_\varepsilon$ .

Step 1. Suppose that  $u \in \mathcal{W}(\Omega)$  and

$$\overline{S}_u = \Omega \cap K$$

with  $K$  a  $(n-1)$ -dimensional simplex. Up to a translation and rotation argument, we can suppose that  $K$  is contained in the hyperplane  $\Pi := \{x_n = 0\}$ . Set

$$h(y) := u^+(y) - u^-(y), \quad y \in \overline{S}_u.$$

By our hypotheses on  $u$ ,  $h$  is regular on  $\overline{S}_u$ ; hence, fixed  $\delta > 0$ , we can find a triangulation  $\{T_i\}_{i=1}^N$  of  $\overline{S}_u$  such that

$$|h(y_1) - h(y_2)| < \delta \quad \text{if } y_1, y_2 \in T_i.$$

Let  $h_\delta : \overline{S}_u \rightarrow \mathbf{R}$  be defined as

$$h_\delta(y) := z_i \quad y \in T_i,$$

where  $z_i := \min \{h(y) : y \in \overline{T}_i\}$ . Since  $\|h - h_\delta\|_\infty < \delta$ , by Remark 4.2 (iii), we have that

$$\int_{S_u} g(h_\delta(y)) d\mathcal{H}^{n-1} \leq \int_{S_u} g(h(y)) d\mathcal{H}^{n-1} + \delta \mathcal{H}^{n-1}(\overline{S}_u).$$

Let  $x_{z_i}$  realize the minimum in (17) for  $z = z_i$ . Fixed  $\eta > 0$ , there exists  $T(\eta) > 0$  such that

$$\min \left\{ \int_0^T (|v'|^2 + W(v)) dt : v \in H^1(0, T), v(0) = x_{z_i}, v(T) = 1 \right\} \leq c_W(x_{z_i}) + \eta \quad (31)$$

for all  $T \geq T(\eta)$  and for any  $i = 1, \dots, N$ . Let  $v(z_i, \cdot)$  realize the minimum in (31).

For  $r > 0, \varepsilon > 0$  and  $i \in \{1, \dots, N\}$ , set

$$B_r := \left\{ (y, t) \in \Omega : y \in \overline{S}_u, |t| < r \right\} \quad \text{and} \quad T_i^\varepsilon := \left\{ y \in T_i : d(y, \partial T_i) > \varepsilon \right\},$$

and let  $\phi_\varepsilon^i : \mathbf{R}^{(n-1)} \rightarrow \mathbf{R}$  be a cut-off function between  $T_i^\varepsilon$  and  $T_i$  such that  $\|\nabla \phi_\varepsilon^i\|_\infty < C\varepsilon^{-1}$ . Fix a sequence  $(\xi_\varepsilon)$  such that  $\lim_{\varepsilon \rightarrow 0+} \frac{\xi_\varepsilon}{\varepsilon} = 0$ , set  $T_\varepsilon := T(\eta)\varepsilon + \xi_\varepsilon$ , and define

$$v_\varepsilon(y, t) := \begin{cases} 1 & \text{if } (y, t) \in \Omega \setminus B_{T_\varepsilon} \\ \phi_\varepsilon^i(y)v_\varepsilon^i(t) + (1 - \phi_\varepsilon^i(y)) & \text{if } y \in T_i, |t| < T_\varepsilon, \end{cases}$$

where

$$v_\varepsilon^i(t) := \begin{cases} x_{z_i} & \text{if } |t| < \xi_\varepsilon \\ v\left(z_i, \frac{|t| - \xi_\varepsilon}{\varepsilon}\right) & \text{if } \xi_\varepsilon < |t| < T_\varepsilon. \end{cases}$$

We have that  $(v_\varepsilon) \in H^1(\Omega)$  and  $v_\varepsilon \rightarrow 1$  in  $L^1(\Omega)$  as  $\varepsilon \rightarrow 0+$ . Hence, we get

$$\begin{aligned} & \int_\Omega \left( \varepsilon |\nabla v_\varepsilon|^2 + \frac{1}{\varepsilon} W(v_\varepsilon) \right) dx \\ &= \sum_{i=1}^N \int_{T_i^\varepsilon} 2 \int_{\xi_\varepsilon}^{T_\varepsilon} \frac{1}{\varepsilon} \left( \left| v' \left( z_i, \frac{|t| - \xi_\varepsilon}{\varepsilon} \right) \right|^2 + W \left( v \left( z_i, \frac{|t| - \xi_\varepsilon}{\varepsilon} \right) \right) \right) dt d\mathcal{H}^{n-1}(y) \\ & \quad + \sum_{i=1}^N \int_{T_i} \int_{-\xi_\varepsilon}^{\xi_\varepsilon} \left( \varepsilon |\nabla \phi_\varepsilon^i(y)|^2 |x_{z_i} - 1|^2 + \frac{1}{\varepsilon} W(v_\varepsilon(y, t)) \right) dt d\mathcal{H}^{n-1}(y) \\ & \quad + \sum_{i=1}^N \int_{T_i \setminus T_i^\varepsilon} \int_{\xi_\varepsilon}^{T_\varepsilon} \left( \varepsilon |\nabla \phi_\varepsilon^i(y)|^2 \left| v \left( z_i, \frac{|t| - \xi_\varepsilon}{\varepsilon} \right) - 1 \right|^2 \right. \end{aligned} \quad (32)$$



$$\begin{aligned}
& + \frac{1}{\varepsilon} |\phi_\varepsilon^i(y)|^2 \left| v' \left( z_i, \frac{|t| - \xi_\varepsilon}{\varepsilon} \right) \right|^2 \Big) dt d\mathcal{H}^{n-1}(y) \\
& + \sum_{i=1}^N \int_{T_i \setminus T_i^\varepsilon} \int_{\xi_\varepsilon}^{T_\varepsilon} \frac{1}{\varepsilon} W(v_\varepsilon(y, t)) dt d\mathcal{H}^{n-1}(y) \\
\leq & \sum_{i=1}^N \int_{T_i^\varepsilon} 2 \int_0^T \left( |v'(z_i, t)|^2 + W(v(z_i, t)) \right) dt d\mathcal{H}^{n-1}(y) \\
& + c \frac{\xi_\varepsilon}{\varepsilon} \mathcal{H}^{n-1}(S_u) + c(\eta) \sum_{i=1}^N \mathcal{H}^{n-1}(T_i \setminus T_i^\varepsilon) \\
\leq & \sum_{i=1}^N 2 \int_{T_i} c_W(x_{z_i}) d\mathcal{H}^{n-1}(y) + 2\eta \mathcal{H}^{n-1}(S_u) + O(\varepsilon).
\end{aligned}$$

We now construct a recovery sequence  $u_\varepsilon$ . Let

$$\tilde{u}_\varepsilon(z_1, z_2, t) = \begin{cases} z_1 & -T_\varepsilon < t < -\xi_\varepsilon \\ \frac{z_2 - z_1}{2\xi_\varepsilon}(t + \xi_\varepsilon) + z_1 & |t| < \xi_\varepsilon \\ z_2 & \xi_\varepsilon < t < T_\varepsilon \end{cases}$$

and set

$$u_\varepsilon(y, t) = \begin{cases} u(y, t) & |t| > T_\varepsilon \\ \tilde{u}_\varepsilon(u(y, -T_\varepsilon), u(y, T_\varepsilon), t) & |t| < T_\varepsilon \end{cases}.$$

It can be easily verified that  $u_\varepsilon \in H^1(\Omega)$  and  $u_\varepsilon \rightarrow u$  in  $L^1(\Omega)$  as  $\varepsilon \rightarrow 0^+$ . Moreover, we have

$$\begin{aligned}
\int_\Omega \psi(v_\varepsilon) |\nabla u_\varepsilon| dx & \leq \sum_{i=1}^N \int_{T_i^\varepsilon} \int_{-\xi_\varepsilon}^{\xi_\varepsilon} \frac{1}{2\xi_\varepsilon} \psi(x_{z_i}) |u(y, T_\varepsilon) - u(y, -T_\varepsilon)| dt d\mathcal{H}^{n-1}(y) \\
& + \int_{\Omega \setminus B_{T_\varepsilon}} |\nabla u| dx + c \mathcal{H}^{n-1}(T_i \setminus T_i^\varepsilon) + O(\varepsilon) \\
& = \int_\Omega |\nabla u| dx + \sum_{i=1}^N \int_{T_i} \psi(x_{z_i}) |u^+ - u^-|(y) d\mathcal{H}^{n-1}(y) + O(\varepsilon).
\end{aligned} \tag{33}$$

Letting, now,  $\varepsilon$  tend to  $0^+$ , we obtain, by (32) and (33),

$$\begin{aligned}
G''(u, 1) & \leq \limsup_{\varepsilon \rightarrow 0^+} G_\varepsilon(u_\varepsilon, v_\varepsilon) \\
& \leq \int_\Omega |\nabla u| dx + \sum_{i=1}^N \int_{T_i} (|u^+ - u^-|(y) \psi(x_{z_i}) + 2c_W(x_{z_i})) d\mathcal{H}^{n-1}(y) + c\eta
\end{aligned}$$

$$\begin{aligned}
&\leq \int_{\Omega} |\nabla u| dx + \sum_{i=1}^N \int_{T_i} (z_i \psi(x_{z_i}) + 2c_W(x_{z_i})) d\mathcal{H}^{n-1}(y) + c(\eta + \delta) \\
&= \int_{\Omega} |\nabla u| dx + \int_{S_u} g(h_{\delta}(y)) d\mathcal{H}^{n-1}(y) + c(\eta + \delta) \\
&\leq \int_{\Omega} |\nabla u| dx + \int_{S_u} g(|u^+ - u^-|(y)) d\mathcal{H}^{n-1}(y) + c(\eta + \delta).
\end{aligned}$$

Letting  $\eta$  and  $\delta$  tend to  $0^+$ , we obtain the required inequality.

In order to use the same construction as above in the case  $\overline{S}_u = \Omega \cap \left(\bigcup_{i=1}^M K_i\right)$ , with  $M > 1$ , we now show that we can replace  $(u_{\varepsilon})$  by a new sequence  $(\hat{u}_{\varepsilon})$  such that  $\hat{u}_{\varepsilon} \neq u$  only in a small neighbourhood of  $K$ . To this end we again use a cut-off argument. Set

$$K_{\varepsilon} := \{y \in \Pi : d(y, K) < \varepsilon\}$$

and let  $\phi_{\varepsilon} : \mathbf{R}^{n-1} \rightarrow \mathbf{R}$  be a cut-off function between  $K$  and  $K_{\varepsilon}$  with  $|\nabla \phi_{\varepsilon}|_{\infty} \leq c\varepsilon^{-1}$ . Define

$$\hat{u}_{\varepsilon}(y, t) := \phi_{\varepsilon}(y)u_{\varepsilon}(y, t) + (1 - \phi_{\varepsilon}(y))u(y, t) \quad (y, t) \in \Omega.$$

We have

$$\begin{aligned}
\hat{u}_{\varepsilon}(y, t) &= u_{\varepsilon}(y, t) && \text{if } (y, t) \in B_{T_{\varepsilon}}, \\
\hat{u}_{\varepsilon}(y, t) &= u(y, t) && \text{if } (y, t) \in \Omega \setminus K_{\varepsilon} \times (-T_{\varepsilon}, T_{\varepsilon}).
\end{aligned} \tag{34}$$

Then

$$\begin{aligned}
\int_{\Omega \setminus B_{T_{\varepsilon}}} |\nabla \hat{u}_{\varepsilon}| dx &\leq \int_{\Omega \setminus K_{\varepsilon} \times (-T_{\varepsilon}, T_{\varepsilon})} |\nabla u| dx \\
&\quad + \int_{\Omega \cap (K_{\varepsilon} \setminus K)} \int_{-T_{\varepsilon}}^{T_{\varepsilon}} \left( |\nabla \phi_{\varepsilon}(y)| |u_{\varepsilon}(y, t) - u(y, t)| \right) dt d\mathcal{H}^{n-1}(y) \\
&\quad + \int_{\Omega \cap (K_{\varepsilon} \setminus K)} \int_{-T_{\varepsilon}}^{T_{\varepsilon}} \left( \phi_{\varepsilon}(y) |\nabla u_{\varepsilon}(y, t)| \right. \\
&\quad \quad \left. + (1 - \phi_{\varepsilon}(y)) |\nabla u(y, t)| \right) dt d\mathcal{H}^{n-1}(y) \\
&\leq \int_{\Omega} |\nabla u| dx + c \frac{T_{\varepsilon}}{\varepsilon} \mathcal{H}^{n-1}(K_{\varepsilon} \setminus K) + O(\varepsilon).
\end{aligned}$$

Thus

$$\limsup_{\varepsilon \rightarrow 0^+} \int_{\Omega \setminus B_{T_{\varepsilon}}} |\nabla \hat{u}_{\varepsilon}| dx = \int_{\Omega} |\nabla u| dx,$$

and, by (34), we still have

$$\limsup_{\varepsilon \rightarrow 0^+} G_{\varepsilon}(\hat{u}_{\varepsilon}, v_{\varepsilon}) \leq G(u, 1) + c(\eta + \delta).$$

Step 2. If  $u \in \mathcal{W}(\Omega)$  with  $\overline{S}_u = \Omega \cap \left( \bigcup_{i=1}^M K_i \right)$ , we can generalize in a very natural way the construction of the recovery sequences  $\hat{u}_\varepsilon$  and  $v_\varepsilon$  in Step 1, since this construction modifies  $u$  and  $v$  only in a small neighbourhood of each sets  $K_i$ .

Step 3. Let  $u \in SBV^2(\Omega) \cap L^\infty(\Omega)$ . Then, applying Theorem 3.1 with  $\phi(a, b, \nu) = g(|a - b|)$ , there exists a sequence  $(w_j) \in \mathcal{W}(\Omega)$  such that

$$w_j \rightarrow u \text{ in } L^1(\Omega), \text{ and } \limsup_{j \rightarrow +\infty} G(w_j, 1) \leq G(u, 1).$$

Then, by the previous steps and by the lower semicontinuity of  $G''$

$$G''(u, 1) \leq \liminf_{j \rightarrow +\infty} G''(w_j, 1) \leq \liminf_{j \rightarrow +\infty} G(w_j, 1) \leq G(u, 1).$$

Step 4. Since  $g$  satisfies the hypotheses of Theorem 3.2, the relaxation with respect to  $L^1(\Omega)$ -topology of the functional

$$F(u) := \begin{cases} G(u, 1) & \text{if } u \in SBV^2(\Omega) \cap L^\infty(\Omega) \\ +\infty & \text{otherwise in } BV(\Omega) \end{cases}$$

is given by

$$\overline{F}(u) = G(u, 1)$$

for all  $u \in BV(\Omega)$ . Then by the previous steps and by the lower semicontinuity of  $G''$  we get

$$G''(u, 1) \leq \overline{F}(u) = G(u, 1)$$

for any  $u \in BV(\Omega)$ .

Step 5. We recover the general case by a truncation argument. Let  $u \in GBV(\Omega)$  and let  $u_j = (-j) \vee (u \wedge j)$ . Then

$$\lim_{j \rightarrow +\infty} G(u_j, 1) = G(u, 1).$$

Since  $u_j \rightarrow u$  in  $L^1(\Omega)$  we get the thesis by the lower semicontinuity of  $G''$ .  $\square$

**Example 4.6** We illustrate with a few simple examples the behaviour of the function  $g$ , given by (17), with different choices of  $\psi$ .

. The terms "bulk energy" and "surface energy" refer to the first and second terms in Equation (17) and what is meant by interaction between the two at each value of  $z$  is relative contributions of these terms to  $g(z)$  at that  $z$ .

Let  $W(v) = (1 - v)^2/4$ , so that  $c_W(x) = (1 - x)^2/2$ . We then have

- (a) if  $\psi(v) = v^2$  then  $g(z) = |z|/(1 + |z|)$ ;
- (b) if  $\psi(v) = v$  then  $g(z) = \begin{cases} |z| - (z^2/4) & \text{if } |z| \leq 2 \\ 1 & \text{if } |z| > 2; \end{cases}$
- (c) if  $\psi(v) = \begin{cases} 0 & \text{if } v = 0 \\ 1 & \text{otherwise,} \end{cases}$  then  $g(z) = \min\{|z|, 1\}$ .

We see that the ‘bulk term’ and of the ‘surface term’ (i.e. the first and the second term in (17)) play different roles in these examples. Note that in (a) we always have interaction between these two terms (i.e. both terms contribute to the value  $g(z)$ ) contrary to what happens in the Ambrosio Tortorelli case. The interaction also occurs in (b) for  $|z| < 2$ . Note moreover that in the third case the minimal  $x$  in the definition of  $g(z)$  does not vary with continuity at  $z = 1$ .

## 5 Approximation of general functionals

In this section we show how Theorem 4.1 can be used to obtain an approximation of general (isotropic) energies defined on  $GSBV$  by a double limit. The set  $\Omega$  will be a bounded open subset of  $\mathbf{R}^n$  with Lipschitz boundary.

**Proposition 5.1** *Let  $W$  and  $\psi$  be defined as in Theorem 4.1, let  $f : [0, +\infty) \rightarrow [0, +\infty)$  be a convex and increasing function satisfying*

$$\lim_{t \rightarrow +\infty} \frac{f(t)}{t} = 1, \quad (35)$$

*and let  $G_\varepsilon : L^1(\Omega) \times L^1(\Omega) \rightarrow [0, +\infty)$  be defined by*

$$G_\varepsilon(u, v) = \begin{cases} \int_{\Omega} \left( \psi(v) f(|\nabla u|) + \frac{1}{\varepsilon} W(v) + \varepsilon |\nabla v|^2 \right) dx & \text{if } u, v \in H^1(\Omega) \\ & \text{and } 0 \leq v \leq 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

*Then there exists the  $\Gamma$ - $\lim_{\varepsilon \rightarrow 0+} G_\varepsilon(u, v) = G(u, v)$  with respect to the  $L^1(\Omega) \times L^1(\Omega)$ -convergence, where*

$$G(u, v) = \begin{cases} \int_{\Omega} f(|\nabla u|) dx + \int_{S_u} g(|u^+ - u^-|) d\mathcal{H}^{n-1} + |D^c u|(\Omega) & \text{if } u \in GBV(\Omega) \\ & \text{and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise,} \end{cases}$$

*and  $g$  is defined in (17).*

**PROOF.** The estimate for the  $\Gamma$ -lim inf can be performed as in Proposition 4.3, noting that in (21) we obtain, by Jensen’s inequality,

$$G_{\varepsilon_j}(u_j, v_j, I_N^{k-i}) \geq \psi(v_j(x_j^i)) |x_2 - x_1| f\left(\frac{u(x_2) - u(x_1)}{x_2 - x_1}\right) + 2 \int_{x_1}^{x_2} \sqrt{W(v_j)} |v_j'| dx,$$

from which the lower bound can be easily obtained taking into account (35). The rest of the proof can be obtained following Propositions 4.4 and 4.5.  $\square$

**Remark 5.2** Let  $K > 0$  and  $N \geq 2$ , let

$$0 = a_0 < a_1 < \dots < a_N = 1, \quad 0 = b_N < b_{N-1} < \dots < b_0 = K,$$

and let  $f$  and  $W$  be as in the previous proposition. Then there exists  $\psi$  satisfying the hypotheses in Theorem 4.1 such that, if  $G_\varepsilon : L^1(\Omega) \times L^1(\Omega) \rightarrow [0, +\infty)$  is defined by

$$G_\varepsilon(u, v) = \begin{cases} \int_{\Omega} \left( \psi(v) f(|\nabla u|) + \frac{K}{\varepsilon} W(v) + \varepsilon K |\nabla v|^2 \right) dx & \text{if } u, v \in H^1(\Omega) \\ & \text{and } 0 \leq v \leq 1 \text{ a.e.} \\ +\infty & \text{otherwise,} \end{cases}$$

then the thesis of the previous proposition holds with  $g : [0, +\infty) \rightarrow [0, +\infty)$  given by

$$g(z) = \min\{a_i z + b_i\}.$$

In fact, in this case the formula for  $g$  can be easily inverted, obtaining  $\psi$  as the piecewise constant function given by  $\psi(0) = 0$  and

$$\psi(\xi) = a_i \quad \text{if } c_W^{-1}(b_{i-1}/2) < \xi \leq c_W^{-1}l(b_i/2),$$

where  $c_W$  is defined in Theorem 4.1.

**Proposition 5.3** Let  $W$  be as in Theorem 4.1. Let  $\varphi, \vartheta : [0, +\infty) \rightarrow [0, +\infty)$  be functions satisfying

- (i)  $\varphi$  is convex and increasing,  $\lim_{t \rightarrow +\infty} \varphi(t)/t = +\infty$ ;
- (ii)  $\vartheta$  is concave,  $\lim_{t \rightarrow 0^+} \vartheta(t)/t = +\infty$ .

Then there exist two increasing sequences of functions  $(\varphi_j)$  and  $(\psi_j)$ , and a sequences of positive real numbers  $(k_j)$ , converging to  $\sup \vartheta$ , such that if we define

$$G_\varepsilon^j(u, v) = \begin{cases} \int_{\Omega} \left( \psi_j(v) \varphi_j(|\nabla u|) + \frac{k_j}{\varepsilon} W(v) + k_j \varepsilon |\nabla v|^2 \right) dx & \text{if } u, v \in H^1(\Omega) \\ & \text{and } 0 \leq v \leq 1 \text{ a.e.} \\ +\infty & \text{otherwise,} \end{cases} \quad (36)$$

then for every  $j \in \mathbb{N}$  there exist the limits

$$\begin{aligned} \Gamma\text{-}\lim_{\varepsilon \rightarrow 0^+} G_\varepsilon^j(u, v) &=: G^j(u, v) \\ \Gamma\text{-}\lim_{j \rightarrow +\infty} G^j(u, v) &= \lim_{j \rightarrow +\infty} G^j(u, v) = G(u, v) \end{aligned}$$

with respect to the  $L^1(\Omega) \times L^1(\Omega)$ -convergence, where

$$G(u, v) = \begin{cases} \int_{\Omega} \varphi(|\nabla u|) dx + \int_{S_u} \vartheta(|u^+ - u^-|) d\mathcal{H}^{n-1} & \text{if } u \in GSBV(\Omega) \\ & \text{and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

PROOF. Let  $\vartheta_j : [0, +\infty) \rightarrow [0, +\infty)$  be functions of the form

$$\vartheta_j(z) = \min\{A_i^j z + B_i^j\},$$

with  $0 = A_0^j < \dots < A_j^j = j$  converging increasingly to  $\vartheta$ , and let  $\varphi_j : [0, +\infty) \rightarrow [0, +\infty)$  be convex increasing functions with

$$\lim_{t \rightarrow +\infty} \frac{\varphi_j(t)}{t} = j,$$

converging increasingly to  $\varphi$ . Let  $k_j = \max \vartheta_j$ .

Set  $g_j = \vartheta_j/j$ ,  $K_j = k_j/j$  and  $f_j = \varphi_j/j$ . By the previous remark, applied with  $g = g_j$ ,  $f = f_j$  and  $K = K_j$ , we can find  $\psi =: \psi_j$  such that if we let  $G_\varepsilon^j : L^1(\Omega) \times L^1(\Omega) \rightarrow [0, +\infty)$  be defined by (36) then there exists the  $\Gamma\text{-}\lim_{\varepsilon \rightarrow 0+} G_\varepsilon^j(u, v) = G^j(u, v)$  with respect to the  $L^1(\Omega) \times L^1(\Omega)$ -convergence, where

$$G^j(u, v) = \begin{cases} \int_{\Omega} \varphi_j(|\nabla u|) dx + \int_{S_u} \vartheta_j(|u^+ - u^-|) d\mathcal{H}^{n-1} + j|D^c u|(\Omega) & \text{if } u \in GBV(\Omega) \text{ and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

Since the functionals  $G^j$  converge increasingly to  $G$ , they also  $\Gamma$ -converge to  $G$  as  $j \rightarrow +\infty$ .  $\square$

**Remark 5.4** If  $\varphi$  is convex and even,  $\vartheta$  is concave and even, and

$$\lim_{t \rightarrow +\infty} \frac{\varphi(t)}{t} = \lim_{t \rightarrow 0+} \frac{\vartheta(t)}{t} = M,$$

then there exist  $(\varphi_j)$ ,  $(\psi_j)$  and  $(k_j)$  such that the functionals  $G^j$  defined above  $\Gamma$ -converge with respect to the  $L^1(\Omega) \times L^1(\Omega)$ -convergence to

$$G(u, v) = \begin{cases} \int_{\Omega} \varphi(|\nabla u|) dx + \int_{S_u} \vartheta(|u^+ - u^-|) d\mathcal{H}^{n-1} + M|D^c u|(\Omega) & \text{if } u \in GBV(\Omega) \text{ and } v = 1 \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

The proof can be obtained directly from Remark 5.2, using the approximation argument of Proposition 5.3.

**Acknowledgements** This paper was written while the second author was visiting the Max-Planck-Institute for Mathematics in the Sciences at Leipzig, on Marie-Curie fellowship ERBFMBICT972023 of the European Union program “Training and Mobility of Researchers”. J. Shah is partially supported under NSF Grant DMS-9531293 and NIH Grant I-R01-NS34189-01.

## References

- [1] L. Ambrosio, A compactness theorem for a new class of functions of bounded variation, *Boll. Un. Mat. Ital.* **3-B** (1989), 857-881.
- [2] L. Ambrosio, N. Fusco and D. Pallara, *Special Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, Oxford, to appear.
- [3] L. Ambrosio and V. M. Tortorelli, Approximation of functionals depending on jumps by elliptic functionals via  $\Gamma$ -convergence, *Comm. Pure Appl. Math.* **43** (1990), 999-1036.
- [4] G. Bouchitté, A. Braides and G. Buttazzo, Relaxation results for some free discontinuity problems, *J. Reine Angew. Math.* **458** (1995), 1-18
- [5] A. Braides, *Approximation of Free-discontinuity Problems*, Lecture Notes in Mathematics, Springer Verlag, Berlin, to appear.
- [6] A. Braides and V. Chiadò Piat, Integral representation results for functionals defined on  $SBV(\Omega; \mathbf{R}^m)$ , *J. Math. Pures Appl.* **75** (1996), 595-626.
- [7] G. Cortesani and R. Toader, A density result in  $SBV$  with respect to non-isotropic energies, *Nonlinear Analysis*, to appear.
- [8] G. Dal Maso, *An Introduction to  $\Gamma$ -convergence*, Birkhäuser, Boston, 1993.
- [9] E. De Giorgi, Free Discontinuity Problems in Calculus of Variations, in: *Frontiers in pure and applied Mathematics, a collection of papers dedicated to J.L.Lions on the occasion of his 60<sup>th</sup> birthday* (R. Dautray ed.), North Holland, 1991.
- [10] E. De Giorgi and L. Ambrosio, Un nuovo funzionale del calcolo delle variazioni, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* **82** (1988), 199-210.
- [11] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, Boca Raton 1992.
- [12] H. Federer, *Geometric Measure Theory*, Springer Verlag, New York, 1969.
- [13] L. Modica and S. Mortola, Un esempio di  $\Gamma$ -convergenza, *Boll. Un. Mat. It.* **14-B** (1977), 285-299.
- [14] D. Mumford and J. Shah, Optimal approximation by piecewise smooth functions and associated variational problems, *Comm. Pure Appl. Math.* **17** (1989), 577-685.