

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Quantum Entanglement: Separability, Measure,
Fidelity of Teleportation and Distillation

by

Ming Li, Shao-Ming Fei, and Xianqing Li-Jost

Preprint no.: 10

2010



Quantum Entanglement: Separability, Measure, Fidelity of Teleportation and Distillation

Ming Li¹, Shao-Ming Fei^{2,3} and Xianqing Li-Jost³

¹ Department of Mathematics, China University of Petroleum, 257061 Dongying, China

² Department of Mathematics, Capital Normal University, 100037 Beijing, China

³ Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

Quantum entanglement plays crucial roles in quantum information processing. Quantum entangled states have become the key ingredient in the rapidly expanding field of quantum information science. Although the nonclassical nature of entanglement has been recognized for many years, considerable efforts have been taken to understand and characterize its properties recently. In this review, we introduce some recent results in the theory of quantum entanglement. In particular separability criteria based on the Bloch representation, covariance matrix, normal form and entanglement witness; lower bounds, subadditivity property of concurrence and tangle; fully entangled fraction related to the optimal fidelity of quantum teleportation and entanglement distillation will be discussed in detail.

Contents

1	Introduction	2
2	Separability criteria and normal form	7
2.1	Separability criteria for mixed states	8
2.1.1	Partial positive transpose criterion	8
2.1.2	Reduced density matrix criterion	9
2.1.3	Realignment criterion	9
2.1.4	Criteria based on Bloch representations	10
2.1.5	Covariance matrix criterion	14
2.2	Normal form of quantum states	20

2.3	Entanglement witness based on correlation matrix criterion	25
3	Concurrence and Tangle	27
3.1	Lower and upper bounds of concurrence	29
3.1.1	Lower bound of concurrence from covariance matrix criterion .	29
3.1.2	Lower bound of concurrence from “two-qubit” decomposition .	30
3.1.3	Estimation of multipartite entanglement	36
3.1.4	Bounds of concurrence and tangle	40
3.2	Concurrence and tangle of two entangled states are strictly larger than that of one	41
3.3	Subadditivity of concurrence and tangle	43
4	Fidelity of teleportation and distillation of entanglement	44
4.1	Fidelity of quantum teleportation	45
4.2	Fully entangled fraction and concurrence	51
4.3	Improvement of entanglement distillation protocol	54
5	Summary and Conclusion	57
	References	57

1 Introduction

Entanglement is the characteristic trait of quantum mechanics, and it reflects the property that a quantum system can simultaneously appear in two or more different states [1]. This feature implies the existence of global states of composite system which cannot be written as a product of the states of individual subsystems. This phenomenon [2], now known as “quantum entanglement”, plays crucial roles in quantum information processing [3]. Quantum entangled states have become the key ingredient in the rapidly expanding field of quantum information science, with remarkable prospective applications such as quantum computation [3, 4], quantum teleportation [5, 6], dense coding [7], quantum cryptographic schemes [8], entan-

glement swapping [9] and remote states preparation (RSP) [10, 11, 12, 13]. All such effects are based on entanglement and have been demonstrated in pioneering experiments.

It has become clear that entanglement is not only the subject of philosophical debates, but also a new quantum resource for tasks which can not be performed by means of classical resources. Although considerable efforts have been taken to understand and characterize the properties of quantum entanglement recently, the physical character and mathematical structure of entangled states have not been satisfactorily understood yet [14, 15]. In this review we mainly introduce some recent results related to our researches on several basic questions in this subject:

(1) Separability of quantum states

We first discuss the separability of a quantum states, namely, for a given quantum state, how can we know whether or not it is entangled.

For pure quantum states, there are many ways to verify the separability. For instance for a bipartite pure quantum state the separability is easily determined in terms of its Schmidt numbers. For multipartite pure states, the generalized concurrence given in [16] can be used to judge if the state is separable or not. In addition separable states must satisfy all possible Bell inequalities [17].

For mixed states we still have no general criterion. The well-known PPT (partial positive transposition) criterion was proposed by Peres in 1996 [18]. It says that for any bipartite separable quantum state the density matrix must be positive under partial transposition. By using the method of positive maps Horodeckis [19] showed that the Peres' criterion is also sufficient for 2×2 and 2×3 bipartite systems. And for higher dimensional states, the PPT criterion is only necessary. Horodecki [20] has constructed some classes entangled states with positive partial transposes for 3×3 and 2×4 systems. States of this kind are said to be bound entangled (BE). Another powerful operational criterion is the realignment criterion [21, 22]. It demonstrates a remarkable ability to detect many bound entangled states and even genuinely tripartite entanglement [23]. Considerable efforts have been made in finding stronger variants and multipartite generalizations for this criterion [24, 25]. It was shown that PPT criterion and realignment criterion are equivalent to the permutations of the density matrix's indices [23]. Another important criterion for separability is the reduction criterion [26, 27]. This criterion is equivalent to the PPT criterion for $2 \times N$ composite systems. Although it is generally weaker than

the PPT, the reduction criteria has tight relation to the distillation of quantum states.

There are also some other necessary criteria for separability. Nielsen et al. [28] presented a necessary criterion called majorization: the decreasing ordered vector of the eigenvalues for ρ is majorized by that of ρ^{A_1} or ρ^{A_2} alone for a separable state. i.e. if a state ρ is separable, then $\lambda_\rho^\downarrow \prec \lambda_{\rho^{A_1}}^\downarrow$, $\lambda_\rho^\downarrow \prec \lambda_{\rho^{A_2}}^\downarrow$. Here λ_ρ^\downarrow denotes the decreasing ordered vector of the eigenvalues of ρ . A d -dimensional vector x^\downarrow is majorized by y^\downarrow , $x^\downarrow \prec y^\downarrow$, if $\sum_{j=1}^k x_j^\downarrow \leq \sum_{j=1}^k y_j^\downarrow$ for $k = 1, \dots, d-1$ and the equality holds for $k = d$. Zeros are appended to the vectors $\lambda_{\rho^{A_1, A_2}}^\downarrow$ such that their dimensions equal to the one of λ_ρ^\downarrow .

In Ref. [20], another necessary criterion called range criterion was given. If a bipartite state ρ acting on the space $\mathcal{H}_A \otimes \mathcal{H}_B$ is separable, then there exists a family of product vectors $\psi_i \otimes \phi_i$ such that: (i) they span the range of ρ ; (ii) the vector $\{\psi_i \otimes \phi_i^*\}_{i=1}^k$ span the range of ρ^{TB} , where $*$ denotes complex conjugation in the basis in which partial transposition was performed, ρ^{TB} is the partially transposed matrix of ρ with respect to the subspace B . In particular, any of the vectors $\psi_i \otimes \phi_i^*$ belongs to the range of ρ .

Recently, some elegant results for the separability problem have been derived. In [29, 30, 31], a separability criteria based on the local uncertainty relations (LUR) was obtained. The authors show that for any separable state $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B$,

$$1 - \sum_k \langle G_k^A \otimes G_k^B \rangle - \frac{1}{2} \langle G_k^A \otimes I - I \otimes G_k^B \rangle^2 \geq 0,$$

where G_k^A or G_k^B are arbitrary local orthogonal and normalized operators (LOOs) in $\mathcal{H}_A \otimes \mathcal{H}_B$. This criterion is strictly stronger than the realignment criterion. Thus more bound entangled quantum states can be recognized by the LUR criterion. The criterion is optimized in [32] by choosing the optimal LOOs. In [33] a criterion based on the correlation matrix of a state has been presented. The correlation matrix criterion is shown to be independent of PPT and realignment criterion [34], i.e. there exist quantum states that can be recognized by correlation criterion while the PPT and realignment criterion fail. The covariance matrix of a quantum state is also used to study separability in [35]. It has been shown that the LUR criterion, including the optimized one, can be derived from the covariance matrix criterion [36].

(2) Measure of quantum entanglement

One of the most difficult and fundamental problems in entanglement theory is to quantify entanglement. The initial idea to quantify entanglement was connected with its usefulness in terms of communication [37]. A good entanglement measure has to fulfill some conditions [38]. For bipartite quantum systems, we have several good entanglement measures such as Entanglement of Formation(EOF), Concurrence, Tangle etc. For two-qubit systems it has been proved that EOF is a monotonically increasing function of the concurrence and an elegant formula for the concurrence was derived analytically by Wootters [39]. However with the increasing dimensions of the subsystems the computation of EOF and concurrence become formidably difficult. A few explicit analytic formulae for EOF and concurrence have been found only for some special symmetric states [40, 41, 42, 43, 44].

The first analytic lower bound of concurrence for arbitrary dimensional bipartite quantum states was derived by Mintert et al. in [45]. By using the positive partial transposition (PPT) and realignment separability criterion analytic lower bounds on EOF and concurrence for any dimensional mixed bipartite quantum states have been derived in [46, 47]. These bounds are exact for some special classes of states and can be used to detect many bound entangled states. In [48] another lower bound on EOF for bipartite states has been presented from a new separability criterion [49]. A lower bound of concurrence based on local uncertainty relations (LURs) criterion is derived in [50]. This bound is further optimized in [32]. The lower bound of concurrence for tripartite systems has been studied in [51]. In [52, 53] the authors presented lower bounds of concurrence for bipartite systems by considering the “two-qubit” entanglement of bipartite quantum states with arbitrary dimensions. It has been shown that this lower bound has a tight relationship with the distillability of bipartite quantum states. Tangle is also a good entanglement measure that has a close relation with concurrence, as it is defined by the square of the concurrence for a pure state. It is also meaningful to derive tight lower and upper bounds for tangle [54].

In [55] Mintert *et al.* proposed an experimental method to measure the concurrence directly by using joint measurements on two copies of a pure state. Then S. P. Walborn *et al.* presented an experimental determination of concurrence for two-qubit states [56], where only one-setting measurement is needed, but two copies of the state have to be prepared in every measurement. In [57] another way of experimental determination of concurrence for two-qubit and multi-qubit states has been presented, in which only one-copy of the state is needed in every measurement. To determine the concurrence of the two-qubit state used in [56], also one-setting

measurement is needed, which avoids the preparation of the twin states or the imperfect copy of the unknown state, and the experimental difficulty is dramatically reduced.

(3) Fidelity of quantum teleportation and distillation

Quantum teleportation, or entanglement-assisted teleportation, is a technique used to transfer information on a quantum level, usually from one particle (or series of particles) to another particle (or series of particles) in another location via quantum entanglement. It does not transport energy or matter, nor does it allow communication of information at super luminal (faster than light) speed.

In [5], Bennett et. al. first presented a protocol to teleport an unknown qubit state by using a pair of maximally entangled pure qubit state. The protocol is generalized to transmit high dimensional quantum states [6]. The optimal fidelity of teleportation is shown to be determined by the fully entangled fraction of the entangled resource which is generally a mixed state. Nevertheless similar to the estimation of concurrence, the computation of the fully entangled fraction for a given mixed state is also very difficult.

The distillation protocol has been presented to get maximally entangled pure states from many entangled mixed states. by means of local quantum operations and classical communication (LQCC) between the parties sharing the pairs of particles in this mixed state [58, 59, 60, 61]. Bennett et. al. first derived a protocol to distill one maximally entangled pure bell state from many copies of not maximally entangled quantum mixed states in [58] in 1996. The protocol is then generalized to distill any bipartite quantum state with higher dimension by Horodeckis in 1999 [62]. It is proven that a quantum state can be always distilled if it violates the reduced matrix separability criterion [62].

This review mainly contains three parts. In section 2 we investigate the separability of quantum states. We first introduce several important separability criteria. Then we discuss the criterions by using the Bloch representation of the density matrix of a quantum state. We also study the covariance matrix of a quantum density matrix and derive separability criterion for multipartite systems. We investigate the normal forms for multipartite quantum states at the end of this section and show that the normal form can be used to improve the power of these criteria. In section 3 we mainly consider the entanglement measure concurrence. We investigate the lower and upper bounds of concurrence for both bipartite and multipartite systems.

We also show that the concurrence and tangle of two entangled quantum states will be always larger than that of one, even both the two states are bound entangled (not distillable). In section 4 we study the fully entangled fraction of an arbitrary bipartite quantum state. We derive precise formula of fully entangled fraction for two qubits system. For bipartite system with higher dimension we obtain tight upper bounds which can not only be used to estimate the optimal teleportation fidelity but also helps to improve the distillation protocol. We further investigate the evolution of the fully entangled fraction when one of the bipartite system undergoes a noisy channel. We give a summary and conclusion in the last section.

2 Separability criteria and normal form

A multipartite pure quantum state $\rho_{12\dots N} \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$ is said to be fully separable if it can be written as

$$\rho_{12\dots N} = \rho_1 \otimes \rho_2 \otimes \dots \otimes \rho_N, \quad (2.1)$$

where ρ_1 and ρ_2, \dots, ρ_N are reduced density matrices defined as $\rho_1 = \text{Tr}_{23\dots N}[\rho_{12\dots N}]$, $\rho_2 = \text{Tr}_{13\dots N}[\rho_{12\dots N}]$, ..., $\rho_N = \text{Tr}_{12\dots N-1}[\rho_{12\dots N}]$. This is equivalent to the condition

$$\rho_{12\dots N} = |\psi_1\rangle\langle\psi_1| \otimes |\phi_2\rangle\langle\phi_2| \otimes \dots \otimes |\mu_N\rangle\langle\mu_N|,$$

where $|\psi_1\rangle \in \mathcal{H}_1$, $|\phi_2\rangle \in \mathcal{H}_2$, ..., $|\mu_N\rangle \in \mathcal{H}_N$.

A multipartite quantum mixed state $\rho_{12\dots N} \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$ is said to be fully separable if it can be written as

$$\rho_{12\dots N} = \sum_i q_i \rho_i^1 \otimes \rho_i^2 \otimes \dots \otimes \rho_i^N, \quad (2.2)$$

where $\rho_i^1, \rho_i^2, \dots, \rho_i^N$ are the reduced density matrices with respect to the systems $1, 2, \dots, N$ respectively, $q_i > 0$ and $\sum_i q_i = 1$. This is equivalent to the condition

$$\rho_{12\dots N} = \sum_i p_i |\psi_i^1\rangle\langle\psi_i^1| \otimes |\phi_i^2\rangle\langle\phi_i^2| \otimes \dots \otimes |\mu_i^N\rangle\langle\mu_i^N|,$$

where $|\psi_i^1\rangle, |\phi_i^2\rangle, \dots, |\mu_i^N\rangle$ are normalized pure states of systems $1, 2, \dots, N$ respectively, $p_i > 0$ and $\sum_i p_i = 1$.

For pure states, the definition (2.1) itself is an operational separability criterion. In particular, for bipartite case, there are Schmidt decompositions:

Theorem 2.1 (*Schmidt decomposition*): Suppose $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ is a pure state of a composite system, AB , then there exist orthonormal states $|i_A\rangle$ for system A , and orthonormal states $|i_B\rangle$ for system B such that

$$|\psi\rangle = \sum_i \lambda_i |i_A\rangle |i_B\rangle,$$

where λ_i are non-negative real numbers satisfying $\sum_i \lambda_i^2 = 1$, known as Schmidt coefficients.

$|i_A\rangle$ and $|i_B\rangle$ are called Schmidt bases with respect to \mathcal{H}_A and \mathcal{H}_B . The number of non-zero values λ_i is called Schmidt number, also known as Schmidt rank, which is invariant under unitary transformations on system A or system B . For a bipartite pure state $|\psi\rangle$, $|\psi\rangle$ is separable if and only if the Schmidt number of $|\psi\rangle$ is one.

For multipartite pure states, one has no such Schmidt decomposition. In [63] it has been verified that any pure three-qubit state $|\Psi\rangle$ can be uniquely written as

$$|\Psi\rangle = \lambda_0 |000\rangle + \lambda_1 e^{i\psi} |100\rangle + \lambda_2 |101\rangle + \lambda_3 |110\rangle + \lambda_4 |111\rangle \quad (2.3)$$

with normalization condition $\lambda_i \geq 0$, $0 \leq \psi \leq \pi$, where $\sum_i \mu_i = 1$, $\mu_i \equiv \lambda_i^2$. Eq. (2.3) is called generalized Schmidt decomposition.

For mixed states it is generally very hard to verify if a decomposition like (2.2) exists. For a given generic separable density matrix, it is also not easy to find the decomposition (2.2) in detail.

2.1 Separability criteria for mixed states

In this section we introduce several separability criteria and the relations among themselves. These criteria have also tight relations with lower bounds of entanglement measures and distillation that will be discussed in the next section.

2.1.1 Partial positive transpose criterion

The positive partial transpose (PPT) criterion provided by Peres [18] says that if a bipartite state $\rho_{AB} \in \mathcal{H}_A \otimes \mathcal{H}_B$ is separable, then the new matrix $\rho_{AB}^{T_B}$ with matrix elements defined in some fixed product basis as:

$$\langle m | \langle \mu | \rho_{AB}^{T_B} | n \rangle | \nu \rangle \equiv \langle m | \langle \nu | \rho_{AB} | n \rangle | \mu \rangle$$

is also a density matrix (i.e. has nonnegative spectrum). The operation T_B , called partial transpose, just corresponds to the transposition of the indices with respect to the second subsystem B . It has an interpretation as a partial time reversal [64].

Afterwards the Horodeckis showed that the Peres' criterion is also sufficient for 2×2 and 2×3 bipartite systems [19]. This criterion is now called PPT or Peres-Horodecki (P-H) criterion. For high-dimensional states, the P-H criterion is only necessary. Horodecki has constructed some classes of families of entangled states with positive partial transposes for 3×3 and 2×4 systems [20]. States of this kind are said to be bound entangled (BE).

2.1.2 Reduced density matrix criterion

Cerf et al. [65] and Horodecki [66] independently, introduced a map $\Gamma : \rho \rightarrow \text{Tr}_B[\rho_{AB}] \otimes I - \rho_{AB}$ ($I \otimes \text{Tr}_A[\rho_{AB}] - \rho_{AB}$), which gives rise to a simple necessary condition for separability in arbitrary dimensions, called the reduction criterion: If ρ_{AB} is separable, then

$$\rho_A \otimes I - \rho_{AB} \geq 0, \quad I \otimes \rho_B - \rho_{AB} \geq 0,$$

where $\rho_A = \text{Tr}_B[\rho_{AB}]$, $\rho_B = \text{Tr}_A[\rho_{AB}]$. This criterion is simply equivalent to the P-H criterion for $2 \times n$ composite systems. It is also sufficient for 2×2 and 2×3 systems. In higher dimensions the reduction criterion is weaker than the P-H criterion.

2.1.3 Realignment criterion

There is yet another class of criteria based on linear contractions on product states. They stem from the new criterion discovered in [67, 22] called computable cross norm (CCN) criterion or matrix realignment criterion which is operational and independent on PPT test [18]. If a state ρ_{AB} is separable then the realigned matrix $\mathcal{R}(\rho)$ with elements $\mathcal{R}(\rho)_{ij,kl} = \rho_{ik,jl}$ has trace norm not greater than one,

$$\|\mathcal{R}(\rho)\|_{KF} \leq 1. \tag{2.4}$$

Quite remarkably, the realignment criterion can detect some PPT entangled (bound entangled) states [67, 22] and can be used for construction of some nondecomposable maps. It also provides nice lower bound for concurrence [47].

2.1.4 Criteria based on Bloch representations

Any Hermitian operator on an N -dimensional Hilbert space \mathcal{H} can be expressed according to the generators of the special unitary group $SU(N)$ [68]. The generators of $SU(N)$ can be introduced according to the transition-projection operators $P_{jk} = |j\rangle\langle k|$, where $|i\rangle$, $i = 1, \dots, N$, are the orthonormal eigenstates of a linear Hermitian operator on \mathcal{H} . Set

$$\omega_l = -\sqrt{\frac{2}{l(l+1)}}(P_{11} + P_{22} + \dots + P_{ll} - lP_{l+1,l+1}),$$

$$u_{jk} = P_{jk} + P_{kj}, \quad v_{jk} = i(P_{jk} - P_{kj}),$$

where $1 \leq l \leq N-1$ and $1 \leq j < k \leq N$. We get a set of $N^2 - 1$ operators

$$\Gamma \equiv \{\omega_l, \omega_2, \dots, \omega_{N-1}, u_{12}, u_{13}, \dots, v_{12}, v_{13}, \dots\},$$

which satisfy the relations

$$\text{Tr}[\lambda_i] = 0, \quad \text{Tr}[\lambda_i \lambda_j] = 2\delta_{ij}, \quad \forall \lambda_i \in \Gamma$$

and thus generate the $SU(N)$ [69].

Any Hermitian operator ρ in \mathcal{H} can be represented in terms of these generators of $SU(N)$,

$$\rho = \frac{1}{N}I_N + \frac{1}{2} \sum_{j=1}^{N^2-1} r_j \lambda_j, \quad (2.5)$$

where I_N is a unit matrix and $\mathbf{r} = (r_1, r_2, \dots, r_{N^2-1}) \in \mathbb{R}^{N^2-1}$. \mathbf{r} is called Bloch vector. The set of all the Bloch vectors that constitute a density operator is known as the Bloch vector space $B(\mathbb{R}^{N^2-1})$.

A matrix of the form (2.5) is of unit trace and Hermitian, but it might not be positive. To guarantee the positivity restrictions must be imposed on the Bloch vector. It is shown that $B(\mathbb{R}^{N^2-1})$ is a subset of the ball $D_R(\mathbb{R}^{N^2-1})$ of radius $R = \sqrt{2(1 - \frac{1}{N})}$, which is the minimum ball containing it, and that the ball $D_r(\mathbb{R}^{N^2-1})$ of radius $r = \sqrt{\frac{2}{N(N-1)}}$ is included in $B(\mathbb{R}^{N^2-1})$ [70], that is,

$$D_r(\mathbb{R}^{N^2-1}) \subseteq B(\mathbb{R}^{N^2-1}) \subseteq D_R(\mathbb{R}^{N^2-1}).$$

Let the dimensions of systems A, B and C be $d_A = N_1, d_B = N_2$ and $d_C = N_3$ respectively. Any tripartite quantum states $\rho_{ABC} \in \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ can be written

as:

$$\begin{aligned}\rho_{ABC} &= I_{N_1} \otimes I_{N_2} \otimes M_0 + \sum_{i=1}^{N_1^2-1} \lambda_i(1) \otimes I_{N_2} \otimes M_i + \sum_{j=1}^{N_2^2-1} I_{N_1} \otimes \lambda_j(2) \otimes \widetilde{M}_j \\ &+ \sum_{i=1}^{N_1^2-1} \sum_{j=1}^{N_2^2-1} \lambda_i(1) \otimes \lambda_j(2) \otimes M_{ij},\end{aligned}\quad (2.6)$$

where $\lambda_i(1)$, $\lambda_j(2)$ are the generators of $SU(N_1)$ and $SU(N_2)$; M_i , \widetilde{M}_j and M_{ij} are operators of \mathcal{H}_C .

Theorem 2.2 *Let $\mathbf{r} \in \mathbb{R}^{N_1^2-1}$, $\mathbf{s} \in \mathbb{R}^{N_2^2-1}$ and $|\mathbf{r}| \leq \sqrt{\frac{2}{N_1(N_1-1)}}$, $|\mathbf{s}| \leq \sqrt{\frac{2}{N_2(N_2-1)}}$. For a tripartite quantum state $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ with representation (2.6), we have*

$$M_0 - \sum_{i=1}^{N_1^2-1} r_i M_i - \sum_{j=1}^{N_2^2-1} s_j \widetilde{M}_j + \sum_{i=1}^{N_1^2-1} \sum_{j=1}^{N_2^2-1} r_i s_j M_{ij} \geq 0. \quad (2.7)$$

[Proof] Since $\mathbf{r} \in \mathbb{R}^{N_1^2-1}$, $\mathbf{s} \in \mathbb{R}^{N_2^2-1}$ and $|\mathbf{r}| \leq \sqrt{\frac{2}{N_1(N_1-1)}}$, $|\mathbf{s}| \leq \sqrt{\frac{2}{N_2(N_2-1)}}$, we have that $A_1 \equiv \frac{1}{2}(\frac{2}{N_1}I - \sum_{i=1}^{N_1^2-1} r_i \lambda_i(1))$ and $A_2 \equiv \frac{1}{2}(\frac{2}{N_2}I - \sum_{j=1}^{N_2^2-1} s_j \lambda_j(2))$ are positive Hermitian operators. Let $A = \sqrt{A_1} \otimes \sqrt{A_2} \otimes I_{N_3}$. Then $A\rho A \geq 0$ and $(A\rho A)^\dagger = A\rho A$. The partial trace of $A\rho A$ over \mathcal{H}_A (and \mathcal{H}_B) should be also positive. Hence

$$\begin{aligned}0 &\leq \text{Tr}_{AB}[A\rho A] \\ &= \text{Tr}_{AB}[A_1 \otimes A_2 \otimes M_0 + \sum_i \sqrt{A_1} \lambda_i(1) \sqrt{A_1} \otimes A_2 \otimes M_i \\ &+ \sum_j A_1 \otimes \sqrt{A_2} \lambda_j(2) \sqrt{A_2} \otimes \widetilde{M}_j + \sum_{ij} \sqrt{A_1} \lambda_i(1) \sqrt{A_1} \otimes \sqrt{A_2} \lambda_j(2) \sqrt{A_2} \otimes M_{ij}] \\ &= M_0 - \sum_{i=1}^{N_1^2-1} r_i M_i - \sum_{j=1}^{N_2^2-1} s_j \widetilde{M}_j + \sum_{i=1}^{N_1^2-1} \sum_{j=1}^{N_2^2-1} r_i s_j M_{ij}.\end{aligned}$$

□

Formula (2.7) is valid for any tripartite states. By setting $\mathbf{s} = 0$ in (2.7), one can get a result for bipartite systems:

Corollary 2.2 *Let $\rho_{AB} \in \mathcal{H}_A \otimes \mathcal{H}_B$, which can be generally written as $\rho_{AB} = I_{N_1} \otimes M_0 + \sum_{j=1}^{N_1^2-1} \lambda_j \otimes M_j$, then for any $\mathbf{r} \in \mathbb{R}^{N_1^2-1}$ with $|\mathbf{r}| \leq \sqrt{\frac{2}{N_1(N_1-1)}}$, $M_0 - \sum_{j=1}^{N_1^2-1} r_j M_j \geq 0$.*

A separable tripartite state ρ_{ABC} can be written as

$$\rho_{ABC} = \sum_i p_i |\psi_i^A\rangle\langle\psi_i^A| \otimes |\phi_i^B\rangle\langle\phi_i^B| \otimes |\omega_i^C\rangle\langle\omega_i^C|.$$

From (2.5) it can also be represented as:

$$\begin{aligned} \rho_{ABC} &= \sum_i p_i \frac{1}{2} \left(\frac{2}{N_1} I_{N_1} + \sum_{k=1}^{N_1^2-1} a_i^{(k)} \lambda_k(1) \right) \otimes \frac{1}{2} \left(\frac{2}{N_2} I_{N_2} + \sum_{l=1}^{N_2^2-1} b_i^{(l)} \lambda_l(2) \right) \otimes |\omega_i^C\rangle\langle\omega_i^C| \\ &= I_{N_1} \otimes I_{N_2} \otimes \frac{1}{N_1 N_2} \sum_i p_i |\omega_i^C\rangle\langle\omega_i^C| \\ &\quad + \sum_{k=1}^{N_1^2-1} \lambda_k(1) \otimes I_{N_2} \otimes \frac{1}{2N_2} \sum_i a_i^{(k)} p_i |\omega_i^C\rangle\langle\omega_i^C| \\ &\quad + \sum_{l=1}^{N_2^2-1} I_{N_1} \otimes \lambda_l(2) \otimes \frac{1}{2N_1} \sum_i b_i^{(l)} p_i |\omega_i^C\rangle\langle\omega_i^C| \\ &\quad + \sum_k \sum_l \lambda_k(1) \otimes \lambda_l(2) \otimes \frac{1}{4} \sum_i a_i^{(k)} b_i^{(l)} p_i |\omega_i^C\rangle\langle\omega_i^C|, \end{aligned} \quad (2.8)$$

where $(a_i^{(1)}, a_i^{(2)} \dots, a_i^{(N_1^2-1)})$ and $(b_i^{(1)}, b_i^{(2)} \dots, b_i^{(N_2^2-1)})$ are real vectors on the Bloch sphere satisfying $|\vec{a}_i|^2 = \sum_{j=1}^{N_1^2-1} (a_i^{(j)})^2 = 2(1 - \frac{1}{N_1})$ and $|\vec{b}_i|^2 = \sum_{j=1}^{N_2^2-1} (b_i^{(j)})^2 = 2(1 - \frac{1}{N_2})$.

Comparing (2.6) with (2.8), we have

$$\begin{aligned} M_0 &= \frac{1}{N_1 N_2} \sum_i p_i |\omega_i^C\rangle\langle\omega_i^C|, & M_k &= \frac{1}{2N_2} \sum_i a_i^{(k)} p_i |\omega_i^C\rangle\langle\omega_i^C|, \\ \widetilde{M}_l &= \frac{1}{2N_1} \sum_i b_i^{(l)} p_i |\omega_i^C\rangle\langle\omega_i^C|, & M_{kl} &= \frac{1}{4} \sum_i a_i^{(k)} b_i^{(l)} p_i |\omega_i^C\rangle\langle\omega_i^C|. \end{aligned} \quad (2.9)$$

For any $(N_1^2 - 1) \times (N_1^2 - 1)$ real matrix $R(1)$ and $(N_2^2 - 1) \times (N_2^2 - 1)$ real matrix $R(2)$ satisfying $\frac{1}{(N_1-1)^2} I - R(1)^T R(1) \geq 0$ and $\frac{1}{(N_2-1)^2} I - R(2)^T R(2) \geq 0$, we define a new matrix

$$\mathcal{R} = \begin{pmatrix} R(1) & 0 & 0 \\ 0 & R(2) & 0 \\ 0 & 0 & T \end{pmatrix}, \quad (2.10)$$

where T is a transformation acting on an $(N_1^2 - 1) \times (N_2^2 - 1)$ matrix M by

$$T(M) = R(1) M R^T(2).$$

Using \mathcal{R} we define a new operator $\gamma_{\mathcal{R}}$,

$$\begin{aligned}\gamma_{\mathcal{R}}(\rho_{ABC}) &= I_{N_1} \otimes I_{N_2} \otimes M'_0 + \sum_{i=1}^{N_1^2-1} \lambda_i(1) \otimes I_{N_2} \otimes M'_i + \sum_{j=1}^{N_2^2-1} I_{N_1} \otimes \lambda_j(2) \otimes \widetilde{M}'_j \\ &+ \sum_{i=1}^{N_1^2-1} \sum_{j=1}^{N_2^2-1} \lambda_i(1) \otimes \lambda_j(2) \otimes M'_{ij},\end{aligned}\quad (2.11)$$

where $M'_0 = M_0$, $M'_k = \sum_{m=1}^{N_1^2-1} R_{km}(1)M_m$, $\widetilde{M}'_l = \sum_{n=1}^{N_2^2-1} R_{ln}(2)\widetilde{M}_n$ and $M'_{ij} = (T(M))_{ij} = (R(1)MR^T(2))_{ij}$.

Theorem 2.3 *If ρ_{ABC} is separable, then $\gamma_{\mathcal{R}}(\rho_{ABC}) \geq 0$.*

[Proof] From (2.9) and (2.11) we get

$$\begin{aligned}M'_0 &= M_0 = \frac{1}{N_1 N_2} \sum_i p_i |\omega_i^C\rangle \langle \omega_i^C|, \quad M'_k = \frac{1}{2N_2} \sum_{mi} R_{km}(1) a_i^{(m)} p_i |\omega_i^C\rangle \langle \omega_i^C|, \\ \widetilde{M}'_l &= \frac{1}{2N_1} \sum_{ni} R_{ln}(2) b_i^{(n)} p_i |\omega_i^C\rangle \langle \omega_i^C|, \quad M'_{kl} = \frac{1}{4} \sum_{mni} R_{km}(1) a_i^{(m)} R_{ln}(2) b_i^{(n)} p_i |\omega_i^C\rangle \langle \omega_i^C|.\end{aligned}$$

A straightforward calculation gives rise to

$$\begin{aligned}\gamma_{\mathcal{R}}(\rho_{ABC}) &= \sum_i p_i \frac{1}{2} \left(\frac{2}{N_1} I_{N_1} + \sum_{k=1}^{N_1^2-1} \sum_{m=1}^{N_1^2-1} R_{km}(1) a_i^{(m)} \lambda_k(1) \right) \\ &\otimes \frac{1}{2} \left(\frac{2}{N_2} I_{N_2} + \sum_{l=1}^{N_2^2-1} \sum_{n=1}^{N_2^2-1} R_{ln}(2) b_i^{(n)} \lambda_l(2) \right) \otimes |\omega_i^C\rangle \langle \omega_i^C|.\end{aligned}$$

As $\frac{1}{(N_1-1)^2} I - R(1)^T R(1) \geq 0$ and $\frac{1}{(N_2-1)^2} I - R(2)^T R(2) \geq 0$, we get

$$\begin{aligned}|\vec{a}_i|^2 &= |R(1)\vec{a}_i|^2 \leq \frac{1}{(N_1-1)^2} |\vec{a}_i|^2 = \frac{2}{N_1(N_1-1)}, \\ |\vec{b}_i|^2 &= |R(2)\vec{b}_i|^2 \leq \frac{1}{(N_2-1)^2} |\vec{b}_i|^2 = \frac{2}{N_2(N_2-1)}.\end{aligned}$$

Therefore $\gamma_{\mathcal{R}}(\rho_{ABC})$ is still a density operator, i.e. $\gamma_{\mathcal{R}}(\rho_{ABC}) \geq 0$. \square

Theorem 2.3 gives a necessary separability criterion for general tripartite systems. The result can be also applied to bipartite systems. Let $\rho_{AB} \in \mathcal{H}_A \otimes \mathcal{H}_B$, $\rho_{AB} = I_{N_1} \otimes M_0 + \sum_{j=1}^{N_1^2-1} \lambda_j \otimes M_j$. For any real $(N_1^2 - 1) \times (N_1^2 - 1)$ matrix \mathcal{R} satisfying $\frac{1}{(N_1-1)^2}I - \mathcal{R}^T \mathcal{R} \geq 0$ and any state ρ_{AB} , we define

$$\gamma_{\mathcal{R}}(\rho_{AB}) = I_{N_1} \otimes M_0 + \sum_{j=1}^{N_1^2-1} \lambda_j \otimes M'_j,$$

where $M'_j = \sum_k \mathcal{R}_{jk} M_k$.

Corollary 2.3 *For $\rho_{AB} \in \mathcal{H}_A \otimes \mathcal{H}_B$, if there exists an \mathcal{R} with $\frac{1}{(N_1-1)^2}I - \mathcal{R}^T \mathcal{R} \geq 0$ such that $\gamma_{\mathcal{R}}(\rho_{AB}) < 0$, then ρ_{AB} must be entangled.*

For $2 \times N$ systems, the above corollary is reduced to the results in [71]. As an example we consider the 3×3 isotropic states,

$$\rho_I = \frac{1-p}{9} I_3 \otimes I_3 + \frac{p}{3} \sum_{i,j=1}^3 |ii\rangle\langle jj| = I_3 \otimes \left(\frac{1}{9} I_3\right) + \sum_{i=1}^5 \lambda_i \otimes \left(\frac{p}{6} \lambda_i\right) - \sum_{i=6}^8 \lambda_i \otimes \left(\frac{p}{6} \lambda_i\right).$$

If we choose \mathcal{R} to be $\text{Diag}\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$, we get that ρ_I is entangled for $0.5 < p \leq 1$.

For tripartite case, we take the following $3 \times 3 \times 3$ mixed state as an example:

$$\rho = \frac{1-p}{27} I_{27} + p |\psi\rangle\langle\psi|,$$

where $|\psi\rangle = \frac{1}{\sqrt{3}}(|000\rangle + |111\rangle + |222\rangle)(\langle 000| + \langle 111| + \langle 222|)$. Taking $R(1) = R(2) = \text{Diag}\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$, we have that ρ is entangled for $0.6248 < p \leq 1$.

In fact the criterion for $2 \times N$ systems [71] is equivalent to the PPT criterion [72]. Similarly theorem 2.3 is also equivalent to the PPT criterion for $2 \times 2 \times N$ systems.

2.1.5 Covariance matrix criterion

In this subsection we study the separability problem by using the covariance matrix approach. We first give a brief review of covariance matrix criterion proposed in [35]. Let \mathcal{H}_d^A and \mathcal{H}_d^B be d -dimensional complex vector spaces, and ρ_{AB} a bipartite quantum state in $\mathcal{H}_d^A \otimes \mathcal{H}_d^B$. Let A_k (resp. B_k) be d^2 observables on \mathcal{H}_d^A (resp.

\mathcal{H}_d^B) such that they form an orthonormal normalized basis of the observable space, satisfying $\text{Tr}[A_k A_l] = \delta_{k,l}$ (resp. $\text{Tr}[B_k B_l] = \delta_{k,l}$). Consider the total set $\{M_k\} = \{A_k \otimes I, I \otimes B_k\}$. It can be proven that [30],

$$\sum_{k=1}^{N^2} (M_k)^2 = dI, \quad \sum_{k=1}^{N^2} \langle M_k \rangle^2 = \text{Tr}[\rho_{AB}^2]. \quad (2.12)$$

The covariance matrix γ is defined with entries

$$\gamma_{ij}(\rho_{AB}, \{M_k\}) = \frac{\langle M_i M_j \rangle + \langle M_j M_i \rangle}{2} - \langle M_i \rangle \langle M_j \rangle, \quad (2.13)$$

which has a block structure [35]:

$$\gamma = \begin{pmatrix} A & C \\ C^T & B \end{pmatrix}, \quad (2.14)$$

where $A = \gamma(\rho_A, \{A_k\})$, $B = \gamma(\rho_B, \{B_k\})$, $C_{ij} = \langle A_i \otimes B_j \rangle_{\rho_{AB}} - \langle A_i \rangle_{\rho_A} \langle B_j \rangle_{\rho_B}$, $\rho_A = \text{Tr}_B[\rho_{AB}]$, $\rho_B = \text{Tr}_A[\rho_{AB}]$. Such covariance matrix has a concavity property: for a mixed density matrix $\rho = \sum_k p_k \rho_k$ with $p_k \geq 0$ and $\sum_k p_k = 1$, one has $\gamma(\rho) \geq \sum_k p_k \gamma(\rho_k)$.

For a bipartite product state $\rho_{AB} = \rho_A \otimes \rho_B$, C in (2.14) is zero. Generally if ρ_{AB} is separable, then there exist states $|a_k\rangle\langle a_k|$ on \mathcal{H}_d^A , $|b_k\rangle\langle b_k|$ on \mathcal{H}_d^B and p_k such that

$$\gamma(\rho) \geq \kappa_A \oplus \kappa_B, \quad (2.15)$$

where $\kappa_A = \sum p_k \gamma(|a_k\rangle\langle a_k|, \{A_k\})$, $\kappa_B = \sum p_k \gamma(|b_k\rangle\langle b_k|, \{B_k\})$.

For a separable bipartite state, it has been shown that [35]

$$\sum_{i=1}^{d^2} |C_{ii}| \leq \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2}. \quad (2.16)$$

Criterion (2.16) depends on the choice of the orthonormal normalized basis of the observables. In fact the term $\sum_{i=1}^{d^2} |C_{ii}|$ has an upper bound $\|C\|_{KF}$ which is invariant under unitary transformation and can be attained by choosing proper local orthonormal observable basis, where $\|C\|_{KF}$ stands for the Ky Fan norm of C , $\|C\|_{KF} = \text{Tr}[\sqrt{CC^\dagger}]$, with \dagger denoting the transpose and conjugation. It has been shown in [32] that if ρ_{AB} is separable, then

$$\|C\|_{KF} \leq \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2}. \quad (2.17)$$

From the covariance matrix approach, we can also get an alternative criterion. From (2.14) and (2.15) we have that if ρ_{AB} is separable, then

$$X \equiv \begin{pmatrix} A - \kappa_A & C \\ C^T & B - \kappa_B \end{pmatrix} \geq 0. \quad (2.18)$$

Hence all the 2×2 minor submatrices of X must be positive. Namely one has

$$\begin{vmatrix} (A - \kappa_A)_{ii} & C_{ij} \\ C_{ji} & (B - \kappa_B)_{jj} \end{vmatrix} \geq 0,$$

i.e. $(A - \kappa_A)_{ii}(B - \kappa_B)_{jj} \geq C_{ij}^2$. Summing over all i, j and using (2.12), we get

$$\begin{aligned} \sum_{i,j=1}^{d^2} C_{i,j}^2 &\leq (\text{Tr}[A] - \text{Tr}[\kappa_A])(\text{Tr}[B] - \text{Tr}[\kappa_B]) \\ &= (d - \text{Tr}[\rho_A^2] - d + 1)(d - \text{Tr}[\rho_B^2] - d + 1) = (1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2]). \end{aligned}$$

That is

$$\|C\|_{HS}^2 \leq (1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2]), \quad (2.19)$$

where $\|C\|_{HS}$ stands for the Euclid norm of C , i.e. $\|C\|_{HS} = \sqrt{\text{Tr}[CC^\dagger]}$.

Formulae (2.17) and (2.19) are independent and could be complement. When

$$\sqrt{(1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2])} < \|C\|_{HS} \leq \|C\|_{KF} \leq \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2},$$

(2.19) can recognize the entanglement but (2.17) can not. When

$$\|C\|_{HS} \leq \sqrt{(1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2])} \leq \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2} < \|C\|_{KF},$$

(2.17) can recognize the entanglement while (2.19) not.

The separability criteria based on covariance matrix approach can be generalized to multipartite systems. We first consider the tripartite case, $\rho_{ABC} \in \mathcal{H}_d^A \otimes \mathcal{H}_d^B \otimes \mathcal{H}_d^C$. Take d^2 observables A_k on \mathcal{H}_A resp. B_k on \mathcal{H}_B resp. C_k on \mathcal{H}_C . Set $\{M_k\} = \{A_k \otimes I \otimes I, I \otimes B_k \otimes I, I \otimes I \otimes C_k\}$. The covariance matrix defined by (2.13) has then the following block structure:

$$\gamma = \begin{pmatrix} A & D & E \\ D^T & B & F \\ E^T & F^T & C \end{pmatrix}, \quad (2.20)$$

where $A = \gamma(\rho_A, \{A_k\})$, $B = \gamma(\rho_B, \{B_k\})$, $C = \gamma(\rho_C, \{C_k\})$, $D_{ij} = \langle A_i \otimes B_j \rangle_{\rho_{AB}} - \langle A_i \rangle_{\rho_A} \langle B_j \rangle_{\rho_B}$, $E_{ij} = \langle A_i \otimes C_j \rangle_{\rho_{AC}} - \langle A_i \rangle_{\rho_A} \langle C_j \rangle_{\rho_C}$, $F_{ij} = \langle B_i \otimes C_j \rangle_{\rho_{BC}} - \langle B_i \rangle_{\rho_B} \langle C_j \rangle_{\rho_C}$.

Theorem 2.4 *If ρ_{ABC} is fully separable, then*

$$\|D\|_{HS}^2 \leq (1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2]), \quad (2.21)$$

$$\|E\|_{HS}^2 \leq (1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_C^2]), \quad (2.22)$$

$$\|F\|_{HS}^2 \leq (1 - \text{Tr}[\rho_B^2])(1 - \text{Tr}[\rho_C^2]), \quad (2.23)$$

and

$$2\|D\|_{KF} \leq (1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2]), \quad (2.24)$$

$$2\|E\|_{KF} \leq (1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_C^2]), \quad (2.25)$$

$$2\|F\|_{KF} \leq (1 - \text{Tr}[\rho_B^2]) + (1 - \text{Tr}[\rho_C^2]). \quad (2.26)$$

[Proof] For a tripartite product state $\rho_{ABC} = \rho_A \otimes \rho_B \otimes \rho_C$, D , E and F in (2.20) are zero. If ρ_{ABC} is fully separable, then there exist states $|a_k\rangle\langle a_k|$ in \mathcal{H}_d^A , $|b_k\rangle\langle b_k|$ in \mathcal{H}_d^B and $|c_k\rangle\langle c_k|$ in \mathcal{H}_d^C , and p_k such that $\gamma(\rho) \geq \kappa_A \oplus \kappa_B \oplus \kappa_C$, where $\kappa_A = \sum p_k \gamma(|a_k\rangle\langle a_k|, \{A_k\})$, $\kappa_B = \sum p_k \gamma(|b_k\rangle\langle b_k|, \{B_k\})$ and $\kappa_C = \sum p_k \gamma(|c_k\rangle\langle c_k|, \{C_k\})$, i.e.

$$Y \equiv \begin{pmatrix} A - \kappa_A & D & E \\ D^T & B - \kappa_B & F \\ E^T & F^T & C - \kappa_C \end{pmatrix} \geq 0. \quad (2.27)$$

Thus all the 2×2 minor submatrices of Y must be positive. Selecting one with two rows and columns from the first two block rows and columns of Y , we have

$$\begin{vmatrix} (A - \kappa_A)_{ii} & D_{ij} \\ D_{ji} & (B - \kappa_B)_{jj} \end{vmatrix} \geq 0, \quad (2.28)$$

i.e. $(A - \kappa_A)_{ii}(B - \kappa_B)_{jj} \geq |D_{ij}|^2$. Summing over all i, j and using (2.12), we get

$$\begin{aligned} \|D\|_{HS}^2 &= \sum_{i,j=1}^d D_{i,j}^2 \leq (\text{Tr}[A] - \text{Tr}[\kappa_A])(\text{Tr}[B] - \text{Tr}[\kappa_B]) \\ &= (d - \text{Tr}[\rho_A^2] - d + 1)(d - \text{Tr}[\rho_B^2] - d + 1) = (1 - \text{Tr}[\rho_A^2])(1 - \text{Tr}[\rho_B^2]), \end{aligned}$$

which proves (2.21). (2.22) and (2.23) can be similarly proved.

From (2.28) we also have $(A - \kappa_A)_{ii} + (B - \kappa_B)_{ii} \geq 2|D_{ii}|$. Therefore

$$\begin{aligned} \sum_i |D_{ii}| &\leq \frac{(\text{Tr}[A] - \text{Tr}[\kappa_A]) + (\text{Tr}[B] - \text{Tr}[\kappa_B])}{2} \\ &= \frac{(d - \text{Tr}[\rho_A^2] - d + 1) + (d - \text{Tr}[\rho_B^2] - d + 1)}{2} \\ &= \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2}. \end{aligned} \quad (2.29)$$

Note that $\sum_{i=1}^{d^2} D_{ii} \leq \sum_{i=1}^{d^2} |D_{ii}|$. By using that $\text{Tr}[MU] \leq \|M\|_{KF} = \text{Tr}[\sqrt{MM^\dagger}]$ for any matrix M and any unitary U [73], we have $\sum_{i=1}^{d^2} D_{ii} \leq \|D\|_{KF}$.

Let $D = U^\dagger \Lambda V$ be the singular value decomposition of D . Make a transformation of the orthonormal normalized basis of the local orthonormal observable space: $\tilde{A}_i = \sum_l U_{il} A_l$ and $\tilde{B}_j = \sum_m V_{jm}^* B_m$. In the new basis we have

$$\tilde{D}_{ij} = \sum_{lm} U_{il} D_{lm} V_{jm} = (UDV^\dagger)_{ij} = \Lambda_{ij}. \quad (2.30)$$

Then (2.29) becomes

$$\sum_{i=1}^{d^2} \tilde{D}_{ii} = \|D\|_{KF} \leq \frac{(1 - \text{Tr}[\rho_A^2]) + (1 - \text{Tr}[\rho_B^2])}{2}$$

which proves (2.24). (2.25) and (2.26) can similarly be treated. \square

We consider now the case that ρ_{ABC} is bi-partite separable.

Theorem 2.5 *If ρ_{ABC} is a bi-partite separable state with respect to the bipartite partition of the sub-systems A and BC (resp. AB and C ; resp. AC and B), then (2.21), (2.22) and (2.24), (2.25) (resp. (2.22), (2.23) and (2.25), (2.26); resp. (2.21), (2.23) and (2.24), (2.26)) must hold.*

[Proof] We prove the case that ρ_{ABC} is bi-partite separable with respect to the A system and BC systems partition. The other cases can be similarly treated. In this case the matrices D and E in the covariance matrix (2.20) are zero. ρ_{ABC} takes the form $\rho_{ABC} = \sum p_m \rho_A^m \otimes \rho_{BC}^m$. Define $\kappa_A = \sum p_m \gamma(\rho_A^m, \{A_k\})$, $\kappa_{BC} = \sum p_m \gamma(\rho_{BC}^m, \{B_k \otimes I, I \otimes C_k\})$. κ_{BC} has a form

$$\kappa_{BC} = \begin{pmatrix} \kappa_B & F' \\ (F')^T & \kappa_C \end{pmatrix},$$

where $\kappa_B = \sum p_k \gamma(|b_k\rangle\langle b_k|, \{B_k\})$ and $\kappa_C = \sum p_k \gamma(|c_k\rangle\langle c_k|, \{C_k\})$, $(F')_{ij} = \sum_m p_m (\langle B_i \otimes C_j \rangle_{\rho_{BC}^m} - \langle B_i \rangle_{\rho_B^m} \langle C_j \rangle_{\rho_C^m})$. By using the concavity of covariance matrix we have

$$\gamma(\rho_{ABC}) \geq \sum_m p_m \gamma(\rho_A^m \otimes \rho_{BC}^m) = \begin{pmatrix} \kappa_A & 0 & 0 \\ 0 & \kappa_B & F' \\ 0 & (F')^T & \kappa_C \end{pmatrix}.$$

Accounting to the method used in proving Theorem 2, we get (2.21), (2.22) and (2.24), (2.25). \square

From Theorem 2.4 and 2.5 we have the following corollary.

Corollary 2.5 *If two of the inequalities (2.21), (2.22) and (2.23) (or (2.24), (2.25) and (2.26)) are violated, the state must be fully entangled.*

The result of Theorem 2.4 can be generalized to general multipartite case $\rho \in \mathcal{H}_d^{(1)} \otimes \mathcal{H}_d^{(2)} \otimes \dots \otimes \mathcal{H}_d^{(N)}$. Define $\hat{A}_\alpha^i = I \otimes I \otimes \dots \otimes \lambda_\alpha \otimes I \otimes \dots \otimes I$, where $\lambda_0 = I/d$, λ_α ($\alpha = 1, 2, \dots, d^2 - 1$) are the normalized generators of $SU(d)$ satisfying $\text{Tr}[\lambda_\alpha \lambda_\beta] = \delta_{\alpha\beta}$ and acting on the i^{th} system $\mathcal{H}_d^{(i)}$, $i = 1, 2, \dots, N$. Denote $\{M_k\}$ the set of all \hat{A}_α^i . Then the covariance matrix of ρ can be written as

$$\gamma(\rho) = \begin{pmatrix} \mathcal{A}_{11} & \mathcal{A}_{12} \cdots & \mathcal{A}_{1N} \\ \mathcal{A}_{12}^T & \mathcal{A}_{22} \cdots & \mathcal{A}_{2N} \\ \vdots & \vdots & \vdots \\ \mathcal{A}_{1N}^T & \mathcal{A}_{2N}^T \cdots & \mathcal{A}_{NN} \end{pmatrix}, \quad (2.31)$$

where $\mathcal{A}_{ii} = \gamma(\rho, \{\hat{A}_k^i\})$ and $(\mathcal{A}_{ij})_{mn} = \langle \hat{A}_m^i \otimes \hat{A}_n^j \rangle - \langle \hat{A}_m^i \rangle \langle \hat{A}_n^j \rangle$ for $i \neq j$.

For a product state $\rho_{12\dots N}$, \mathcal{A}_{ij} , $i \neq j$, in (2.31) are zero matrices. Define

$$\kappa_{\mathcal{A}_{ii}} = \sum_k p_k \gamma(|\psi_k^i\rangle\langle\psi_k^i|, \{\hat{A}_l^i\}). \quad (2.32)$$

Then for a fully separable multipartite state $\rho = \sum_k p_k |\psi_k^1\rangle\langle\psi_k^1| \otimes |\psi_k^2\rangle\langle\psi_k^2| \otimes \dots \otimes |\psi_k^N\rangle\langle\psi_k^N|$ one has

$$Z = \begin{pmatrix} \mathcal{A}_{11} - \kappa_{\mathcal{A}_{11}} & \mathcal{A}_{12} \cdots & \mathcal{A}_{1N} \\ \mathcal{A}_{12}^T & \mathcal{A}_{22} - \kappa_{\mathcal{A}_{22}} \cdots & \mathcal{A}_{2N} \\ \vdots & \vdots & \vdots \\ \mathcal{A}_{1N}^T & \mathcal{A}_{2N}^T \cdots & \mathcal{A}_{NN} - \kappa_{\mathcal{A}_{NN}} \end{pmatrix} \geq 0. \quad (2.33)$$

From which we have the following separability criterion for multipartite systems:

Theorem 2.6 *If a state $\rho \in \mathcal{H}_d^{(1)} \otimes \mathcal{H}_d^{(2)} \otimes \dots \otimes \mathcal{H}_d^{(N)}$ is fully separable, the following inequalities*

$$\|\mathcal{A}_{ij}\|_{HS}^2 \leq (1 - \text{Tr}[\rho_i^2])(1 - \text{Tr}[\rho_j^2]), \quad (2.34)$$

$$\|\mathcal{A}_{ij}\|_{KF} \leq \frac{(1 - \text{Tr}[\rho_i^2]) + (1 - \text{Tr}[\rho_j^2])}{2} \quad (2.35)$$

must be fulfilled for any $i \neq j$.

2.2 Normal form of quantum states

In this subsection we show that the correlation matrix (CM) criterion can be improved from the normal form obtained under filtering transformations. Based on CM criterion entanglement witness in terms of local orthogonal observables (LOOs) [74] for both bipartite and multipartite systems can be also constructed.

For bipartite case, $\rho \in \mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$ with $\dim \mathcal{H}_A = M$, $\dim \mathcal{H}_B = N$, $M \leq N$, is mapped to the following form under local filtering transformations [75]:

$$\rho \rightarrow \tilde{\rho} = \frac{(F_A \otimes F_B)\rho(F_A \otimes F_B)^\dagger}{\text{Tr}[(F_A \otimes F_B)\rho(F_A \otimes F_B)^\dagger]}, \quad (2.36)$$

where $F_{A/B} \in GL(M/N, \mathbb{C})$ are arbitrary invertible matrices. This transformation is also known as stochastic local operations assisted by classical communication (SLOCC). By the definition it is obvious that filtering transformation will preserve the separability of a quantum state.

It has been shown that under local filtering operations one can transform a strictly positive ρ into a normal form [76],

$$\tilde{\rho} = \frac{(F_A \otimes F_B)\rho(F_A \otimes F_B)^\dagger}{\text{Tr}[(F_A \otimes F_B)\rho(F_A \otimes F_B)^\dagger]} = \frac{1}{MN} \left(I + \sum_{i=1}^{M^2-1} \xi_i G_i^A \otimes G_i^B \right), \quad (2.37)$$

where $\xi_i \geq 0$, G_i^A and G_i^B are some traceless orthogonal observables. The matrices F_A and F_B can be obtained by minimizing the function

$$f(A, B) = \frac{\text{Tr}[\rho(A \otimes B)]}{(\det A)^{1/M}(\det B)^{1/N}}, \quad (2.38)$$

where $A = F_A^\dagger F_A$ and $B = F_B^\dagger F_B$. In fact, one can choose $F_A^0 \equiv |\det(\rho_A)|^{1/2M} (\sqrt{\rho_A})^{-1}$, and $F_B^0 \equiv |\det(\rho_B')|^{1/2N} (\sqrt{\rho_B'})^{-1}$, where $\rho_B' = \text{Tr}_A[I \otimes (\sqrt{\rho_A})^{-1} \rho I \otimes (\sqrt{\rho_A})^{-1}]$. Then by iterations one can get the optimal A and B. In particular, there is a matlab code available in [77].

For bipartite separable states ρ , the CM separability criterion [78] says that

$$\|T\|_{KF} \leq \sqrt{MN(M-1)(N-1)}, \quad (2.39)$$

where T is an $(M^2 - 1) \times (N^2 - 1)$ matrix with $T_{ij} = MN \cdot \text{Tr}[\rho \lambda_i^A \otimes \lambda_j^B]$, $\|T\|_{KF}$ stands for the trace norm of T , $\lambda_k^{A/B}$ s are the generators of $SU(M/N)$ and have been chosen to be normalized, $\text{Tr}[\lambda_k^{(A/B)} \lambda_l^{(A/B)}] = \delta_{kl}$.

As the filtering transformation does not change the separability of a state, one can study the separability of $\tilde{\rho}$ instead of ρ . Under the normal form (2.37) the criterion (2.39) becomes

$$\sum_i \xi_i \leq \sqrt{MN(M-1)(N-1)}. \quad (2.40)$$

In [30] a separability criterion based on local uncertainty relation (LUR) has been obtained. It says that for any separable state ρ ,

$$1 - \sum_k \langle G_k^A \otimes G_k^B \rangle - \frac{1}{2} \langle G_k^A \otimes I - I \otimes G_k^B \rangle^2 \geq 0, \quad (2.41)$$

where $G_k^{A/B}$ s are LOOs such as the normalized generators of $SU(M/N)$ and $G_k^A = 0$ for $k = M^2 + 1, \dots, N^2$. The criterion is shown to be strictly stronger than the realignment criterion [47]. Under the normal form (2.37) criterion (2.41) becomes

$$\begin{aligned} 1 & - \sum_k \langle G_k^A \otimes G_k^B \rangle - \frac{1}{2} \langle G_k^A \otimes I - I \otimes G_k^B \rangle^2 \\ & = 1 - \frac{1}{MN} \sum_k \xi_k - \frac{1}{2} \left(\frac{1}{M} + \frac{1}{N} \right) \geq 0, \end{aligned}$$

i.e.

$$\sum_k \xi_k \leq MN - \frac{M+N}{2}. \quad (2.42)$$

As $\sqrt{MN(M-1)(N-1)} \leq MN - \frac{M+N}{2}$ holds for any M and N , from (2.40) and (2.42) it is obvious that the CM criterion recognizes entanglement better when the normal form is taken into account.

We now consider multipartite systems. Let ρ be a strictly positive density matrix in $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$, $\dim \mathcal{H}_i = d_i$. ρ can be generally expressed in terms of the $SU(n)$ generators λ_{α_k} [79],

$$\begin{aligned} \rho & = \frac{1}{\prod_i^N d_i} \left(\otimes_j^N I_{d_j} + \sum_{\{\mu_1\}} \sum_{\alpha_1} \mathcal{T}_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_1}^{\{\mu_1\}} + \sum_{\{\mu_1\mu_2\}} \sum_{\alpha_1\alpha_2} \mathcal{T}_{\alpha_1\alpha_2}^{\{\mu_1\mu_2\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \right. \\ & + \sum_{\{\mu_1\mu_2\mu_3\}} \sum_{\alpha_1\alpha_2\alpha_3} \mathcal{T}_{\alpha_1\alpha_2\alpha_3}^{\{\mu_1\mu_2\mu_3\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \lambda_{\alpha_3}^{\{\mu_3\}} \\ & + \dots + \sum_{\{\mu_1\mu_2\dots\mu_M\}} \sum_{\alpha_1\alpha_2\dots\alpha_M} \mathcal{T}_{\alpha_1\alpha_2\dots\alpha_M}^{\{\mu_1\mu_2\dots\mu_M\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \dots \lambda_{\alpha_M}^{\{\mu_M\}} \\ & \left. + \dots + \sum_{\alpha_1\alpha_2\dots\alpha_N} \mathcal{T}_{\alpha_1\alpha_2\dots\alpha_N}^{\{1,2,\dots,N\}} \lambda_{\alpha_1}^{\{1\}} \lambda_{\alpha_2}^{\{2\}} \dots \lambda_{\alpha_N}^{\{N\}} \right), \end{aligned} \quad (2.43)$$

where $\lambda_{\alpha_k}^{\{\mu_k\}} = I_{d_1} \otimes I_{d_2} \otimes \cdots \otimes \lambda_{\alpha_k} \otimes I_{d_{\mu_k+1}} \otimes \cdots \otimes I_{d_N}$ with λ_{α_k} appears at the μ_k th position and

$$\mathcal{T}_{\alpha_1 \alpha_2 \cdots \alpha_M}^{\{\mu_1 \mu_2 \cdots \mu_M\}} = \frac{\prod_{i=1}^M d_{\mu_i}}{2^M} \text{Tr}[\rho \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \cdots \lambda_{\alpha_M}^{\{\mu_M\}}].$$

The generalized CM criterion says that: if ρ in (2.43) is fully separable, then

$$\|\mathcal{T}^{\{\mu_1, \mu_2, \dots, \mu_M\}}\|_{KF} \leq \sqrt{\frac{1}{2^M} \prod_{k=1}^M d_{\mu_k} (d_{\mu_k} - 1)}, \quad (2.44)$$

for $2 \leq M \leq N$, $\{\mu_1, \mu_2, \dots, \mu_M\} \subset \{1, 2, \dots, N\}$. The KF norm is defined by

$$\|\mathcal{T}^{\{\mu_1, \mu_2, \dots, \mu_M\}}\|_{KF} = \max_{m=1, 2, \dots, M} \|\mathcal{T}_{(m)}\|_{KF},$$

where $\mathcal{T}_{(m)}$ is a kind of matrix unfolding of $\mathcal{T}^{\{\mu_1, \mu_2, \dots, \mu_M\}}$.

The criterion (2.44) can be improved by investigating the normal form of (2.43).

Theorem 2.7 *By filtering transformations of the form*

$$\tilde{\rho} = F_1 \otimes F_2 \otimes \cdots \otimes F_N \rho F_1^\dagger \otimes F_2^\dagger \otimes F_N^\dagger, \quad (2.45)$$

where $F_i \in GL(d_i, \mathbb{C})$, $i = 1, 2, \dots, N$, followed by normalization, any strictly positive state ρ can be transformed into a normal form

$$\begin{aligned} \rho = & \frac{1}{\prod_i^N d_i} \left(\otimes_j^N I_{d_j} + \sum_{\{\mu_1 \mu_2\}} \sum_{\alpha_1 \alpha_2} \mathcal{T}_{\alpha_1 \alpha_2}^{\{\mu_1 \mu_2\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \right. \\ & + \sum_{\{\mu_1 \mu_2 \mu_3\}} \sum_{\alpha_1 \alpha_2 \alpha_3} \mathcal{T}_{\alpha_1 \alpha_2 \alpha_3}^{\{\mu_1 \mu_2 \mu_3\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \lambda_{\alpha_3}^{\{\mu_3\}} \\ & + \cdots + \sum_{\{\mu_1 \mu_2 \cdots \mu_M\}} \sum_{\alpha_1 \alpha_2 \cdots \alpha_M} \mathcal{T}_{\alpha_1 \alpha_2 \cdots \alpha_M}^{\{\mu_1 \mu_2 \cdots \mu_M\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \cdots \lambda_{\alpha_M}^{\{\mu_M\}} \\ & \left. + \cdots + \sum_{\alpha_1 \alpha_2 \cdots \alpha_N} \mathcal{T}_{\alpha_1 \alpha_2 \cdots \alpha_N}^{\{1, 2, \dots, N\}} \lambda_{\alpha_1}^{\{1\}} \lambda_{\alpha_2}^{\{2\}} \cdots \lambda_{\alpha_N}^{\{N\}} \right). \end{aligned} \quad (2.46)$$

[Proof] Let D_1, D_2, \dots, D_N be the sets of density matrices of the N subsystems. The cartesian product $D_1 \times D_2 \times \cdots \times D_N$ consisting of all product density matrices $\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N$ with normalization $\text{Tr}[\rho_i] = 1$, $i = 1, 2, \dots, N$, is a compact set of matrices on the full Hilbert space \mathcal{H} . For the given density matrix ρ we define the following function of ρ_i

$$f(\rho_1, \rho_2, \dots, \rho_N) = \frac{\text{Tr}[\rho(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N)]}{\prod_{i=1}^N \det(\rho_i)^{1/d_i}}.$$

The function is well-defined on the interior of $D_1 \times D_2 \times \cdots \times D_N$ where $\det \rho_i > 0$. As ρ is assumed to be strictly positive, we have $\text{Tr}[\rho(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N)] > 0$. Since $D_1 \times D_2 \times \cdots \times D_N$ is compact, we have $\text{Tr}[\rho(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N)] \geq C > 0$ with a lower bound C depending on ρ .

It follows that $f \rightarrow \infty$ on the boundary of $D_1 \times D_2 \times \cdots \times D_N$ where at least one of the ρ_i s satisfies that $\det \rho_i = 0$. It follows further that f has a positive minimum on the interior of $D_1 \times D_2 \times \cdots \times D_N$ with the minimum value attained for at least one product density matrix $\tau_1 \otimes \tau_2 \otimes \cdots \otimes \tau_N$ with $\det \tau_i > 0$, $i = 1, 2, \dots, N$. Any positive density matrix τ_i with $\det \tau_i > 0$ can be factorized in terms of Hermitian matrices F_i as

$$\tau_i = F_i^\dagger F_i \quad (2.47)$$

where $F_i \in GL(d_i, \mathbb{C})$. Denote $F = F_1 \otimes F_2 \otimes \cdots \otimes F_N$, so that $\tau_1 \otimes \tau_2 \otimes \cdots \otimes \tau_N = F^\dagger F$. Set $\tilde{\rho} = F \rho F^\dagger$ and define

$$\begin{aligned} \tilde{f}(\rho_1, \rho_2, \dots, \rho_N) &= \frac{\text{Tr}[\tilde{\rho}(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N)]}{\prod_{i=1}^N \det(\rho_i)^{1/d_i}} \\ &= \prod_{i=1}^N \det(\tau_i)^{1/d_i} \cdot \frac{\text{Tr}[\rho(F_1^\dagger \rho_1 F_1 \otimes F_2^\dagger \rho_2 F_2 \otimes \cdots \otimes F_N^\dagger \rho_N F_N)]}{\prod_{i=1}^N \det(\tau_i)^{1/d_i} \det(\rho_i)^{1/d_i}} \\ &= \prod_{i=1}^N \det(\tau_i)^{1/d_i} \cdot f(F_1^\dagger \rho_1 F_1, F_2^\dagger \rho_2 F_2, \dots, F_N^\dagger \rho_N F_N). \end{aligned}$$

We see that when $F_i^\dagger \rho_i F_i = \tau_i$, \tilde{f} has a minimum and

$$\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N = (F^\dagger)^{-1} \tau_1 \otimes \tau_2 \otimes \cdots \otimes \tau_N F^{-1} = I.$$

Since \tilde{f} is stationary under infinitesimal variations about the minimum it follows that

$$\text{Tr}[\tilde{\rho} \delta(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N)] = 0$$

for all infinitesimal variations,

$$\begin{aligned} \delta(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_N) &= \delta \rho_1 \otimes I_{d_2} \otimes \cdots \otimes I_{d_N} + I_{d_1} \otimes \delta \rho_2 \otimes I_{d_3} \otimes \cdots \otimes I_{d_N} \\ &\quad + \cdots + I_{d_1} \otimes I_{d_2} \otimes \cdots \otimes I_{d_{N-1}} \otimes \delta \rho_N, \end{aligned}$$

subjected to the constraint $\det(I_{d_i} + \delta \rho_i) = 1$, which is equivalent to $\text{Tr}[\delta \rho_i] = 0$, $i = 1, 2, \dots, N$, using $\det(e^A) = e^{\text{Tr}[A]}$ for a given matrix A . Thus, $\delta \rho_i$ can be represented by the SU generators, $\delta \rho_i = \sum_k \delta c_k^i \lambda_k^i$. It follows that $\text{Tr}[\tilde{\rho} \lambda_{\alpha_k}^{\{\mu_k\}}] = 0$ for any α_k and μ_k . Hence the terms proportional to $\lambda_{\alpha_k}^{\{\mu_k\}}$ in (2.43) disappear. \square

Corollary 2.7 *The normal form of a product state in \mathcal{H} must be proportional to the identity.*

[Proof] Let ρ be such a state. From (2.46), we get that

$$\tilde{\rho}_i = \text{Tr}_{1,2,\dots,i-1,i+1,\dots,N}[\rho] = \frac{1}{d_i} I_{d_i}. \quad (2.48)$$

Therefore for a product state ρ we have

$$\tilde{\rho} = \rho_1 \otimes \rho_2 \otimes \dots \otimes \rho_N = \frac{1}{\prod_{i=1}^N d_i} \otimes_{i=1}^N I_{d_i}.$$

□

As an example for separability of multipartite states in terms of their normal forms (2.46), we consider the PPT entangled edge state [63]

$$\rho = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & b & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & c & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{c} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{b} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{a} & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad (2.49)$$

mixed with noises:

$$\rho_p = p\rho + \frac{(1-p)}{8} I_8.$$

Select $a = 2, b = 3$, and $c = 0.6$. Using the criterion in [79] we get that ρ_p is entangled for $0.92744 < p \leq 1$. But after transforming ρ_p to its normal form (2.46), the criterion can detect entanglement for $0.90285 < p \leq 1$.

Here we indicate that the filtering transformation does not change the PPT property. Let $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B$ be PPT, i.e. $\rho^{T_A} \geq 0$, and $\rho^{T_B} \geq 0$. Let $\tilde{\rho}$ be the normal form of ρ . From (2.36) we have

$$\tilde{\rho}^{T_A} = \frac{(F_A^* \otimes F_B) \rho^{T_A} (F_A^T \otimes F_B^\dagger)}{\text{Tr}[(F_A \otimes F_B) \rho (F_A \otimes F_B)^\dagger]}.$$

For any vector $|\psi\rangle$, we have

$$\langle \psi | \tilde{\rho}^{T_A} | \psi \rangle = \frac{\langle \psi | (F_A^* \otimes F_B) \rho^{T_A} (F_A^T \otimes F_B^\dagger) | \psi \rangle}{\text{Tr}[(F_A \otimes F_B) \rho (F_A \otimes F_B)^\dagger]} \equiv \langle \psi' | \rho^{T_A} | \psi' \rangle \geq 0,$$

where $|\psi'\rangle = \frac{(F_A^T \otimes F_B^\dagger) |\psi\rangle}{\sqrt{\text{Tr}[(F_A \otimes F_B) \rho (F_A \otimes F_B)^\dagger]}}$. $\tilde{\rho}^{T_B} \geq 0$ can be proved similarly. This property is also valid for multipartite case. Hence a bound entangled state will be bound entangled under filtering transformations.

2.3 Entanglement witness based on correlation matrix criterion

Entanglement witness (EW) is another way to describe separability. Based on CM criterion we can further construct entanglement witness in terms of LOOs. EW [74] is an observable of the composite system such that (i) nonnegative expectation values in all separable states, (ii) at least one negative eigenvalue (can recognize at least one entangled state). Consider bipartite systems in $\mathcal{H}_A^M \otimes \mathcal{H}_B^N$ with $M \leq N$.

Theorem 2.8 For any LOOs G_k^A and G_k^B ,

$$W = I - \alpha \sum_{k=0}^{N^2-1} G_k^A \otimes G_k^B$$

is an EW, where $\alpha = \frac{\sqrt{MN}}{\sqrt{(M-1)(N-1)+1}}$ and

$$G_0^A = \frac{1}{\sqrt{M}} I_M, \quad G_0^B = \frac{1}{\sqrt{N}} I_N. \quad (2.50)$$

[Proof] Let $\rho = \sum_{l,m=0}^{N^2-1} T_{lm} \lambda_l^A \otimes \lambda_m^B$ be a separable state, where $\lambda_k^{A/B}$ are normalized generators of $SU(M/N)$ with $\lambda_0^A = \frac{1}{\sqrt{M}} I_M$, $\lambda_0^B = \frac{1}{\sqrt{N}} I_N$. Any other LOOs $G_k^{A/B}$ fulfill (2.50) can be obtained from these λ s through orthogonal transformations $\mathcal{O}^{A/B}$, $G_k^{A/B} = \sum_{l=0}^{N^2-1} \mathcal{O}_{kl}^{A/B} \lambda_l$, where $\mathcal{O}^{A/B} = \begin{pmatrix} 1 & 0 \\ 0 & \mathcal{R}^{A/B} \end{pmatrix}$, $\mathcal{R}^{A/B}$ are $(N^2 - 1) \times (N^2 - 1)$ orthogonal matrices. We have

$$\begin{aligned} \text{Tr}[\rho W] &= 1 - \alpha \frac{1}{\sqrt{MN}} - \alpha \sum_{k=1}^{N^2-1} \sum_{l,m=1}^{N^2-1} \mathcal{R}_{kl}^A \mathcal{R}_{km}^B \text{Tr}[\rho(\lambda_l^A \otimes \lambda_m^B)] \\ &= \frac{\sqrt{(M-1)(N-1)}}{\sqrt{(M-1)(N-1)+1}} - \frac{1}{\sqrt{MN}(\sqrt{(M-1)(N-1)+1})} \sum_{k=1}^{N^2-1} \sum_{l,m=1}^{N^2-1} \mathcal{R}_{kl}^A T_{lm} \mathcal{R}_{km}^B \\ &\geq \frac{\sqrt{MN(M-1)(N-1)} - \|T\|_{KF}}{\sqrt{MN}(\sqrt{(M-1)(N-1)+1})} \geq 0, \end{aligned}$$

where we have used $\text{Tr}[\mathcal{R}T] \leq \|T\|_{KF}$ for any unitary \mathcal{R} in the first inequality and the CM criterion in the second inequality.

Now let $\rho = \frac{1}{MN} (I_{MN} + \sum_{i=1}^{M^2-1} s_i \lambda_i^A \otimes I_N + \sum_{j=1}^{N^2-1} r_j I_M \otimes \lambda_j^B + \sum_{i=1}^{M^2-1} \sum_{j=1}^{N^2-1} T_{ij} \lambda_i^A \otimes \lambda_j^B)$ be a state in $\mathcal{H}_A^M \otimes \mathcal{H}_B^N$ which violates the CM criterion. Denote $\sigma_k(T)$ the singular

values of T . By singular value decomposition, one has $T = U^\dagger \Lambda V^*$, where Λ is a diagonal matrix with $\Lambda_{kk} = \sigma_k(T)$. Now choose LOOs to be $G_k^A = \sum_l U_{kl} \lambda_l^A$, $G_k^B = \sum_m V_{km} \lambda_m^B$ for $k = 1, 2, \dots, N^2 - 1$ and $G_0^A = \frac{1}{M} I_M, G_0^B = \frac{1}{N} I_N$. We obtain

$$\begin{aligned} \text{Tr}[\rho W] &= 1 - \alpha \frac{1}{\sqrt{MN}} - \alpha \sum_{k=1}^{N^2-1} \sum_{l,m=1}^{N^2-1} U_{kl} V_{km} \text{Tr}[\rho(\lambda_l^A \otimes \lambda_m^B)] \\ &= \frac{\sqrt{(M-1)(N-1)}}{\sqrt{(M-1)(N-1)+1}} - \frac{1}{\sqrt{MN}(\sqrt{(M-1)(N-1)+1})} \text{Tr}[UTV^T] \\ &= \frac{\sqrt{MN(M-1)(N-1)} - \|T\|_{KF}}{\sqrt{MN}(\sqrt{(M-1)(N-1)+1})} < 0 \end{aligned}$$

where the CM criterion has been used in the last step. \square

As the CM criterion can be generalized to multipartite form [79], we can also define entanglement witness for multipartite system in $\mathcal{H}_1^{d_1} \otimes \mathcal{H}_2^{d_2} \otimes \dots \otimes \mathcal{H}_N^{d_N}$. Set $d(M) = \max\{d_{\mu_i}, i = 1, 2, \dots, M\}$. Choose LOOs $G_k^{\{\mu_i\}}$ for $0 \leq k \leq d(M)^2 - 1$ with $G_0^{\{\mu_i\}} = \frac{1}{d_{\mu_i}} I_{d_{\mu_i}}$ and define

$$W^{(M)} = I - \beta^{(M)} \sum_{k=0}^{d(M)^2-1} G_k^{\{\mu_1\}} \otimes G_k^{\{\mu_2\}} \otimes \dots \otimes G_k^{\{\mu_M\}}, \quad (2.51)$$

where $\beta^{(M)} = \frac{\sqrt{\prod_{i=1}^M d_{\mu_i}}}{1 + \sqrt{\prod_{i=1}^M (d_{\mu_i} - 1)}}$, $2 \leq M \leq N$. One can prove that (2.51) is an EW candidate for multipartite states. First we assume $\|\mathcal{T}^{(M)}\|_{KF} = \|\mathcal{T}_{(m_0)}\|_{KF}$. Note that for any $\mathcal{T}_{(m_0)}$, there must exist an elementary transformation P such that $\sum_{k=1}^{d(M)^2-1} \mathcal{T}_{kk \dots k}^{\{\mu_1 \mu_2 \dots \mu_M\}} = \text{Tr}[\mathcal{T}_{(m_0)} P]$. Then for an N-partite separable state we have

$$\begin{aligned} \text{Tr}[\rho W^{(M)}] &= 1 - \beta^{(M)} \frac{1}{\sqrt{\prod_{i=1}^M d_{\mu_i}}} - \beta^{(M)} \frac{1}{\prod_{i=1}^M d_{\mu_i}} \text{Tr}[\mathcal{T}_{(m_0)} P] \\ &\geq 1 - \beta^{(M)} \frac{1}{\sqrt{\prod_{i=1}^M d_{\mu_i}}} - \beta^{(M)} \frac{1}{\prod_{i=1}^M d_{\mu_i}} \|\mathcal{T}_{(m_0)}\|_{KF} \\ &\geq 1 - \beta^{(M)} \frac{1}{\sqrt{\prod_{i=1}^M d_{\mu_i}}} - \beta^{(M)} \frac{1}{\prod_{i=1}^M d_{\mu_i}} \sqrt{\prod_{k=1}^M d_{\mu_k} (d_{\mu_k} - 1)} \\ &= 0 \end{aligned}$$

for any $2 \leq M \leq N$, where we have taken into account that P is orthogonal and $\text{Tr}[MU] \leq \|M\|_{KF}$ for any unitary U at the first inequality. The second inequality is due to the generalized CM criterion.

By choosing proper LOOs it is also easy to show that $W^{(M)}$ has negative eigenvalues. For example for three qubits case, taking the normalized pauli matrices as LOOs, one find a negative eigenvalue of $W^{(M)}$, $(1 - \sqrt{3})/2$.

3 Concurrence and Tangle

In this section, we focus on two important measures: concurrence and tangle (see, [80]). An elegant formula for concurrence of two-qubit states is derived analytically by Wootters [39, 81]. This quantity has recently been shown to play an essential role in describing quantum phase transition in various interacting quantum many-body systems [82] and may affect macroscopic properties of solids significantly [83]. Furthermore, concurrence also provides an estimation [84] for the entanglement of formation (EOF) [61], which quantifies the required minimally physical resources to prepare a quantum state.

Let \mathcal{H}_A (resp. \mathcal{H}_B) be an M (resp. N)-dimensional complex vector space with $|i\rangle$, $i = 1, \dots, M$ (resp. $|j\rangle$, $j = 1, \dots, N$), as an orthonormal basis. A general pure state on $\mathcal{H}_A \otimes \mathcal{H}_B$ is of the form

$$|\Psi\rangle = \sum_{i=1}^M \sum_{j=1}^N a_{ij} |i\rangle \otimes |j\rangle, \quad (3.52)$$

where $a_{ij} \in \mathbb{C}$ satisfy the normalization $\sum_{i=1}^M \sum_{j=1}^N a_{ij} a_{ij}^* = 1$.

The concurrence of (3.52) is defined by [85, 16]

$$C(|\psi\rangle) = \sqrt{2(1 - \text{Tr}[\rho_A^2])}, \quad (3.53)$$

where $\rho_A = \text{Tr}_B[|\psi\rangle\langle\psi|]$. The definition is extended to general mixed states $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ by the convex roof,

$$C(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C(|\psi_i\rangle). \quad (3.54)$$

For two qubits systems, the concurrence of $|\Psi\rangle$ is given by:

$$C(|\Psi\rangle) = |\langle\Psi|\tilde{\Psi}\rangle| = 2|a_{11}a_{22} - a_{12}a_{21}|, \quad (3.55)$$

where $|\tilde{\Psi}\rangle = \sigma_y \otimes \sigma_y |\Psi^*\rangle$, $|\Psi^*\rangle$ is the complex conjugate of $|\Psi\rangle$, σ_y is the Pauli matrix, $\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

For a mixed two-qubit quantum state ρ , the entanglement of formation $E(\rho)$ has a simple relation with the concurrence [39, 81]

$$E(\rho) = h\left(\frac{1 + \sqrt{1 - C(\rho)^2}}{2}\right),$$

where $h(x) = -x \log_2 x - (1 - x) \log_2(1 - x)$,

$$C(\rho) = \max\{\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0\}, \quad (3.56)$$

where the λ_i s are the eigenvalues, in decreasing order, of the Hermitian matrix $\sqrt{\sqrt{\rho}\tilde{\rho}\sqrt{\rho}}$ and $\tilde{\rho} = (\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$.

Another entanglement measure called tangle is defined by

$$\tau(|\psi\rangle) = C^2(|\psi\rangle) = 2(1 - \text{Tr}[\rho_A^2]) \quad (3.57)$$

for a pure state $|\psi\rangle$. For mixed state $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$, the definition is given by

$$\tau(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \tau(|\psi_i\rangle). \quad (3.58)$$

For multipartite state $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, $\dim\mathcal{H}_i = d_i$, $i = 1, \dots, N$, the concurrence of $|\psi\rangle$ is defined by [86]

$$C_N(|\psi\rangle\langle\psi|) = 2^{1-\frac{N}{2}} \sqrt{(2^N - 2) - \sum_{\alpha} \text{Tr}[\rho_{\alpha}^2]}, \quad (3.59)$$

where α labels all different reduced density matrices.

Up to constant factor (3.59) can be also expressed in another way. Let H denotes a d -dimensional vector space with basis $|i\rangle$, $i = 1, 2, \dots, d$. An N -partite pure state in $H \otimes \cdots \otimes H$ is generally of the form,

$$|\Psi\rangle = \sum_{i_1, i_2, \dots, i_N=1}^d a_{i_1, i_2, \dots, i_N} |i_1, i_2, \dots, i_N\rangle, \quad a_{i_1, i_2, \dots, i_N} \in \mathbb{C}. \quad (3.60)$$

Let α and α' (resp. β and β') be subsets of the subindices of a , associated to the same sub Hilbert spaces but with different summing indices. α (or α') and β (or β') span the whole space of the given sub-index of a . The generalized concurrence of $|\Psi\rangle$ is then given by [16]

$$C_d^N(|\Psi\rangle) = \sqrt{\frac{d}{2m(d-1)} \sum_p \sum_{\{\alpha, \alpha', \beta, \beta'\}} |a_{\alpha\beta} a_{\alpha'\beta'} - a_{\alpha\beta'} a_{\alpha'\beta}|^2}, \quad (3.61)$$

where $m = 2^{N-1} - 1$, \sum_p stands for the summation over all possible combinations of the indices of α and β .

For a mixed multipartite quantum state, $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ in $\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, the corresponding concurrence is given by the convex roof:

$$C_N(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C_N(|\psi_i\rangle). \quad (3.62)$$

3.1 Lower and upper bounds of concurrence

Calculation of the concurrence for general mixed states are extremely difficult. However, one can try to find the lower and the upper bounds to estimate the exact values of the concurrence [47, 51, 50, 32].

3.1.1 Lower bound of concurrence from covariance matrix criterion

In [47] a lower bound of $C(\rho)$ has been obtained,

$$C(\rho) \geq \sqrt{\frac{2}{M(M-1)}} [Max(\|T_A(\rho)\|, \|R(\rho)\|) - 1], \quad (3.63)$$

where T_A and R stand for partial transpose with respect to subsystem A and the realignment respectively. This bound is further improved based on local uncertainty relations [50],

$$C(\rho) \geq \frac{M + N - 2 - \sum_i \Delta_\rho^2(G_i^A \otimes I + I \otimes G_i^B)}{\sqrt{2M(M-1)}}, \quad (3.64)$$

where G_i^A and G_i^B are any set of local orthonormal observables, $\Delta_\rho^2(X) = \text{Tr}[X^2\rho] - (\text{Tr}[X\rho])^2$.

Bound (3.64) again depends on the choice of the local orthonormal observables. This bound can be optimized, in the sense that a local orthonormal observable-independent up bound of the right hand side of (3.64) can be obtained.

Theorem 3.1 *Let ρ be a bipartite state in $\mathcal{H}_M^A \otimes \mathcal{H}_N^B$. Then $C(\rho)$ satisfies*

$$C(\rho) \geq \frac{2\|C\|_{KF} - (1 - \text{Tr}[\rho_A^2]) - (1 - \text{Tr}[\rho_B^2])}{\sqrt{2M(M-1)}}. \quad (3.65)$$

[Proof] The other orthonormal normalized basis of the local orthonormal observable space can be obtained from A_i and B_i by unitary transformations U and V : $\tilde{A}_i = \sum_l U_{il} A_l$ and $\tilde{B}_j = \sum_m V_{jm}^* B_m$. Select U and V so that $C = U^\dagger \Lambda V$ is the singular value decomposition of C . Then the new observables can be written as $\tilde{A}_i = \sum_l U_{il} A_l$, $\tilde{B}_j = -\sum_m V_{jm}^* B_m$. We have

$$\begin{aligned} \sum_i \Delta_\rho^2(\tilde{A}_i \otimes I + I \otimes \tilde{B}_i) &= \sum_i [\Delta_{\rho_A}^2(\tilde{A}_i) + \Delta_{\rho_B}^2(\tilde{B}_i) + 2(\langle \tilde{A}_i \otimes \tilde{B}_i \rangle - \langle \tilde{A}_i \rangle \langle \tilde{B}_i \rangle)] \\ &= M - \text{Tr}[\rho_A^2] + N - \text{Tr}[\rho_B^2] - 2 \sum_i (UCV^\dagger)_{ii} \\ &= M - \text{Tr}[\rho_A^2] + N - \text{Tr}[\rho_B^2] - 2\|C\|_{KF}. \end{aligned}$$

Substituting above relation to (3.64) one gets (3.65). \square

Bound (3.65) does not depend on the choice of local orthonormal observables. It can be easily applied and realized by direct measurements in experiments. It is in accord with the result in [32] where optimization of entanglement witness based on local uncertainty relation has been taken into account. As an example let us consider the 3×3 bound entangled state [61],

$$\rho = \frac{1}{4}(I_9 - \sum_{i=0}^4 |\xi_i\rangle\langle\xi_i|), \quad (3.66)$$

where I_9 is the 9×9 identity matrix, $|\xi_0\rangle = \frac{1}{\sqrt{2}}|0\rangle(|0\rangle - |1\rangle)$, $|\xi_1\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)|2\rangle$, $|\xi_2\rangle = \frac{1}{\sqrt{2}}|2\rangle(|1\rangle - |2\rangle)$, $|\xi_3\rangle = \frac{1}{\sqrt{2}}(|1\rangle - |2\rangle)|0\rangle$, $|\xi_4\rangle = \frac{1}{3}(|0\rangle + |1\rangle + |2\rangle)(|0\rangle + |1\rangle + |2\rangle)$. We simply choose the local orthonormal observables to be the normalized generators of $SU(3)$. Formula (3.63) gives $C(\rho) \geq 0.050$. Formula (3.64) gives $C(\rho) \geq 0.052$ [50], while formula (3.65) yields a better lower bound $C(\rho) \geq 0.0555$.

If we mix the bound entangled state (3.66) with $|\psi\rangle = \frac{1}{\sqrt{3}} \sum_{i=0}^2 |ii\rangle$, $\rho' = (1-x)\rho + x|\psi\rangle\langle\psi|$, it is easily seen that (3.65) gives a better lower bound of concurrence than formula (3.63) (Fig. 1).

3.1.2 Lower bound of concurrence from “two-qubit” decomposition

In [53] the authors derived an analytical lower bound of concurrence for arbitrary bipartite quantum states by decomposing the joint Hilbert space into many $2 \otimes 2$ dimensional subspaces, which does not involve any optimization procedure and gives an effective evaluation of entanglement together with an operational sufficient condition for the distillability of any bipartite quantum states.




Figure 1: Lower bounds from (3.65) (dashed line) and (3.63) (solid line)

(1) Lower bound of concurrence for bipartite states

The lower bound τ_2 of concurrence for bipartite states has been obtained in [53]. For a bipartite quantum state ρ in $H \otimes H$, the concurrence $C(\rho)$ satisfies

$$\tau_2(\rho) \equiv \frac{d}{2(d-1)} \sum_{m,n=1}^{\frac{d(d-1)}{2}} C_{mn}^2(\rho) \leq C^2(\rho), \quad (3.67)$$

where $C_{mn}(\rho) = \max\{0, \lambda_{mn}^{(1)} - \lambda_{mn}^{(2)} - \lambda_{mn}^{(3)} - \lambda_{mn}^{(4)}\}$ with $\lambda_{mn}^{(1)}, \dots, \lambda_{mn}^{(4)}$ the square roots of the four nonzero eigenvalues, in decreasing order, of the non-Hermitian matrix $\rho \tilde{\rho}_{mn}$ with $\tilde{\rho}_{mn} = (L_m \otimes L_n) \rho^* (L_m \otimes L_n)$, L_m and L_n are the generators of $SO(d)$.

The lower bound τ_2 in Eq.(3.67) in fact characterizes all two-qubit's entanglement in a high dimensional bipartite state. One can directly verify that there are at most $4 \times 4 = 16$ nonzero elements in each matrix $\rho \tilde{\rho}_{mn}$. These elements constitute a 4×4 matrix $\varrho(\sigma_y \otimes \sigma_y) \varrho^*(\sigma_y \otimes \sigma_y)$, where σ_y is the Pauli matrix, the matrix ϱ is a submatrix of the original ρ :

$$\varrho = \begin{pmatrix} \rho_{ik,ik} & \rho_{ik,il} & \rho_{ik,jk} & \rho_{ik,jl} \\ \rho_{il,ik} & \rho_{il,il} & \rho_{il,jk} & \rho_{il,jl} \\ \rho_{jk,ik} & \rho_{jk,il} & \rho_{jk,jk} & \rho_{jk,jl} \\ \rho_{jl,ik} & \rho_{jl,il} & \rho_{jl,jk} & \rho_{jl,jl} \end{pmatrix}, \quad (3.68)$$

$i \neq j$ and $k \neq l$, with subindices i and j associated with the first space, k and l with the second space. The two-qubit submatrix ϱ is not normalized but positive semidefinite. C_{mn} are just the concurrences of these states (3.68).

The bound τ_2 provides a much clearer structure of entanglement, which not only yields an effective separability criterion and an easy evaluation of entanglement, but also helps one to classify mixed-state entanglement.

(2) Lower bound of concurrence for multipartite states

We first consider tripartite case. A general pure state on $H \otimes H \otimes H$ is of the form

$$|\Psi\rangle = \sum_{i,j,k=1}^d a_{ijk} |ijk\rangle, \quad a_{ijk} \in \mathbb{C}, \quad \sum_{i,j,k=1}^d a_{ijk} a_{ijk}^* = 1 \quad (3.69)$$

with

$$C_d^3(|\Psi\rangle) = \sqrt{\frac{d}{6(d-1)}} \times \sqrt{\sum (|a_{ijk}a_{pqm} - a_{ijm}a_{pqk}|^2 + |a_{ijk}a_{pqm} - a_{iqk}a_{pjm}|^2 + |a_{ijk}a_{pqm} - a_{pjk}a_{iqm}|^2)}$$

or equivalently

$$C_d^3(|\Psi\rangle) = \sqrt{\frac{d}{6(d-1)}} (3 - (\text{Tr}[\rho_1^2] + \text{Tr}[\rho_2^2] + \text{Tr}[\rho_3^2])), \quad (3.70)$$

where $\rho_1 = \text{Tr}_{23}[\rho]$, $\rho_2 = \text{Tr}_{13}[\rho]$, $\rho_3 = \text{Tr}_{12}[\rho]$ are the reduced density matrices of $\rho = |\Psi\rangle\langle\Psi|$.

Define $C_{\alpha\beta}^{12|3}(|\Psi\rangle) = |a_{ijk}a_{pqm} - a_{ijm}a_{pqk}|$, $C_{\alpha\beta}^{13|2}(|\Psi\rangle) = |a_{ijk}a_{pqm} - a_{iqk}a_{pjm}|$, $C_{\alpha\beta}^{23|1}(|\Psi\rangle) = |a_{ijk}a_{pqm} - a_{pjk}a_{iqm}|$, where α and β of $C_{\alpha\beta}^{12|3}$ (resp. $C_{\alpha\beta}^{13|2}$ resp. $C_{\alpha\beta}^{23|1}$) stand for the sub-indices of a associated with the subspaces 1, 2 and 3 (resp. 1, 3 and 2 resp. 2, 3 and 1). Let $L^{i_1 i_2 \dots i_N}$ denote the generators of group $SO(d_{i_1} d_{i_2} \dots d_{i_N})$ associated to the subsystems i_1, i_2, \dots, i_N . Then for a tripartite pure state (3.69), one has

$$\begin{aligned} C_d^3(|\Psi\rangle) &= \sqrt{\frac{d}{6(d-1)} \sum_{\alpha}^{\frac{d^2(d^2-1)}{2}} \sum_{\beta}^{\frac{d(d-1)}{2}} [(C_{\alpha\beta}^{12|3}(|\Psi\rangle))^2 + (C_{\alpha\beta}^{13|2}(|\Psi\rangle))^2 + (C_{\alpha\beta}^{23|1}(|\Psi\rangle))^2]} \\ &= \sqrt{\frac{d}{6(d-1)} \sum_{\alpha\beta} [(|\langle\Psi|S_{\alpha\beta}^{12|3}|\Psi^*\rangle|)^2 + (|\langle\Psi|S_{\alpha\beta}^{13|2}|\Psi^*\rangle|)^2 + (|\langle\Psi|S_{\alpha\beta}^{23|1}|\Psi^*\rangle|)^2]}, \end{aligned}$$

where $S_{\alpha\beta}^{12|3} = (L_{\alpha}^{12} \otimes L_{\beta}^3)$, $S_{\alpha\beta}^{13|2} = (L_{\alpha}^{13} \otimes L_{\beta}^2)$ and $S_{\alpha\beta}^{23|1} = (L_{\beta}^1 \otimes L_{\alpha}^{23})$.

Theorem 3.2 For an arbitrary mixed state ρ in $H \otimes H \otimes H$, the concurrence $C(\rho)$ satisfies

$$\tau_3(\rho) \equiv \frac{d}{6(d-1)} \sum_{\alpha}^{\frac{d^2(d^2-1)}{2}} \sum_{\beta}^{\frac{d(d-1)}{2}} [(C_{\alpha\beta}^{12|3}(\rho))^2 + (C_{\alpha\beta}^{13|2}(\rho))^2 + (C_{\alpha\beta}^{23|1}(\rho))^2] \leq C^2(\rho) \quad (3.71)$$

where $\tau_3(\rho)$ is a lower bound of $C(\rho)$,

$$C_{\alpha\beta}^{12|3}(\rho) = \max\{0, \lambda(1)_{\alpha\beta}^{12|3} - \lambda(2)_{\alpha\beta}^{12|3} - \lambda(3)_{\alpha\beta}^{12|3} - \lambda(4)_{\alpha\beta}^{12|3}\}, \quad (3.72)$$

$\lambda(1)_{\alpha\beta}^{12|3}, \lambda(2)_{\alpha\beta}^{12|3}, \lambda(3)_{\alpha\beta}^{12|3}, \lambda(4)_{\alpha\beta}^{12|3}$ are the square roots of the four nonzero eigenvalues, in decreasing order, of the non-Hermitian matrix $\rho \tilde{\rho}_{\alpha\beta}^{12|3}$ with $\tilde{\rho}_{\alpha\beta}^{12|3} = S_{\alpha\beta}^{12|3} \rho^* S_{\alpha\beta}^{12|3}$. $C_{\alpha\beta}^{13|2}(\rho)$ and $C_{\alpha\beta}^{23|1}(\rho)$ are defined in a similar way to $C_{\alpha\beta}^{12|3}(\rho)$.

[Proof] Set $|\xi_i\rangle = \sqrt{p_i}|\psi_i\rangle$, $x_{\alpha\beta}^i = |\langle \xi_i | S_{\alpha\beta}^{12|3} | \xi_i^* \rangle|$, $y_{\alpha\beta}^i = |\langle \xi_i | S_{\alpha\beta}^{13|2} | \xi_i^* \rangle|$ and $z_{\alpha\beta}^i = |\langle \xi_i | S_{\alpha\beta}^{12|3} | \xi_i^* \rangle|$. We have, from Minkowski inequality

$$\begin{aligned} C(\rho) &= \min_i \sqrt{\frac{d}{6(d-1)} \sum_{\alpha\beta} [(x_{\alpha\beta}^i)^2 + (y_{\alpha\beta}^i)^2 + (z_{\alpha\beta}^i)^2]} \\ &\geq \min \sqrt{\frac{d}{6(d-1)} \sum_{\alpha\beta} \left(\sum_i [(x_{\alpha\beta}^i)^2 + (y_{\alpha\beta}^i)^2 + (z_{\alpha\beta}^i)^2]^{\frac{1}{2}} \right)^2}. \end{aligned}$$

Noting that for nonnegative real variables $x_{\alpha}, y_{\alpha}, z_{\alpha}$ and given $X = \sum_{\alpha=1}^N x_{\alpha}$, $Y = \sum_{\alpha=1}^N Y_{\alpha}$ and $Z = \sum_{\alpha=1}^N z_{\alpha}$, by using Lagrange multipliers one obtains that the following inequality holds,

$$\sum_{\alpha=1}^N (x_{\alpha}^2 + y_{\alpha}^2 + z_{\alpha}^2)^{\frac{1}{2}} \geq (X^2 + Y^2 + Z^2)^{\frac{1}{2}}. \quad (3.73)$$

Therefore we have

$$\begin{aligned} C(\rho) &\geq \min \sqrt{\frac{d}{6(d-1)} \sum_{\alpha\beta} [(\sum_i x_{\alpha\beta}^i)^2 + (\sum_i y_{\alpha\beta}^i)^2 + (\sum_i z_{\alpha\beta}^i)^2]} \\ &\geq \sqrt{\frac{d}{6(d-1)} \sum_{\alpha\beta} [(\min_i \sum_i x_{\alpha\beta}^i)^2 + (\min_i \sum_i y_{\alpha\beta}^i)^2 + (\min_i \sum_i z_{\alpha\beta}^i)^2]}. \end{aligned} \quad (3.74)$$

The values of $C_{\alpha\beta}^{12|3}(\rho) \equiv \min_i \sum_i x_{\alpha\beta}^i$, $C_{\alpha\beta}^{13|2}(\rho) \equiv \min_i \sum_i y_{\alpha\beta}^i$ and $C_{\alpha\beta}^{23|1}(\rho) \equiv \min_i \sum_i z_{\alpha\beta}^i$ can be calculated by using the similar procedure in [39]. Here we compute

the value of $C_{\alpha\beta}^{12|3}(\rho)$ in detail. The values of $C_{\alpha\beta}^{13|2}(\rho)$ and $C_{\alpha\beta}^{23|1}(\rho)$ can be obtained analogously.

Let λ_i and $|\chi_i\rangle$ be eigenvalues and eigenvectors of ρ respectively. Any decomposition of ρ can be obtained from a unitary $d^3 \times d^3$ matrix V_{ij} , $|\xi_j\rangle = \sum_{i=1}^{d^3} V_{ij}^*(\sqrt{\lambda_i}|\chi_i\rangle)$. Therefore one has $\langle \xi_i | S_{\alpha\beta}^{12|3} | \xi_j^* \rangle = (VY_{\alpha\beta}V^T)_{ij}$, where the matrix $Y_{\alpha\beta}$ is defined by $(Y_{\alpha\beta})_{ij} = \langle \chi_i | S_{\alpha\beta}^{12|3} | \chi_j^* \rangle$. Namely $C_{\alpha\beta}^{12|3}(\rho) = \min \sum_i |[VY_{\alpha\beta}V^T]_{ii}|$, which has an analytical expression [39], $C_{\alpha\beta}^{12|3}(\rho) = \max\{0, \lambda(1)_{\alpha\beta}^{12|3} - \sum_{j>1} \lambda(j)_{\alpha\beta}^{12|3}\}$, where $\lambda_{\alpha\beta}^{12|3}(k)$ are the square roots of the eigenvalues of the positive Hermitian matrix $Y_{\alpha\beta}Y_{\alpha\beta}^\dagger$, or equivalently the non-Hermitian matrix $\rho\tilde{\rho}_{\alpha\beta}$, in decreasing order. Here as the matrix $S_{\alpha\beta}^{12|3}$ has $d^2 - 4$ rows and $d^2 - 4$ columns that are identically zero, the matrix $\rho\tilde{\rho}_{\alpha\beta}$ has a rank no greater than 4, i.e., $\lambda_{\alpha\beta}^{12|3}(j) = 0$ for $j \geq 5$. From Eq.(3.74) we have Eq.(3.71). \square

Theorem 3.2 can be directly generalized to arbitrary multipartite case.

Theorem 3.3 *For an arbitrary N-partite state $\rho \in H \otimes H \otimes \dots \otimes H$, the concurrence defined in (4.112) satisfies:*

$$\tau_N(\rho) \equiv \frac{d}{2m(d-1)} \sum_p \sum_{\alpha\beta} (C_{\alpha\beta}^p(\rho))^2 \leq C^2(\rho), \quad (3.75)$$

where $\tau_N(\rho)$ is the lower bound of $C(\rho)$, \sum stands for the summation over all possible combinations of the indices of α, β , $C_{\alpha\beta}^p(\rho) = \max\{0, \lambda(1)_{\alpha\beta}^p - \lambda(2)_{\alpha\beta}^p - \lambda(3)_{\alpha\beta}^p - \lambda(4)_{\alpha\beta}^p\}$, $\lambda(i)_{\alpha\beta}^p$, $i = 1, 2, 3, 4$, are the square roots of the four nonzero eigenvalues, in decreasing order, of the non-Hermitian matrix $\rho\tilde{\rho}_{\alpha\beta}^p$ where $\tilde{\rho}_{\alpha\beta}^p = S_{\alpha\beta}^p \rho^* S_{\alpha\beta}^p$.

(3) Lower bound and separability

An N-partite quantum state ρ is fully separable if and only if there exist p_i with $p_i \geq 0$, $\sum_i p_i = 1$ and pure states $\rho_i^j = |\psi_i^j\rangle\langle\psi_i^j|$ such that

$$\rho = \sum_i p_i \rho_i^1 \otimes \rho_i^2 \otimes \dots \otimes \rho_i^N. \quad (3.76)$$

It is easily verified that for a fully separable multipartite state ρ , $\tau_N(\rho) = 0$. Thus $\tau_N(\rho) > 0$ indicates that there must be some kinds of entanglement inside the quantum state, which shows that the lower bound $\tau_N(\rho)$ can be used to recognize entanglement.

As an example we consider a tripartite quantum state [63], $\rho = \frac{1-p}{8}I_8 + p|W\rangle\langle W|$, where I_8 is the 8×8 identity matrix, and $|W\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle)$ is the tripartite W-state. Select an entanglement witness operator to be $\mathcal{W} = \frac{1}{2}I_8 - |GHZ\rangle\langle GHZ|$, where $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ to be the tripartite GHZ-state. From the condition $\text{Tr}[\mathcal{W}\rho] < 0$, the entanglement of ρ is detected for $\frac{3}{5} < p \leq 1$ in [63]. In [79] the authors have obtained the generalized correlation matrix criterion which says if an N-qubit quantum state is fully separable then the inequality $\|\mathcal{T}^N\|_{KF} \leq 1$ must hold, where $\|\mathcal{T}^N\|_{KF} = \max\{\|\mathcal{T}_n^N\|_{KF}\}$, \mathcal{T}_n^N is a kind of matrix unfold of $t_{\alpha_1\alpha_2\cdots\alpha_N}$ defined by $t_{\alpha_1\alpha_2\cdots\alpha_N} = \text{Tr}[\rho\sigma_{\alpha_1}^{(1)}\sigma_{\alpha_2}^{(2)}\cdots\sigma_{\alpha_N}^{(N)}]$ and $\sigma_{\alpha_i}^{(i)}$ stands for the pauli matrix. Now using the generalized correlation matrix criterion the entanglement of ρ is detected for $0.3068 < p \leq 1$. From theorem 3.2, we have that the lower bound $\tau_3(\rho) > 0$ for $0.2727 < p \leq 1$. Therefore the bound (3.110) detects entanglement better than these two criteria in this case. If we replace W with GHZ state in ρ , the criterion in [79] detects the entanglement of ρ for $0.35355 < p \leq 1$, while $\tau_3(\rho)$ detects, again better, the entanglement for $0.2 < p \leq 1$.

Nevertheless for PPT states ρ , we have $\tau_3(\rho) = 0$, which can be seen in the following way. A density matrix ρ is called PPT if the partial transposition of ρ over any subsystem(s) is still positive. Let ρ^{T_i} denote the partial transposition with respect to the i -th subsystem. Assume that there is a PPT state ρ with $\tau(\rho) > 0$. Then at least one term in (3.71), say $C_{\alpha_0\beta_0}^{12|3}(\rho)$, is not zero. Define $\rho_{\alpha_0\beta_0} = L_{\alpha_0}^{12} \otimes L_{\beta_0}^3 \rho (L_{\alpha_0}^{12} \otimes L_{\beta_0}^3)^\dagger$. By using the PPT property of ρ , we have:

$$\rho_{\alpha_0\beta_0}^{T_3} = L_{\alpha_0}^{12} \otimes (L_{\beta_0}^3)^* \rho^{T_3} (L_{\alpha_0}^{12})^\dagger \otimes (L_{\beta_0}^3)^T \geq 0. \quad (3.77)$$

Noting that both $L_{\alpha_0}^{12}$ and $L_{\beta_0}^3$ are projectors to two-dimensional subsystems, $\rho_{\alpha_0\beta_0}$ can be considered as a 4×4 density matrix. While a PPT 4×4 density matrix $\rho_{\alpha_0\beta_0}$ must be a separable state, which contradicts with $C_{\alpha_0\beta_0}^{12|3}(\rho) \neq 0$.

(4) Relation between lower bounds of bi- and tripartite concurrence

τ_3 is basically different from τ_2 as τ_3 characterizes also genuine tripartite entanglement that can not be described by bipartite decompositions. Nevertheless, there are interesting relations between them.

Theorem 3.4 *For any pure tripartite state (3.69), the following inequality holds:*

$$\tau_2(\rho_{12}) + \tau_2(\rho_{13}) + \tau_2(\rho_{23}) \leq 3\tau_3(\rho), \quad (3.78)$$

where τ_2 is the lower bound of bipartite concurrence (3.67), τ_3 is the lower bound of tripartite concurrence (3.71) and $\rho_{12} = \text{Tr}_3[\rho]$, $\rho_{13} = \text{Tr}_2[\rho]$, $\rho_{23} = \text{Tr}_1[\rho]$, $\rho = |\Psi\rangle_{123}\langle\Psi|$.

[Proof] Since $C_{\alpha\beta}^2 \leq (\lambda_{\alpha\beta}(1))^2 \leq \sum_{i=1}^4 (\lambda_{\alpha\beta}(i))^2 = \text{Tr}[\rho\tilde{\rho}_{\alpha\beta}]$ for $\rho = \rho_{12}$, $\rho = \rho_{13}$ and $\rho = \rho_{23}$, we have

$$\begin{aligned} & \tau_2(\rho_{12}) + \tau_2(\rho_{13}) + \tau_2(\rho_{23}) \\ \leq & \frac{d}{2(d-1)} \left(\sum_{\alpha,\beta=1}^{\frac{d(d-1)}{2}} \text{Tr}[\rho_{12}(\tilde{\rho}_{12})_{\alpha\beta}] + \sum_{\alpha,\beta=1}^{\frac{d(d-1)}{2}} \text{Tr}[\rho_{13}(\tilde{\rho}_{13})_{\alpha\beta}] + \sum_{\alpha,\beta=1}^{\frac{d(d-1)}{2}} \text{Tr}[\rho_{23}(\tilde{\rho}_{23})_{\alpha\beta}] \right) \\ = & \frac{d}{2(d-1)} (3 - \text{Tr}[\rho_1^2] - \text{Tr}[\rho_2^2] - \text{Tr}[\rho_3^2]) = 3C^2(\rho) = 3\tau_3(\rho), \end{aligned} \quad (3.79)$$

where we have used the similar analysis in [53, 87] to obtain the equality $\sum_{\alpha,\beta} \text{Tr}[\rho_{12}(\tilde{\rho}_{12})_{\alpha\beta}] = 1 - \text{Tr}[\rho_1^2] - \text{Tr}[\rho_2^2] + \text{Tr}[\rho_3^2]$, $\sum_{\alpha,\beta} \text{Tr}[\rho_{13}(\tilde{\rho}_{13})_{\alpha\beta}] = 1 - \text{Tr}[\rho_1^2] + \text{Tr}[\rho_2^2] - \text{Tr}[\rho_3^2]$, $\sum_{\alpha,\beta} \text{Tr}[\rho_{23}(\tilde{\rho}_{23})_{\alpha\beta}] = 1 + \text{Tr}[\rho_1^2] - \text{Tr}[\rho_2^2] - \text{Tr}[\rho_3^2]$. The last equality is due to that ρ is a pure state. \square

In fact, the bipartite entanglement inside a tripartite state is useful for distilling maximally entangled states. Assume that there are two of the quantities $\{\tau(\rho_{12}), \tau(\rho_{13}), \tau(\rho_{23})\}$ larger than zero, say $\tau(\rho_{12}) > 0$ and $\tau(\rho_{13}) > 0$. According to [53], one can distill two maximal entangled states $|\psi_{12}\rangle$ and $|\psi_{13}\rangle$ which belong to $\mathcal{H}_1 \otimes \mathcal{H}_2$ and $\mathcal{H}_1 \otimes \mathcal{H}_3$ respectively. In terms of the result in [88], one can use them to produce a GHZ state.

3.1.3 Estimation of multipartite entanglement

For a pure N-partite quantum state $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, $\dim\mathcal{H}_i = d_i$, $i = 1, \dots, N$, the concurrence of bipartite decomposition between subsystems $12 \cdots M$ and $M+1 \cdots N$ is defined by

$$C_2(|\psi\rangle) = \sqrt{2(1 - \text{Tr}[\rho_{12 \cdots M}^2])} \quad (3.80)$$

where $\rho_{12 \cdots M}^2 = \text{Tr}_{M+1 \cdots N}[|\psi\rangle\langle\psi|]$ is the reduced density matrix of $\rho = |\psi\rangle\langle\psi|$ by tracing over subsystems $M+1 \cdots N$. On the other hand, the concurrence of $|\psi\rangle$ is defined by (3.59).

For a mixed multipartite quantum state, $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i| \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, the corresponding concurrence of (3.80) and (3.59) are then given by the convex roof:

$$C_2(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C_2(|\psi_i\rangle\langle\psi_i|), \quad (3.81)$$

and (4.112). We now investigate the relation between these two kinds of concurrences.

Lemma 3.1 *For a bipartite density matrix $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B$, one has*

$$1 - \text{Tr}[\rho^2] \leq 1 - \text{Tr}[\rho_A^2] + 1 - \text{Tr}[\rho_B^2], \quad (3.82)$$

where $\rho_{A/B} = \text{Tr}_{B/A}[\rho]$ be the reduced density matrices of ρ .

[Proof] Let $\rho = \sum_{ij} \lambda_{ij} |ij\rangle\langle ij|$ be the spectral decomposition, where $\lambda_{ij} \geq 0$, $\sum_{ij} \lambda_{ij} = 1$. Then $\rho_1 = \sum_{ij} \lambda_{ij} |i\rangle\langle i|$, $\rho_2 = \sum_{ij} \lambda_{ij} |j\rangle\langle j|$. Therefore

$$\begin{aligned} & 1 - \text{Tr}[\rho_A^2] + 1 - \text{Tr}[\rho_B^2] - 1 + \text{Tr}[\rho^2] = 1 - \text{Tr}[\rho_A^2] - \text{Tr}[\rho_B^2] + \text{Tr}[\rho^2] \\ &= \left(\sum_{ij} \lambda_{ij} \right)^2 - \sum_{i,j'} \lambda_{ij} \lambda_{ij'} - \sum_{i',j} \lambda_{ij} \lambda_{i'j} + \sum_{ij} \lambda_{ij}^2 \\ &= \left(\sum_{i=i',j=j'} \lambda_{ij}^2 + \sum_{i=i',j \neq j'} \lambda_{ij} \lambda_{ij'} + \sum_{i \neq i',j=j'} \lambda_{ij} \lambda_{i'j} + \sum_{i \neq i',j \neq j'} \lambda_{ij} \lambda_{i'j'} \right) \\ &= \sum_{i \neq i',j \neq j'} \lambda_{ij} \lambda_{i'j'} \geq 0. \end{aligned}$$

□

This lemma can be also derived in another way [32, 89].

Theorem 3.5 *For a multipartite quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ with $N \geq 3$, the following inequality holds,*

$$C_N(\rho) \geq \max 2^{\frac{3-N}{2}} C_2(\rho), \quad (3.83)$$

where the maximum is taken over all kinds of bipartite concurrence.

[Proof] Without lose of generality, we suppose that the maximal bipartite concurrence is attained between subsystems $12 \cdots M$ and $(M+1) \cdots N$.

For a pure multipartite state $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, $\text{Tr}[\rho_{12 \cdots M}^2] = \text{Tr}[\rho_{(M+1) \cdots N}^2]$. From (3.82) we have

$$\begin{aligned} C_N^2(|\psi\rangle\langle\psi|) &= 2^{2-N} \left((2^N - 2) - \sum_{\alpha} \text{Tr}[\rho_{\alpha}^2] \right) \geq 2^{3-N} \left(N - \sum_{k=1}^N \text{Tr}[\rho_k^2] \right) \\ &\geq 2^{3-N} (1 - \text{Tr}[\rho_{12 \cdots M}^2] + 1 - \text{Tr}[\rho_{(M+1) \cdots N}^2]) \\ &= 2^{3-N} * 2(1 - \text{Tr}[\rho_{12 \cdots M}^2]) = 2^{3-N} C_2^2(|\psi\rangle\langle\psi|), \end{aligned}$$

i.e. $C_N(|\psi\rangle\langle\psi|) \geq 2^{\frac{3-N}{2}} C_2(|\psi\rangle\langle\psi|)$.

Let $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ attain the minimal decomposition of the multipartite concurrence. One has

$$\begin{aligned} C_N(\rho) &= \sum_i p_i C_N(|\psi_i\rangle\langle\psi_i|) \geq 2^{\frac{3-N}{2}} \sum_i p_i C_2(|\psi_i\rangle\langle\psi_i|) \\ &\geq 2^{\frac{3-N}{2}} \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C_2(|\psi_i\rangle\langle\psi_i|) = 2^{\frac{3-N}{2}} C_2(\rho). \end{aligned}$$

□

Corollary 3.5 *For a tripartite quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$, the following inequality holds:*

$$C_3(\rho) \geq \max C_2(\rho) \quad (3.84)$$

where the maximum is taken over all kinds of bipartite concurrence.

In [50, 32], from the separability criteria related to local uncertainty relation, covariance matrix and correlation matrix, the following lower bounds for bipartite concurrence are obtained:

$$C_2(\rho) \geq \frac{2\|C(\rho)\| - (1 - \text{Tr}[\rho_A^2]) - (1 - \text{Tr}[\rho_B^2])}{\sqrt{2d_A(d_A - 1)}} \quad (3.85)$$

and

$$C_2(\rho) \geq \sqrt{\frac{8}{d_A^3 d_B^2 (d_A - 1)} \left(\|T(\rho)\| - \frac{\sqrt{d_A d_B (d_A - 1)(d_B - 1)}}{2} \right)}, \quad (3.86)$$

where the entries of the matrix C , $C_{ij} = \langle \lambda_i^A \otimes \lambda_j^B \rangle - \langle \lambda_i^A \otimes I_{d_B} \rangle \langle I_{d_A} \otimes \lambda_j^B \rangle$, $T_{ij} = \frac{d_A d_B}{2} \langle \lambda_i^A \otimes \lambda_j^B \rangle$, $\lambda_k^{A/B}$ stands for the normalized generator of $SU(d_A/d_B)$, i.e. $\text{Tr}[\lambda_k^{A/B} \lambda_l^{A/B}] = \delta_{kl}$ and $\langle X \rangle = \text{Tr}[\rho X]$. It is shown that the lower bounds (3.85) and (3.86) are independent of (3.63).

Now we consider a multipartite quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ as a bipartite state belonging to $\mathcal{H}^A \otimes \mathcal{H}^B$ with the dimensions of the subsystems A and B being $d_A = d_{s_1} d_{s_2} \cdots d_{s_m}$ and $d_B = d_{s_{m+1}} d_{s_{m+2}} \cdots d_{s_N}$ respectively. By using the corollary, (3.63), (3.85) and (3.86) one has the following lower bound:

Theorem 3.6 *For any N -partite quantum state ρ ,*

$$C_N(\rho) \geq 2^{\frac{3-N}{2}} \max\{B_1, B_2, B_3\}, \quad (3.87)$$

where

$$\begin{aligned}
B_1 &= \max_{\{i\}} \sqrt{\frac{2}{M_i(M_i - 1)}} [\max(\|\mathcal{T}_A(\rho^i)\|, \|R(\rho^i)\|) - 1], \\
B_2 &= \max_{\{i\}} \frac{2\|C(\rho^i)\| - (1 - \text{Tr}[(\rho_A^i)^2]) - (1 - \text{Tr}[(\rho_B^i)^2])}{\sqrt{2M_i(M_i - 1)}}, \\
B_3 &= \max_{\{i\}} \sqrt{\frac{8}{M_i^3 N_i^2 (M_i - 1)}} (\|T(\rho^i)\| - \frac{\sqrt{M_i N_i (M_i - 1)(N_i - 1)}}{2}),
\end{aligned}$$

ρ^i s are all possible bipartite decompositions of ρ , and

$$\begin{aligned}
M_i &= \min \{d_{s_1} d_{s_2} \cdots d_{s_m}, d_{s_{m+1}} d_{s_{m+2}} \cdots d_{s_N}\}, \\
N_i &= \max \{d_{s_1} d_{s_2} \cdots d_{s_m}, d_{s_{m+1}} d_{s_{m+2}} \cdots d_{s_N}\}.
\end{aligned}$$

In [32, 84, 90], it is shown that the upper and lower bound of multipartite concurrence satisfy

$$\sqrt{(4 - 2^{3-N})\text{Tr}[\rho^2] - 2^{2-N} \sum_{\alpha} \text{Tr}[\rho_{\alpha}^2]} \leq C_N(\rho) \leq \sqrt{2^{2-N}[(2^N - 2) - \sum_{\alpha} \text{Tr}[\rho_{\alpha}^2]]} \quad (3.88)$$

In fact one can obtain a more effective upper bound for multi-partite concurrence. Let $\rho = \sum_i \lambda_i |\psi_i\rangle\langle\psi_i| \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, where $|\psi_i\rangle$ s are the orthogonal pure states and $\sum_i \lambda_i = 1$. We have

$$C_N(\rho) = \min_{\{p_i, |\varphi_i\rangle\}} \sum_i p_i C_N(|\varphi_i\rangle\langle\varphi_i|) \leq \sum_i \lambda_i C_N(|\psi_i\rangle\langle\psi_i|). \quad (3.89)$$

The right side of (3.89) gives a new upper bound of $C_N(\rho)$. Since

$$\begin{aligned}
\sum_i \lambda_i C_N(|\psi_i\rangle\langle\psi_i|) &= 2^{1-\frac{N}{2}} \sum_i \lambda_i \sqrt{(2^N - 2) - \sum_{\alpha} \text{Tr}[(\rho_{\alpha}^i)^2]} \\
&\leq 2^{1-\frac{N}{2}} \sqrt{(2^N - 2) - \sum_{\alpha} \text{Tr}[\sum_i \lambda_i (\rho_{\alpha}^i)^2]} \\
&\leq 2^{1-\frac{N}{2}} \sqrt{(2^N - 2) - \sum_{\alpha} \text{Tr}[(\rho_{\alpha})^2]},
\end{aligned}$$

the upper bound obtained in (3.89) is better than that in (3.88).

3.1.4 Bounds of concurrence and tangle

In [54], a lower bound for tangle defined in (3.58) has been derived:

$$\tau(\rho) \geq \frac{8}{MN(M+N)} (\|T(\rho)\|_{HS}^2 - \frac{MN(M-1)(N-1)}{4}), \quad (3.90)$$

where $\|X\|_{HS} = \sqrt{\text{Tr}[XX^\dagger]}$ denotes the Frobenius or Hilbert-Schmidt norm. Experimentally measurable lower and upper bounds for concurrence have been also given by Mintert and Zhang et.al. in [84, 32]:

$$\sqrt{2(\text{Tr}[\rho^2] - \text{Tr}[\rho_A^2])} \leq C(\rho) \leq \sqrt{2(1 - \text{Tr}[\rho_A^2])}. \quad (3.91)$$

Since the convexity of $C^2(\rho)$, we have that $\tau(\rho) \geq C^2(\rho)$ always holds. For two qubits quantum systems, tangle τ is always equal to the square of concurrence C^2 [44, 91], as a decomposition $\{p_i, |\psi_i\rangle\}$ achieving the minimum in Eq. (3.54) has the property that $C(|\psi_i\rangle) = C(|\psi_j\rangle) \forall i, j$. For higher dimensional systems we do not have similar relations. Thus it is meaningful to derive valid upper bound for tangle and lower bound for concurrence.

Theorem 3.7 *For any quantum state $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B$, we have*

$$\tau(\rho) \leq \min\{1 - \text{Tr}[\rho_A^2], 1 - \text{Tr}[\rho_B^2]\}, \quad (3.92)$$

$$C(\rho) \geq \sqrt{\frac{8}{MN(M+N)} (\|T(\rho)\|_{HS} - \frac{\sqrt{MN(M-1)(N-1)}}{2})}, \quad (3.93)$$

where ρ_A is the reduced matrix of ρ , and $T(\rho)$ is the correlation matrix of ρ defined in (3.86).

[Proof] We assume $1 - \text{Tr}[\rho_A^2] \leq 1 - \text{Tr}[\rho_B^2]$ for convenience. By the definition of τ , we have that for a pure state $|\psi\rangle$, $\tau(|\psi\rangle) = 2(1 - \text{Tr}[(\rho_A^{|\psi\rangle})^2])$. Let $\rho = \sum_i p_i \rho_i$ be the optimal decomposition such that $\tau(\rho) = \sum_i p_i \tau(\rho_i)$. We get

$$\tau(\rho) = \sum_i p_i \tau(\rho_i) = \sum_i p_i 2[1 - \text{Tr}[(\rho_A^{|\psi_i\rangle})^2]] = 2[1 - \text{Tr}[\sum_i p_i (\rho_A^{|\psi_i\rangle})^2]] \leq 2[1 - \text{Tr}[\rho_A^2]]. \quad (3.94)$$

Note that for pure state $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ [54],

$$C(|\psi\rangle) = \sqrt{\frac{8}{MN(M+N)} (\|T(|\psi\rangle)\|^2 - \frac{MN(M-1)(N-1)}{4})}. \quad (3.95)$$

Using the inequality $\sqrt{a-b} \geq \sqrt{a} - \sqrt{b}$ for any $a \geq b$, we get

$$C(|\psi\rangle) \geq \sqrt{\frac{8}{MN(M+N)}} (\|T(|\psi\rangle)\|_{HS} - \frac{\sqrt{MN(M-1)(N-1)}}{2}). \quad (3.96)$$

Now let $\rho = \sum_i p_i \rho_i$ be the optimal decomposition such that $C(\rho) = \sum_i p_i C(\rho_i)$. We get

$$\begin{aligned} C(\rho) &= \sum_i p_i C(\rho_i) \geq \sum_i p_i \sqrt{\frac{8}{MN(M+N)}} (\|T(\rho_i)\|_{HS} - \frac{\sqrt{MN(M-1)(N-1)}}{2}) \\ &= \sqrt{\frac{8}{MN(M+N)}} (\sum_i p_i \|T(\rho_i)\|_{HS} - \frac{\sqrt{MN(M-1)(N-1)}}{2}) \\ &\geq \sqrt{\frac{8}{MN(M+N)}} (\|T(\rho)\|_{HS} - \frac{\sqrt{MN(M-1)(N-1)}}{2}) \end{aligned}$$

which ends the proof. \square

The upper bound (3.92), together with the lower bound (3.93), (3.85), (3.86), (3.90) and (3.91), can allow for estimations of entanglement for arbitrary quantum states. Moreover, since the upper bound is exactly the value of tangle for pure states, the upper bound can be a good estimation when the state is very weakly mixed.

3.2 Concurrence and tangle of two entangled states are strictly larger than that of one

In this subsection we show that although bound entangled states can not be distilled, the concurrence and tangle of two entangled states will be always strictly larger than that of one, even the two entangled states are both bound entangled.

Let $\rho = \sum_{ijkl} \rho_{ij,kl} |ij\rangle\langle kl| \in \mathcal{H}_A \otimes \mathcal{H}_B$ and $\sigma = \sum_{i'j'k'l'} \sigma_{i'j',k'l'} |i'j'\rangle\langle k'l'| \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$ be two quantum states shared by subsystems AA' and BB' . We use $\rho \oplus \sigma = \sum_{ijkl,i'j'k'l'} |ii'\rangle_{AA'} \langle kk'| \otimes |jj'\rangle_{BB'} \langle ll'|$ to denote the state of the whole system.

Lemma 3.2 *For pure states $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ and $|\varphi\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$, the inequalities*

$$C(|\psi\rangle \oplus |\varphi\rangle) \geq \max\{C(|\psi\rangle), C(|\varphi\rangle)\} \quad (3.97)$$

and

$$\tau(|\psi\rangle \oplus |\varphi\rangle) \geq \max\{\tau(|\psi\rangle), \tau(|\varphi\rangle)\} \quad (3.98)$$

always hold, and " = " in the two inequalities hold if and only if at least one of $\{|\psi\rangle, |\varphi\rangle\}$ is separable.

[Proof] Without loss of generality we assume $C(|\psi\rangle) \geq C(|\varphi\rangle)$. First note that

$$\rho_{AA'}^{|\psi\rangle \oplus |\varphi\rangle} = \rho_A^{|\psi\rangle} \otimes \rho_{A'}^{|\varphi\rangle}. \quad (3.99)$$

Let $\rho_A^{|\psi\rangle} = \sum_i \lambda_i |i\rangle\langle i|$ and $\rho_{A'}^{|\varphi\rangle} = \sum_j \pi_j |j\rangle\langle j|$ be the spectral decomposition of $\rho_A^{|\psi\rangle}$ and $\rho_{A'}^{|\varphi\rangle}$, with $\sum_i \lambda_i = 1$ and $\sum_j \pi_j = 1$ respectively. By using (3.99) one obtains that

$$\text{Tr}[(\rho_{AA'}^{|\psi\rangle \oplus |\varphi\rangle})^2] = \sum \lambda_i \pi_j \lambda_{i'} \pi_{j'} |ij\rangle\langle ij| i'j'\rangle\langle i'j'| = \sum \lambda_i^2 \pi_j^2 \quad (3.100)$$

while

$$\text{Tr}[(\rho_A^{|\psi\rangle})^2] = \sum_i \lambda_i^2. \quad (3.101)$$

Now using the definition of concurrence and the normalization conditions of λ_i and π_j one immediately gets

$$C(|\psi\rangle \oplus |\varphi\rangle) = \sqrt{2(1 - \text{Tr}[(\rho_{AA'}^{|\psi\rangle \oplus |\varphi\rangle})^2])} \geq \sqrt{2(1 - \text{Tr}[(\rho_A^{|\psi\rangle})^2])} = C(|\psi\rangle). \quad (3.102)$$

If one of $\{|\psi\rangle, |\varphi\rangle\}$ is separable, say $|\varphi\rangle$, then the rank of $\rho_{A'}^{|\varphi\rangle}$ must be one, which means that there is only one item in the spectral decomposition in $\rho_{A'}^{|\varphi\rangle}$. Using the normalization condition of π_j we obtain $\text{Tr}[(\rho_{AA'}^{|\psi\rangle \oplus |\varphi\rangle})^2] = \text{Tr}[(\rho_A^{|\psi\rangle})^2]$. Then the inequality (3.102) becomes an equality.

On the other hand, if both $|\psi\rangle$ and $|\varphi\rangle$ are entangled (not separable), there must be at least two items in the decomposition of their reduced density matrices $\rho_A^{|\psi\rangle}$ and $\rho_{A'}^{|\varphi\rangle}$, which means that $\text{Tr}[(\rho_{AA'}^{|\psi\rangle \oplus |\varphi\rangle})^2]$ is strictly larger than $\text{Tr}[(\rho_A^{|\psi\rangle})^2]$.

The inequality (3.98) also holds because that for pure quantum state ρ , $\tau(\rho) = C^2(\rho)$. \square

From the lemma, we have, for mixed states,

Theorem 3.8 For any quantum states $\rho \in \mathcal{H}_A \otimes \mathcal{H}_B$ and $\sigma \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$, the inequalities

$$C(\rho \oplus \sigma) \geq \max\{C(\rho), C(\sigma)\} \quad (3.103)$$

and

$$\tau(\rho \oplus \sigma) \geq \max\{\tau(\rho), \tau(\sigma)\} \quad (3.104)$$

always hold, and the " = " in the two inequalities hold if and only if at least one of $\{\rho, \sigma\}$ is separable, i.e. if both ρ and σ are entangled (even bound entangled), $C(\rho \oplus \sigma) > \max\{C(\rho), C(\sigma)\}$ and $\tau(\rho \oplus \sigma) > \max\{\tau(\rho), \tau(\sigma)\}$ always hold.

[Proof] We still assume $C(\rho) \geq C(\sigma)$ for convenience. Let $\rho = \sum_i p_i \rho_i$ and $\sigma = \sum_j q_j \sigma_j$ be the optimal decomposition such that $C(\rho \oplus \sigma) = \sum_i p_i q_j C(\rho_i \oplus \sigma_j)$. By using the inequality obtained in lemma 3.2 we have

$$C(\rho \oplus \sigma) = \sum_i p_i q_j C(\rho_i \oplus \sigma_j) \geq \sum_i p_i q_j C(\rho_i) = \sum_i p_i C(\rho_i) \geq C(\rho). \quad (3.105)$$

Case 1: Now let one of $\{\rho, \sigma\}$ be separable, say σ , with ensemble representation $\sigma = \sum_j q_j \sigma_j$, where $\sum_j q_j = 1$ and σ_j is the density matrix of separable pure state. Suppose $\rho = \sum_i p_i \rho_i$ be the optimal decomposition of ρ such that $C(\rho) = \sum_i p_i C(\rho_i)$. Using lemma 3.2 we have

$$C(\rho \oplus \sigma) \leq \sum_i p_i q_j C(\rho_i \oplus \sigma_j) = \sum_i p_i q_j C(\rho_i) = \sum_i p_i C(\rho_i) = C(\rho). \quad (3.106)$$

The inequalities (3.105) and (3.106) show that if σ is separable, then $C(\rho \oplus \sigma) = C(\rho)$.

Case 2: If both ρ and σ are inseparable, i.e. there is at least one pure state in the ensemble decomposition of ρ (and σ respectively), using lemma 3.2 we have

$$C(\rho \oplus \sigma) = \sum_i p_i q_j C(\rho_i \oplus \sigma_j) > \sum_i p_i q_j C(\rho_i) = \sum_i p_i C(\rho_i) \geq C(\rho). \quad (3.107)$$

The inequality for tangle τ can be proved in a similar way. \square

Remark : In [92] it is shown that any entangled state ρ can enhance the teleportation power of another state σ . This holds even if the state ρ is bound entangled. But if ρ is bound entangled, the corresponding σ must be free entangled (distillable). By theorem 3.8, we can see that even two entangled quantum states ρ and σ are bound entangled, their concurrence and tangle are strictly larger than that of one state.

3.3 Subadditivity of concurrence and tangle

We now give a proof of the subadditivity of concurrence and tangle, which illustrates that concurrence and tangle may be proper entanglement measurements.

Theorem 3.9 Let ρ and σ be quantum states in $\mathcal{H}_A \otimes \mathcal{H}_B$, we have

$$C(\rho \otimes \sigma) \leq C(\rho) + C(\sigma) \quad \text{and} \quad \tau(\rho \otimes \sigma) \leq \tau(\rho) + \tau(\sigma). \quad (3.108)$$

Proof: We first prove that the theorem holds for pure states, i.e. for $|\psi\rangle$ and $|\phi\rangle$ in $\mathcal{H}_A \otimes \mathcal{H}_B$,

$$C(|\psi\rangle \otimes |\phi\rangle) \leq C(|\psi\rangle) + C(|\phi\rangle) \quad \text{and} \quad \tau(|\psi\rangle \otimes |\phi\rangle) \leq \tau(|\psi\rangle) + \tau(|\phi\rangle). \quad (3.109)$$

Assume that $\rho_A^{|\psi\rangle} = \sum_i \lambda_i |i\rangle\langle i|$ and $\rho_A^{|\phi\rangle} = \sum_j \pi_j |j\rangle\langle j|$ be the spectral decomposition of the reduced matrices $\rho_A^{|\psi\rangle}$ and $\rho_A^{|\phi\rangle}$. One has

$$\begin{aligned} & \frac{1}{2} [C(|\psi\rangle) + C(|\phi\rangle)]^2 \geq 1 - \text{Tr}[(\rho_A^{|\psi\rangle})^2] + 1 - \text{Tr}[(\rho_A^{|\phi\rangle})^2] \\ & = 1 - \sum_i \lambda_i^2 + 1 - \sum_j \pi_j^2 \geq 1 - \sum_{ij} \lambda_i^2 \pi_j^2 = \frac{1}{2} C^2(|\psi\rangle \otimes |\phi\rangle). \end{aligned} \quad (3.110)$$

Now we prove that (3.108) holds for any mixed quantum states ρ and σ . Let $\rho = \sum_i p_i \rho_i$ and $\sigma = \sum_j q_j \sigma_j$ be the optimal decomposition such that $C(\rho) = \sum_i p_i C(\rho_i)$ and $C(\sigma) = \sum_j q_j C(\sigma_j)$. We have

$$C(\rho) + C(\sigma) = \sum_{ij} p_i q_j [C(\rho_i) + C(\sigma_j)] \geq \sum_{ij} p_i q_j C(\rho_i \otimes \sigma_j) \geq C(\rho \otimes \sigma). \quad (3.111)$$

The inequality for τ can be derived in a similar way. □

4 Fidelity of teleportation and distillation of entanglement

Quantum teleportation is an important subject in quantum information processing. In terms of a classical communication channel and a quantum resource (a nonlocal entangled state like an EPR-pair of particles), the teleportation protocol gives ways to transmit an unknown quantum state from a sender traditionally named ‘‘Alice’’ to a receiver ‘‘Bob’’ who are spatially separated. These teleportation processes can be viewed as quantum channels. The nature of a quantum channel is determined by the particular protocol and the state used as a teleportation resource. The standard teleportation protocol T_0 proposed by Bennett et.al in 1993 uses *Bell* measurements and *Pauli* rotations. When the maximally entangled pure state $|\phi\rangle = \frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} |ii\rangle$ is used as the quantum resource, it provides an ideal noiseless quantum channel

$\Lambda_{T_0}^{(|\phi\rangle\langle\phi|)}(\rho) = \rho$. However in realistic situation, instead of the pure maximally entangled states, Alice and Bob usually share a mixed entangled state due to the decoherence. Teleportation using mixed state as an entangled resource is, in general, equivalent to having a noisy quantum channel. An explicit expression for the output state of the quantum channel associated with the standard teleportation protocol T_0 with an arbitrary mixed state resource has been obtained [93, 94].

It turns out that by local quantum operations (including collective actions over all members of pairs in each lab) and classical communication (LOCC) between Alice and Bob, it is possible to obtain a number of pairs in nearly maximally entangled state $|\psi_+\rangle$ from many pairs of non-maximally entangled states. Such a procedure proposed in [58, 59, 60, 61, 62] is called distillation. In [58] the authors give operational protocol to distill an entangled two-qubit state whose single fraction F , defined by $F(\rho) = \langle\psi_+|\rho|\psi_+\rangle$, is larger than $\frac{1}{2}$. The protocol is then generalized in [62] to distill any d -dimensional bipartite entangled quantum states with $F(\rho) > \frac{1}{d}$. It is shown that a quantum state ρ violating the reduction criterion can always be distilled. For such states if their single fraction of entanglement $F(\rho) = \langle\psi_+|\rho|\psi_+\rangle$ is greater than $\frac{1}{d}$, one can distill these states directly by using the generalized distillation protocol, otherwise a proper filtering operation has to be used at first to transform ρ to another state ρ' so that $F(\rho') > \frac{1}{d}$.

4.1 Fidelity of quantum teleportation

Let \mathcal{H} be a d -dimensional complex vector space with computational basis $|i\rangle$, $i = 1, \dots, d$. The fully entangled fraction (FEF) of a density matrix $\rho \in \mathcal{H} \otimes \mathcal{H}$ is defined by

$$\mathcal{F}(\rho) = \max_U \langle\psi_+|(I \otimes U^\dagger)\rho(I \otimes U)|\psi_+\rangle \quad (4.112)$$

under all unitary transformations U , where $|\psi_+\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle$ is the maximally entangled state and I is the corresponding identity matrix.

In [95], the authors give a optimal teleportation protocol by using a mixed entangled quantum state. The optimal teleportation fidelity is given by

$$f_{\max}(\rho) = \frac{d\mathcal{F}(\rho)}{d+1} + \frac{1}{d+1}, \quad (4.113)$$

which solely depends the FEF of the entangled resource state ρ .

In fact the fully entangled fraction is tightly related to many quantum information processing such as dense coding [7], teleportation [5], entanglement swapping [9], and quantum cryptography (Bell inequalities) [8]. As the optimal fidelity of teleportation is given by FEF [6], experimentally measurement of FEF can be also used to determine the entanglement of the non-local source used in teleportation. Thus an analytic formula for FEF is of great importance. In [96] an elegant formula of FEF for two-qubit system is derived analytically by using the method of Lagrange multipliers. For high dimensional quantum states the analytical computation of FEF remains formidable and less results have been known. In the following we give an estimation on the values of FEF by giving some upper bounds of FEF .

Let $\lambda_i, i = 1, \dots, d^2 - 1$, be the generators of the $SU(d)$ algebra. A bipartite state $\rho \in \mathcal{H} \otimes \mathcal{H}$ can be expressed as

$$\rho = \frac{1}{d^2} I \otimes I + \frac{1}{d} \sum_{i=1}^{d^2-1} r_i(\rho) \lambda_i \otimes I + \frac{1}{d} \sum_{j=1}^{d^2-1} s_j(\rho) I \otimes \lambda_j + \sum_{i,j=1}^{d^2-1} m_{ij}(\rho) \lambda_i \otimes \lambda_j, \quad (4.114)$$

where $r_i(\rho) = \frac{1}{2} \text{Tr}[\rho \lambda_i(1) \otimes I]$, $s_j(\rho) = \frac{1}{2} \text{Tr}[\rho I \otimes \lambda_j(2)]$ and $m_{ij}(\rho) = \frac{1}{4} \text{Tr}[\rho \lambda_i(1) \otimes \lambda_j(2)]$. Let $M(\rho)$ denote the correlation matrix with entries $m_{ij}(\rho)$.

Theorem 4.1 *For any $\rho \in \mathcal{H} \otimes \mathcal{H}$, the fully entangled fraction $\mathcal{F}(\rho)$ satisfies*

$$\mathcal{F}(\rho) \leq \frac{1}{d^2} + 4 \|M^T(\rho)M(P_+)\|_{KF}, \quad (4.115)$$

where M^T stands for the transpose of M and $\|M\|_{KF} = \text{Tr}[\sqrt{MM^\dagger}]$ is the Ky Fan norm of M .

[Proof] First, we note that

$$P_+ = \frac{1}{d^2} I \otimes I + \sum_{i,j=1}^{d^2-1} m_{ij}(P_+) \lambda_i \otimes \lambda_j,$$

where $m_{ij}(P_+) = \frac{1}{4} \text{Tr}[P_+ \lambda_i \otimes \lambda_j]$. By definition (4.112), one obtains

$$\begin{aligned} \mathcal{F}(\rho) &= \max_U \langle \psi_+ | (I \otimes U^\dagger) \rho (I \otimes U) | \psi_+ \rangle = \max_U \text{Tr}[\rho (I \otimes U) P_+ (I \otimes U^\dagger)] \\ &= \max_U \left\{ \frac{1}{d^2} \text{Tr}[\rho] + \sum_{i,j=1}^{d^2-1} m_{ij}(P_+) \text{Tr}[\rho \lambda_i \otimes U \lambda_j U^\dagger] \right\}. \end{aligned}$$

Since $U \lambda_i U^\dagger$ is a traceless Hermitian operator, it can be expanded according to the $SU(d)$ generators,

$$U \lambda_i U^\dagger = \sum_{j=1}^{d^2-1} \frac{1}{2} \text{Tr}[U \lambda_i U^\dagger \lambda_j] \lambda_j \equiv \sum_{j=1}^{d^2-1} O_{ij} \lambda_j. \quad (4.116)$$

Entries O_{ij} defines a real $(d^2 - 1) \times (d^2 - 1)$ matrix O . From the completeness relation of $SU(d)$ generators

$$\sum_{j=1}^{d^2-1} (\lambda_j)_{ki} (\lambda_j)_{mn} = 2\delta_{im}\delta_{kn} - \frac{2}{d}\delta_{ki}\delta_{mn}, \quad (4.117)$$

one can show that O is an orthonormal matrix. Using (4.116) we have

$$\begin{aligned} \mathcal{F}(\rho) &\leq \frac{1}{d^2} + \max_O \sum_{i,j,k} m_{ij}(P_+) O_{jk} \text{Tr}[\rho \lambda_i \otimes \lambda_k] \\ &= \frac{1}{d^2} + 4 \max_O \sum_{i,j,k} m_{ij}(P_+) O_{jk} m_{ik}(\rho) = \frac{1}{d^2} + 4 \max_O \text{Tr}[M(\rho)^T M(P_+) O] \\ &= \frac{1}{d^2} + 4 \|M(\rho)^T M(P_+)\|_{KF}. \end{aligned}$$

□

For the case $d = 2$, we can get an exact result from (4.115):

Corollary 4.1 *For two qubits system, we have*

$$\mathcal{F}(\rho) = \frac{1}{4} + 4 \|M(\rho)^T M(P_+)\|_{KF}, \quad (4.118)$$

i.e. the upper bound derived in Theorem 4.1 is exactly the FEF.

[Proof] We have shown in (4.116) that given an arbitrary unitary U , one can always obtain an orthonormal matrix O . Now we show that in two-qubit case, for any 3×3 orthonormal matrix O there always exists 2×2 unitary matrix U such that (4.116) holds.

For any vector $\mathbf{t} = \{t_1, t_2, t_3\}$ with unit norm, define an operator $X \equiv \sum_{i=1}^3 t_i \sigma_i$, where σ_i s are Pauli matrices. Given an orthonormal matrix O one obtains a new operator $X' \equiv \sum_{i=1}^3 t'_i \sigma_i = \sum_{i,j=1}^3 O_{ij} t_j \sigma_i$.

X and X' are both hermitian traceless matrices. Their eigenvalues are given by the norms of the vectors \mathbf{t} and $\mathbf{t}' = \{t'_1, t'_2, t'_3\}$ respectively. As the norms are invariant under orthonormal transformations O , they have the same eigenvalues: $\pm \sqrt{t_1^2 + t_2^2 + t_3^2}$. Thus there must be a unitary matrix U such that $X' = UXU^\dagger$. Hence the inequality in the proof of Theorem 4.1 becomes an equality. The upper bound (4.115) then becomes exact at this situation, which is in accord with the result in [96]. □

Remark : The upper bound of FEF (4.115) and the FEF (4.118) depend on the correlation matrices $M(\rho)$ and $M(P_+)$. They can be calculated directly according to a given set of $SU(d)$ generators λ_i , $i = 1, \dots, d^2 - 1$. As an example, for $d = 3$, if we

$$\text{choose } \lambda_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_2 = \begin{pmatrix} \frac{1}{\sqrt{3}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{3}} & 0 \\ 0 & 0 & -\frac{2}{\sqrt{3}} \end{pmatrix}, \lambda_3 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \lambda_5 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \lambda_6 = \begin{pmatrix} 0 & i & 0 \\ -i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_7 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \text{ and} \\ \lambda_8 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & i \\ 0 & -i & 0 \end{pmatrix}, \text{ then we have}$$

$$M(P_+) = \text{Diag}\left\{\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, -\frac{1}{6}, -\frac{1}{6}, -\frac{1}{6}\right\}. \quad (4.119)$$

Nevertheless the *FEF* and its upper bound do not depend on the choice of the $SU(d)$ generators.

The usefulness of the bound depends on detailed states. In the following we give two new upper bounds which is different from theorem 4.1. These bounds work for different states.

Let h and g be $n \times n$ matrices such that $h|j\rangle = |(j+1) \bmod n\rangle$, $g|j\rangle = \omega^j|j\rangle$, with $\omega = \exp\{\frac{-2i\pi}{n}\}$. We can introduce n^2 linear-independent $n \times n$ -matrices $U_{st} = h^t g^s$, which satisfy

$$U_{st} U_{s't'} = \omega^{st'-ts'} U_{s't'} U_{st}, \quad \text{Tr}[U_{st}] = n \delta_{s0} \delta_{t0}. \quad (4.120)$$

One can also check that $\{U_{st}\}$ satisfy the condition of *bases of the unitary operators* in the sense of [97], i.e.

$$\text{Tr}[U_{st} U_{s't'}^+] = n \delta_{tt'} \delta_{ss'}, \quad U_{st} U_{st}^+ = I_{n \times n}, \quad (4.121)$$

where $I_{n \times n}$ is the $n \times n$ identity matrix. $\{U_{st}\}$ form a complete basis of $n \times n$ -matrices, namely, for any $n \times n$ matrix W , W can be expressed as

$$W = \frac{1}{n} \sum_{s,t} \text{Tr}[U_{st}^+ W] U_{st}. \quad (4.122)$$

From $\{U_{st}\}$, we can introduce the generalized Bell-states,

$$|\Phi_{st}\rangle = (I \otimes U_{st}^*) |\psi_+\rangle = \frac{1}{\sqrt{d}} \sum_{i,j} (U_{st})_{ij}^* |ij\rangle, \quad \text{and} \quad |\Phi_{00}\rangle = |\psi_+\rangle, \quad (4.123)$$

$|\Phi_{st}\rangle$ are all maximally entangled states and form a complete orthogonal normalized basis of $\mathcal{H}_d \otimes \mathcal{H}_d$.

Theorem 4.2 For any quantum state $\rho \in \mathcal{H}_d \otimes \mathcal{H}_d$, the fully entangled fraction defined in (4.112) fulfills the following inequality:

$$\mathcal{F}(\rho) \leq \max_j \{\lambda_j\}, \quad (4.124)$$

where λ_j s are the eigenvalues of the real part of matrix $M = \begin{pmatrix} T & iT \\ -iT & T \end{pmatrix}$, T is a $d^2 \times d^2$ matrix with entries $T_{n,m} = \langle \Phi_n | \rho | \Phi_m \rangle$ and Φ_j are the maximally entangled basis states defined in (4.123).

[Proof] From (4.122), any $d \times d$ unitary matrix U can be represented by $U = \sum_{k=1}^{d^2} z_k U_k$, where $z_k = \frac{1}{d} \text{Tr}[U_k^\dagger U]$. Define

$$x_l = \begin{cases} \text{Re}[z_l], & 1 \leq l \leq d^2; \\ \text{Im}[z_l], & d^2 < l \leq 2d^2 \end{cases} \quad \text{and} \quad U'_l = \begin{cases} U_l, & 1 \leq l \leq d^2; \\ i * U_l, & d^2 < l \leq 2d^2. \end{cases} \quad (4.125)$$

Then the unitary matrix U can be rewritten as $U = \sum_{k=1}^{2d^2} z_k U'_k$. The necessary condition for the unitary property of U implies that $\sum_k x_k^2 = 1$. Thus we have

$$F(\rho) \equiv \langle \psi_+ | (I \otimes U^\dagger) \rho (I \otimes U) | \psi_+ \rangle = \sum_{m,n=1}^{2d^2} x_m x_n M_{mn}, \quad (4.126)$$

where M_{mn} is defined in the theorem. One can deduce that

$$M_{mn}^* = M_{nm} \quad (4.127)$$

from the hermiticity of ρ .

Taking into account the constraint with an undetermined Lagrange multiplier λ , we have

$$\frac{\partial}{\partial x_k} \{F(\rho) + \lambda (\sum_l x_l^2 - 1)\} = 0. \quad (4.128)$$

Accounting to (4.127) we have the eigenvalue equation

$$\sum_{n=1}^{2d^2} \text{Re}[M_{k,n}] x_n = -\lambda x_k. \quad (4.129)$$

Inserting (4.129) into (4.126) results in

$$\mathcal{F}(\rho) = \max_U F \leq \max_j \{\eta_j\}, \quad (4.130)$$

where $\eta_j = -\lambda_j$ is the corresponding eigenvalues of the real part of the matrix M .

□




Figure 2: Upper bound of $\mathcal{F}(\rho(a))$ from (4.124) (solid line) and upper bound from (4.115)(dashed line).

Example: Horodecki gives a very interesting bound entangled state in [20],

$$\rho(a) = \frac{1}{8a+1} \begin{pmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & a & 0 \\ a & 0 & 0 & 0 & a & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{pmatrix}. \quad (4.131)$$

One can easily compare the upper bound obtained in (4.124) and that in (4.115). From Fig. 2 we see that for $0 \leq a < 0.572$, the upper bound in (4.124) is larger than that in (4.115). But for $0.572 < a < 1$ the upper bound in (4.124) is always lower than that in (4.115), which means the upper bound (4.124) is tighter than (4.115).

In fact, we can drive another upper bound for FEF which will be very tight for weakly mixed quantum states.

Theorem 4.3 *For any bipartite quantum state $\rho \in \mathcal{H}_d \otimes \mathcal{H}_d$, the following inequality holds:*

$$\mathcal{F}(\rho) \leq \frac{1}{d} (\text{Tr}[\sqrt{\rho_A}])^2, \quad (4.132)$$

where ρ_A is the reduced matrix of ρ .

[Proof] Note that in [62] the authors have obtained the FEF for pure state $|\psi\rangle$,

$$\mathcal{F}(|\psi\rangle) = \frac{1}{d}(\text{Tr}[\sqrt{\rho_A^{|\psi\rangle}}])^2, \quad (4.133)$$

where $\rho_A^{|\psi\rangle}$ is the reduced matrix of $|\psi\rangle\langle\psi|$.

For mixed state $\rho = \sum_i p_i \rho^i$, we have

$$\begin{aligned} \mathcal{F}(\rho) &= \max_U \langle \psi_+ | (I \otimes U^\dagger) \rho (I \otimes U) | \psi_+ \rangle \leq \sum_i p_i \max_U \langle \psi_+ | (I \otimes U^\dagger) \rho^i (I \otimes U) | \psi_+ \rangle \\ &= \frac{1}{d} \sum_i p_i (\text{Tr}[\sqrt{\rho_A^i}])^2 = \frac{1}{d} \sum_i (\text{Tr}[\sqrt{p_i \rho_A^i}])^2. \end{aligned} \quad (4.134)$$

Let λ_{ij} be the real and nonnegative eigenvalues of the matrix $p_i \rho_A^i$. Recall that for any function $F = \sum_i (\sum_j x_{ij}^2)^{\frac{1}{2}}$ subjected to the constraints $z_j = \sum_i x_{ij}$ with x_{ij} being real and nonnegative, the inequality $\sum_j z_j^2 \leq F^2$ holds, from which it follows that

$$\mathcal{F}(\rho) \leq \frac{1}{d} \sum_i (\sum_j \sqrt{\lambda_{ij}})^2 \leq \frac{1}{d} (\sum_j \sqrt{\sum_i \lambda_{ij}})^2 = \frac{1}{d} (\text{Tr}[\sqrt{\rho_A}])^2, \quad (4.135)$$

which ends the proof. \square

4.2 Fully entangled fraction and concurrence

The upper bound of FEF has also interesting relations to the entanglement measure concurrence. As shown in [96], the concurrence of a two-qubit quantum state has some kinds of relation with the optimal teleportation fidelity. For quantum state with high dimension, we have the similar relation between them too.

Theorem 4.4 *For any bipartite quantum state $\rho \in \mathcal{H}_d \otimes \mathcal{H}_d$, we have*

$$C(\rho) \geq \sqrt{\frac{2d}{d-1}} [\mathcal{F}(\rho) - \frac{1}{d}]. \quad (4.136)$$

[Proof] In [98], the authors show that for any pure state $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$, the following inequality holds:

$$C(|\psi\rangle) \geq \sqrt{\frac{2d}{d-1}} (\max_{|\phi\rangle \in \varepsilon} |\langle \psi | \phi \rangle|^2 - \frac{1}{d}), \quad (4.137)$$

where ε denotes the set of $d \times d$ -dimensional maximally entangled states.

Let $\rho = \sum_i p_i |\phi_i\rangle\langle\phi_i|$ be the optimal decomposition such that $C(\rho) = \sum_i p_i C(|\psi_i\rangle)$. We have

$$\begin{aligned} C(\rho) &= \sum_i p_i C(|\psi_i\rangle) \geq \sum_i p_i \sqrt{\frac{2d}{d-1}} (\max_{|\phi\rangle \in \mathcal{E}} |\langle\psi_i|\phi\rangle|^2 - \frac{1}{d}) \\ &\geq \sqrt{\frac{2d}{d-1}} (\max_{|\phi\rangle \in \mathcal{E}} \sum_i p_i |\langle\psi_i|\phi\rangle|^2 - \frac{1}{d}) \\ &= \sqrt{\frac{2d}{d-1}} (\max_{|\phi\rangle \in \mathcal{E}} \langle\phi|\rho|\phi\rangle - \frac{1}{d}) = \sqrt{\frac{2d}{d-1}} (\mathcal{F}(\rho) - \frac{1}{d}), \end{aligned}$$

which ends the proof. \square

The inequality (4.136) has demonstrated the relation between the lower bound of concurrence and the fully entangled fraction (thus the optimal teleportation fidelity), i.e. the fully entangled fraction of a quantum state ρ is limited by its concurrence.

We now consider tripartite case. Let ρ_{ABC} be a state of three-qubit systems denoted by A , B and C . We study the upper bound of the FEF , $\mathcal{F}(\rho_{AB})$, between qubits A and B , and its relations to the concurrence under bipartite partition AB and C . For convenience we normalize $\mathcal{F}(\rho_{AB})$ to be

$$\mathcal{F}_N(\rho_{AB}) = \max\{2\mathcal{F}(\rho_{AB}) - 1, 0\}. \quad (4.138)$$

Let $C(\rho_{AB|C})$ denote the concurrence between subsystems AB and C .

Theorem 4.5 *For any triqubit state ρ_{ABC} , $\mathcal{F}_N(\rho_{AB})$ satisfies*

$$\mathcal{F}_N(\rho_{AB}) \leq \sqrt{1 - C^2(\rho_{AB|C})}. \quad (4.139)$$

[Proof] We first consider the case that ρ_{ABC} is pure, $\rho_{ABC} = |\psi\rangle_{ABC}\langle\psi|$. By using the Schmidt decomposition between qubits A, B and C , $|\psi\rangle_{ABC}$ can be written as:

$$|\psi\rangle_{AB|C} = \sum_{i=1}^2 \eta_i |i_{AB}\rangle |i_C\rangle, \quad \eta_1^2 + \eta_2^2 = 1, \quad \eta_1 \geq \eta_2 \quad (4.140)$$

for some orthonormalized bases $|i_{AB}\rangle$, $|i_C\rangle$ of subsystems AB , C respectively. The reduced density matrix ρ_{AB} has the form

$$\rho_{AB} = \text{Tr}_C[\rho_{ABC}] = \sum_{i=1}^2 \eta_i^2 |i_{AB}\rangle\langle i_{AB}| = U^T \Lambda U^*,$$

where Λ is a 4×4 diagonal matrix with diagonal elements $\{\eta_1^2, \eta_2^2, 0, 0\}$, U is a unitary matrix and U^* denotes the conjugation of U .

The *FEF* of the two-qubit state ρ_{AB} can be calculated by using formula (4.118) or the one in [96]. Let

$$M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & i \\ 0 & i & -1 & 0 \\ 0 & i & 1 & 0 \\ 1 & 0 & 0 & -i \end{pmatrix}$$

be the 4×4 matrix constituted by the four Bell bases. The *FEF* of ρ_{AB} can be written as

$$\begin{aligned} \mathcal{F}(\rho_{AB}) &= \eta_{\max}(\text{Re}\{M^\dagger \rho_{AB} M\}) = \frac{1}{2} \eta_{\max}(M^\dagger \rho_{AB} M + M^T \rho_{AB}^* M^*) \\ &\leq \frac{1}{2} [\eta_{\max}(M^\dagger U^T \Lambda U^* M) + \eta_{\max}(M^T U^\dagger \Lambda U M^*)] = \eta_1^2 \end{aligned} \quad (4.141)$$

where $\eta_{\max}(X)$ stands for the maximal eigenvalues of the matrix X .

For pure state (4.140) in bipartite partition AB and C , we have

$$C(|\psi\rangle_{AB|C}) = \sqrt{2(1 - \text{Tr}[\rho_{AB}^2])} = 2\eta_1\eta_2. \quad (4.142)$$

From (4.138), (4.141) and (4.142) we get

$$\mathcal{F}_N(\rho_{AB}) \leq \sqrt{1 - C^2(|\psi\rangle_{AB|C})}. \quad (4.143)$$

We now prove that the above inequality (4.143) also holds for mixed state ρ_{ABC} . Let $\rho_{ABC} = \sum_i p_i |\psi_i\rangle_{ABC} \langle \psi_i|$ be the optimal decomposition of ρ_{ABC} such that $C(\rho_{AB|C}) = \sum_i p_i C(|\psi_i\rangle_{AB|C})$. We have

$$\begin{aligned} \mathcal{F}_N(\rho_{AB}) &\leq \sum_i p_i \mathcal{F}_N(\rho_{AB}^i) \leq \sum_i p_i \sqrt{1 - C^2(\rho_{AB|C}^i)} \\ &\leq \sqrt{1 - \sum_i p_i C^2(\rho_{AB|C}^i)} \leq \sqrt{1 - C^2(\rho_{AB|C})}, \end{aligned}$$

where $\rho_{AB|C}^i = |\psi_i\rangle_{ABC} \langle \psi_i|$ and $\rho_{AB}^i = \text{Tr}_C[\rho_{AB|C}^i]$. \square

From Theorem 4.5 we see that the *FEF* of qubits A and B are bounded by the concurrence between qubits A , B and qubit C . The upper bound of *FEF* for ρ_{AB} decreases when the entanglement between qubits A , B and C increases. As an example, we consider the generalized W state defined by $|W'\rangle = \alpha|100\rangle + \beta|010\rangle + \gamma|001\rangle$, $|\alpha|^2 + |\beta|^2 + |\gamma|^2 = 1$. The reduced density matrix is given by

$$\rho_{AB}^{W'} = \begin{pmatrix} |\gamma|^2 & 0 & 0 & 0 \\ 0 & |\beta|^2 & \alpha^* \beta & 0 \\ 0 & \alpha \beta^* & |\alpha|^2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$




Figure 3: $\mathcal{F}_N(\rho_{AB}^{W'})$ (dashed line) and Upper bound $\sqrt{1 - C^2(|W'\rangle_{AB|C})}$ (solid line) of state $|W'\rangle_{AB|C}$ at $|\alpha| = |\beta|$.

The *FEF* of $\rho_{AB}^{W'}$ is given by

$$\mathcal{F}_N(\rho_{AB}^{W'}) = -\frac{1}{2} + 2|\alpha||\beta| + \frac{1}{2}||\alpha|^2 + |\beta|^2 - |\gamma|^2|.$$

While the concurrence of $|W'\rangle$ has the form $C_{AB|C}(|W'\rangle) = 2|\gamma|\sqrt{|\alpha|^2 + |\beta|^2}$. We see that (4.139) always holds. In particular for $|\alpha| = |\beta|$ and $|\gamma| \leq \frac{\sqrt{2}}{2}$, the inequality (4.139) is saturated (see Fig. 3).

4.3 Improvement of entanglement distillation protocol

The upper bound can give rise to not only an estimation of the fidelity in quantum information processing such as teleportation, but also an interesting application in entanglement distillation of quantum states. In [62] a generalized distillation protocol has been presented. It is shown that a quantum state ρ violating the reduction criterion can always be distilled. For such states if their single fraction of entanglement $F(\rho) = \langle \psi_+ | \rho | \psi_+ \rangle$ is greater than $\frac{1}{d}$, then one can distill these states directly by using the generalized distillation protocol. If the *FEF* (the largest value of single fraction of entanglement under local unitary transformations) is less than or equal to $\frac{1}{d}$, then a proper filtering operation has to be used at first to transform ρ to another state ρ' so that $F(\rho') > \frac{1}{d}$. For $d = 2$, one can compute *FEF* analytically according to the corollary. For $d \geq 3$ our upper bound (4.115) can supply a necessary condition in the distillation:

Theorem 4.6 *For an entangled state $\rho \in \mathcal{H} \otimes \mathcal{H}$ violating the reduction criterion,*




Figure 4: Upper bound of $\mathcal{F}(\rho) - \frac{1}{3}$ from (4.115) (solid line) and fidelity $F(\rho) - \frac{1}{3}$ (dashed line).

if the upper bound (4.115) is less than or equal to $\frac{1}{3}$, then the filtering operation has to be applied before using the generalized distillation protocol.

As an example we consider a 3×3 state

$$\rho = \frac{8}{9}\sigma + \frac{1}{9}|\psi_+\rangle\langle\psi_+|, \quad (4.144)$$

where $\sigma = (x|0\rangle\langle 0| + (1-x)|1\rangle\langle 1|) \otimes (x|0\rangle\langle 0| + (1-x)|1\rangle\langle 1|)$. It is direct to verify that ρ violates the reduction criterion for $0 \leq x \leq 1$, as $(\rho_1 \otimes I) - \rho$ has a negative eigenvalue $-\frac{2}{27}$. Therefore the state is distillable. From Fig. 4 we see that for $0 \leq x < 0.0722$ and $0.9278 < x \leq 1$, the fidelity is already greater than $\frac{1}{3}$, thus the generalized distillation protocol can be applied without the filtering operation. However for $0.1188 \leq x \leq 0.8811$, even the upper bound of the fully entangled fraction is less than or equal to $\frac{1}{3}$, hence the filtering operation has to be applied first, before using the generalized distillation protocol.

Moreover, the lower bounds of concurrence can be also used to study the distillability of quantum states. Based on the positive partial transpose (PPT) criterion, a necessary and sufficient condition for the distillability was proposed in [99], which is not operational in general. An alternative distillability criterion based on the bound τ_2 in (3.67) can be obtained to improve the operatinality.

Theorem 4.7 *A bipartite quantum state ρ is distillable if and only if $\tau_2(\rho^{\otimes N}) > 0$ for some number N .*

[Proof] It was shown in [99] that a density matrix ρ is distillable if and only if there are some projectors P, Q that map high dimensional spaces to two-dimensional ones and some number N such that the state $P \otimes Q \rho^{\otimes N} P \otimes Q$ is entangled [99]. Thus if $\tau_2(\rho^{\otimes N}) > 0$, there exists one submatrix of matrix $\rho^{\otimes N}$, similar to Eq. (3.68), which has nonzero τ_2 and is entangled in a $2 \otimes 2$ space, hence ρ is distillable. \square

Corollary 4.7 *The lower bound $\tau_2(\rho) > 0$ is a sufficient condition for the distillability of any bipartite state ρ .*

Corollary 4.7 *The lower bound $\tau_2(\rho) = 0$ is a necessary condition for separability of any bipartite state ρ .*

Remark: Corollary 4.7 directly follows from Theorem 4.7 and this case is referred to as one-distillable [100]. The problem of whether non-PPT (NPPT) nondistillable states exist is studied numerically in [100, 101]. By using Theorem 4.7, although it seems impossible to solve the problem completely, it is easy to judge the distillability of a state under condition that it is one-distillable.

The lower bound τ_2 , PPT criterion, separability and distillability for any bipartite quantum state ρ have the following relations: if $\tau_2(\rho) > 0$, ρ is entangled. If ρ is separable, it is PPT. If $\tau_2(\rho) > 0$, ρ is distillable. If ρ is distillable, it is NPPT. From the last two propositions it follows that if ρ is PPT, $\tau_2(\rho) = 0$, i.e., if $\tau_2(\rho) > 0$, ρ is NPPT.

Theorem 4.8 *For any pure tripartite state $|\phi\rangle_{ABC}$ in arbitrary $d \otimes d \otimes d$ dimensional spaces, bound τ_2 satisfies*

$$\tau_2(\rho_{AB}) + \tau_2(\rho_{AC}) \leq \tau_2(\rho_{A:BC}), \quad (4.145)$$

where $\rho_{AB} = \text{Tr}_C(|\phi\rangle_{ABC}\langle\phi|)$, $\rho_{AC} = \text{Tr}_B(|\phi\rangle_{ABC}\langle\phi|)$, and $\rho_{A:BC} = \text{Tr}_{BC}(|\phi\rangle_{ABC}\langle\phi|)$.

[Proof] Since $C_{mn}^2 \leq \left(\lambda_{mn}^{(1)}\right)^2 \leq \sum_{i=1}^4 \left(\lambda_{mn}^{(i)}\right)^2 = \text{Tr}(\rho \tilde{\rho}_{mn})$, one can derive the inequality:

$$\tau(\rho_{AB}) + \tau(\rho_{AC}) \leq \sum_{l,k}^D \text{Tr} [\rho_{AB}(\tilde{\rho}_{AB})_{lk}] + \sum_{p,q}^D \text{Tr} [\rho_{AC}(\tilde{\rho}_{AC})_{pq}], \quad (4.146)$$

where $D = d(d-1)/2$. Note that $\sum_{lk} \text{Tr} [\rho_{AB}(\tilde{\rho}_{AB})_{lk}] \leq 1 - \text{Tr} \rho_A^2 - \text{Tr} \rho_B^2 + \text{Tr} \rho_C^2$ and $\sum_{pq} \text{Tr} [\rho_{AC}(\tilde{\rho}_{AC})_{pq}] \leq 1 - \text{Tr} \rho_A^2 + \text{Tr} \rho_B^2 - \text{Tr} \rho_C^2$, where $l, pk, q, = 1, \dots, D$. By

using the similar analysis in [102] one has that the right-hand side of Eq. (4.146) is equal to $2(1 - \text{Tr}\rho_A^2) = C^2(\rho_{A:BC})$. Taking into account that $\tau_2(\rho_{A:BC}) = C^2(\rho_{A:BC})$ for a pure state, one obtains the inequality (4.145). \square

Generally for any pure multipartite quantum state $\rho_{AB_1B_2\dots b_n}$, one has the following monogamy inequality:

$$\tau_2(\rho_{AB_1}) + \tau_2(\rho_{AB_2}) + \dots + \tau_2(\rho_{AB_n}) \leq \tau_2(\rho_{A:B_1B_2\dots B_n}).$$

5 Summary and Conclusion

We have introduced some recent results on three aspects in quantum information theory. The first one is the separability of quantum states. New criteria to detect more entanglements have been discussed. The normal form of quantum states have been also studied, which helps in investigating the separability of quantum states. Moreover, since many kinds of quantum states can be transformed into the same normal forms, quantum states can be classified in terms of the normal forms. For the well known entanglement measure concurrence, we have discussed the tight lower and upper bounds. It turns out that although one can not distill a singlet from many pairs of bound entangled states, the concurrence and tangle of two entangled quantum states are always larger than that of one, even both two entangled quantum states are bound entangled. Related to the optimal teleportation fidelity, upper bounds for the fully entangled fraction have been studied, which can be used to improve the distillation protocol. Interesting relations between fully entangled fraction and concurrence have been also introduced. All these related problems in the theory of quantum entanglement have not been completely solved yet. Many problems remain open concerning the physical properties and mathematical structures of quantum entanglement, and the applications of entangled states in information processing.

References

- [1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. 47, 777(1935).
- [2] A. Peres, Quantum Mechanics: Concepts and Methods, Kluwer, Dordrecht(1993).

- [3] M.A. Nielsen and I.L. Chuang, Quantum Computation and Quantum Information. Cambridge: Cambridge University Press, (2000).
- [4] See, for example, D.P. Di Vincenzo, Science 270,255(1995)
- [5] C.H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres and W.K. Wootters, Phys. Rev. Lett. 70, 1895 (1993);
S. Albeverio and S.M. Fei, Phys. Lett. A 276, 8(2000);
G.M. D'Ariano, P. Lo Presti and M.F. Sacchi, Phys. Lett. A 272, 32(2000).
- [6] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. A, 60, 1888(1999);
S. Albeverio, S.M. Fei, and W. L. Yang, Phys. Rev. A, 66, 012301(2002).
- [7] C.H. Bennett and S.J. Wiesner, Phys. Rev. Lett. **69**, 2881(1992).
- [8] A. Ekert, Phys. Rev. Lett. 67, 661(1991);
D. Deutsch, A. Ekert, P. Rozas, C. Macchicavello, S. Popescu and A. Sanpera, Phys. Rev. Lett. 77, 2818(1996);
C.A. Fuchs, N. Gisin, R.B. Griffiths, C.S. Niu and A. Peres, Phys. Rev. A 56, 1163(1997).
- [9] M. Żukowski, A. Zeilinger, M.A. Horne and A.K. Ekert, Phys. Rev. Lett. 71, 4287(1993);
S. Bose, V. Vedral and P.L. Knight, Phys. Rev. A 57, 822(1998); 60, 194(1999);
B.S. Shi, Y.K. Jiang, G.C. Guo, Phys. Rev. A 62, 054301(2000);
L. Hardy and D.D. Song, Phys. Rev. A 62, 052315(2000).
- [10] C.H. Bennett, D.P. DiVincenzo, P.W. Shor, J.A. Smolin, B.M. Terhal and W.K. Wootters, Phys. Rev. Lett. 87, 077902(2001).
- [11] B.S. Shi and A. Tomita, J. Opt. B: Quant. Semiclass. Opt. 4, 380(2002);
J.M. Liu and Y.Z. Wang, Chinese Phys. 13, 147(2004).
- [12] D.W. Leung and P.W. Shor, Phys. Rev. Lett. 90, 127905(2003);
Sanders, Phys. Rev. Lett. 90, 057901(2003);
S.A. Babichev, B. Brezger and A.I. Lvovsky, Phys. Rev. Lett. 92, 047903 (2004).
- [13] M.Y. Ye, Y.S. Zhang and G.C. Guo, Phys. Rev. A 69, 022310(2004).
- [14] R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, Rev. Modern Phys. 81, 865(2009).

- [15] O. Guhne and G. Toth, Phys. Reports 474, 1(2009).
- [16] S. Albeverio and S.M. Fei, J. Opt. B: Quantum Semiclass Opt, 3, 223(2001).
- [17] R.F. Werner, Phys. Rev. A 40, 4277(1989).
- [18] A. Peres, Phys. Rev. Lett. 77, 1413(1996).
- [19] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Lett. A 223, 1(1996).
- [20] P. Horodecki, Phys. Lett. A 232, 333(1997).
- [21] O. Rudolph, Phys. Rev. A 67, 032312 (2003).
- [22] K. Chen and L.A. Wu, Quant. Inf. Comput. 3, 193 (2003).
- [23] M. Horodecki, P. Horodecki and R. Horodecki, Open Syst. Inf. Dyn. 13, 103 (2006).
- [24] K. Chen and L.A. Wu, Phys. Lett. A 306, 14 (2002);
O. Rudolph, Phys. Rev. A 67, 032312 (2003);
K. Chen and L.A. Wu, Phys. Rev. A 69, 022312 (2004);
P. Wocjan and M. Horodecki, Quant. Inf. and Comput. 6, 277 (2006).
- [25] S. Albeverio, K. Chen and S.M. Fei, Phys. Rev. A 68, 062313 (2003).
- [26] M. Horodecki and P. Horodecki, Phys. Rev. A 59, 4206 (1999).
- [27] N.J. Cerf, C. Adami and R.M. Gingrich, Phys. Rev. A 60, 898 (1999).
- [28] M.A. Nielsen and J. Kempe, Phys. Rev. Lett. 86, 5184(2001).
- [29] H. F. Hofmann and S. Takeuchi, Phys. Rev. A 68, 032103 (2003).
- [30] O. Gühne, M. Mechler, G. Tóth and P. Adam, Phys. Rev. A 74, 010301(R)(2006).
- [31] O. Gühne, Phys. Rev. Lett. 92, 117903 (2004).
- [32] C.J. Zhang, Y.S. Zhang, S. Zhang and G.C. Guo. Phys. Rev. A 76, 012334 (2007).
- [33] J.D. Vicente, Quantum Inf. Comput. 7, 624(2007).
- [34] J.D. Vicente, J. Phys. A: Math. and Theor., 41, 065309(2008).

- [35] O. Gühne, P. Hyllus, O. Gittsovich, and J. Eisert, Phys. Rev. Lett. 99, 130504 (2007).
- [36] O. Gittsovich, O. Gühne, P. Hyllus, and J. Eisert, Phys. Rev. A **78**, 052319(2008).
- [37] C.H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. Smolin, and W.K. Wootters, Phys. Rev. Lett. 78, 2031(1996).
- [38] D. Brub, J. Math. Phys. 43, 4237(2002).
- [39] W.K. Wootters, Phys. Rev. Lett. 80, 2245 (1998).
- [40] B.M. Terhal, K.G.H. Vollbrecht, Phys. Rev. Lett. 85, 2625-2628(2000).
- [41] S.M. Fei, J. Jost, X.Q. Li-Jost and G.F. Wang, Phys. Lett. A 310, 333-338(2003).
- [42] S.M. Fei and X.Q. Li-Jost, Rep. Math. Phys. 53, 195-210(2004).
- [43] S.M. Fei, Z.X. Wang and H. Zhao, Phys. Lett. A 329, 414-419(2004).
- [44] P. Rungta and C.M. Caves, Phys Rev A 67, 012307(2003).
- [45] F. Mintert, M. Kus, A. Buchleitner, Phys. Rev. Lett. 92, 167902(2004).
- [46] K. Chen, S. Albeverio and S.M. Fei, Phys. Rev. Lett. 95, 210501(2005).
- [47] K. Chen, S. Albeverio, and S. M. Fei, Phys. Rev. Lett. 95, 040504(2005).
- [48] H.P. Breuer, J. Phys. A 39, 11847(2006).
- [49] H.P. Breuer, Phys. Rev. Lett. 97, 080501(2006).
- [50] J.D. Vicente, Phys, Rev. A 75, 052320(2007).
- [51] X.H. Gao, S.M. Fei and K. Wu, Phys. Rev. A 74, 050303(R) (2006).
- [52] E. Gerjuoy, Phys. Rev. A 67, 052308(2003).
- [53] Y.C. Ou, H. Fan and S.M. Fei, Phys. Rev. A 78, 012311(2008).
- [54] J.I. de Vicente, J. Phys. A: Math. Theor. 41, 065309 (2008).
- [55] F. Mintert, M. Kus, A. Buchleitner, Phys. Rev. Lett. 95, 260502(2005).

- [56] S.P. Walborn, P.H. Souto Ribeiro, L. Davidovich, F. Mintert, and A. Buchleitner, *Nature (London)* 440, 1022(2006);
S.P. Walborn, P.H. Souto Ribeiro, L. Davidovich, F. Mintert, and A. Buchleitner, *Phys. Rev. A* 75, 032338 (2007).
- [57] S.M. Fei, M.J. Zhao, K. Chen and Z.X. Wang, *Experimental Determination of Entanglement for Arbitrary Pure States*, to appear in *Phys. Rev. A*, (2009).
- [58] C.H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. Smolin, and W.K. Wootters, *Phys. Rev. Lett.* 76, 722(1996).
- [59] M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* 78, 574(1997).
- [60] D. Deutsch, A. Ekert, R. Jozsa, C. Macchiavello, S. Popescu, and A. Sanpera, *Phys. Rev. Lett.* 77, 2818(1996).
- [61] C.H. Bennett, D.P. DiVincenzo, J.A. Smolin, and W.K. Wootters, *Phys. Rev. A* 54, 3824 (1996).
- [62] M. Horodecki and P. Horodecki, *Phys. Rev. A*, 59, 4206(2002).
- [63] A. Acin, A. Andrianov, L. Costa, E. Jane, J.I. Latorre, R. Tarrach, *Phys. Rev. Lett.* 85, 1560 (2000).
- [64] A. Sanpera, R. Tarrach, and G. Vidal, *Phys. Rev. A* 58, 826(1998).
- [65] N.J. Cerf, C. Adami, and R.M. Gingrich, *Phys. Rev. A* 60, 898 (1999).
- [66] M. Horodecki and P. Horodecki, *Phys. Rev. A* 54, 2406 (1999).
- [67] O. Rudolph, *J. Phys. A* 36, 5825(2003).
- [68] F.T. Hioe and J.H. Eberly, *Phys. Rev. Lett.* 47, 838(1981).
- [69] Walter Greiner and Berndt Muller. *Quantum Mechanics. Symmetries*, Springer-Verlag Berlin Heidelberg, 1989.
- [70] J.E. Harriman, *Phys. Rev. A* 17, 1249(1978).
- [71] S.J. Wu and J. Anandan, *Phys. Lett. A* 297, 4(2002).
- [72] O. Rudolph, *Quan. Inform. Proc.* 4, 219(2005).
- [73] R.A. Horn and C.R. Johnson, *Matrix Analysis* (Cambridge University Press, Cambridge, U.K., 1985), Theorem 7.4.9.

- [74] S.X. Yu and N.L. Liu, Phys. Rev. Lett. 95, 150504 (2005).
- [75] F. Verstraete, J. Dehaene, and B. De Moor, Phys. Rev. A 68, 012103(2003).
- [76] J.M. Leinaas, J.M. and E. Ovrum, Phys. Rev. A 74, 012313 (2006).
- [77] F. Verstraete, Ph. K. Thesis, Katholieke Universiteit Leuven (2002).
- [78] J.I. de Vicente, Quantum Inf. Comput. 7, 624 (2007).
- [79] A.S.M. Hassan and P.S. Joag, Quant. Inform. Comput. 8, 0773(2008).
- [80] G. Vidal, D. Jonathan, and M.A. Nielsen, Phys. Rev. A (3) 62, 012304 (2000);
V. Vedral and M.B. Plenio, Phys. Rev. A 57, 1619 (1998).
- [81] S. Hill and W.K. Wootters, Phys. Rev. Lett. 78, 5022(1997).
- [82] A. Osterloh et al., Nature 416, 608 (2002);
L.A. Wu, M.S. Sarandy, and D.A. Lidar, Phys. Rev. Lett. 93, 250404 (2004).
- [83] S. Ghosh, T.F. Rosenbaum, G. Aeppli, S.N. Coppersmith, Nature 425, 48
(2003);
V. Vedral, Nature 425, 28 (2003).
- [84] F. Mintert, M. Kus, and A. Buchleitner, Phys. Rev. Lett. 92, 167902 (2004);
F. Mintert, Ph.D. thesis, Munich University, 2004.
- [85] P. Rungta, V. Bužek, C.M. Caves, M. Hillery, and G.J. Milburn, Phys. Rev. A
64, 042315 (2001).
A. Uhlmann, Phys. Rev. A 62, 032307(2000).
- [86] L. Aolita and F. Mintert, Phys. Rev. Lett. 97, (2006) 050501;
A.R.R. Carvalho, F. Mintert, and A. Buchleitner, Phys. Rev. Lett. 93,
230501(2004).
- [87] V. Coffman, J. Kundu, and W.K. Wootters, Phys. Rev. A 61, 052306 (2000).
- [88] M. Zukowski, A. Zeilinger, M.A. Horne and A.K. Ekert, Phys. Rev. Lett. 71,
4287 (1993).
- [89] J.M. Cai, Z.W. Zhou, S. Zhang, and G.C. Guo, Phys. Rev. A 75, 052324(2007).
- [90] L. Aolita, A. Buchleitner, and F. Mintert, Phys. Rev. A 78, 022308(2008).

- [91] V. Coffman, J. Kundu, and W.K. Wootters, Phys. Rev. A 61, 052306 (2000); Tobias J. Osborne, Phys. Rev. A 72, 022309(2005).
- [92] L. Masanes, Phys. Rev. Lett. 96, 150501 (2006).
- [93] G. Bowen, S. Bose, Phys. Rev. Lett. 87, 267901(2001).
- [94] S. Albeverio, S.M. Fei and W.L. Yang, Commun. Theor. Phys. 38, 301(2002).
- [95] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. A, 60, 1888(1999); S. Albeverio, S.M. Fei, and W.L. Yang, Phys. Rev. A, 66, 012301(2002).
- [96] J. Grondalski and D.M. Etlinger, D. F. V. James, Phys. Lett. A 300, 573(2002).
- [97] R.F. Werner, J. Phys. A: Math. Gen. 34, 7081(2001).
- [98] C.S. Yu, X.X. Yi, and H.S. Song, Phys. Rev. A 78, 062330 (2008).
- [99] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 80, 5239(1996).
- [100] W. Dr, J. I. Cirac, M. Lewenstein, and D. Bruss, Phys. Rev. A 61, 062313(2000).
- [101] D. P. DiVincenzo, P. W. Shor, J. A. Smolin, B. M. Terhal, and A. V. Thapliyal, Phys. Rev. A 61, 062312(2000).
- [102] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A 61, 052306(2000).