

Quantum conjugacy classes of simple matrix groups*

A. Mudrov

Dedicated to the memory of Joseph Donin

Emmy Noether Mathematics Institute, 52900 Ramat Gan, Israel,
Max-Planck Institut für Mathematik, Vivatsgasse 7, D-53111 Bonn, Germany.
e-mail: mudrova@macs.biu.ac.il, mudrov@mpim-bonn.mpg.de

Abstract

Let G be a simple complex classical group and \mathfrak{g} its Lie algebra. Let $\mathcal{U}_\hbar(\mathfrak{g})$ be the Drinfeld-Jimbo quantization of the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$. We construct an explicit $\mathcal{U}_\hbar(\mathfrak{g})$ -equivariant quantization of conjugacy classes of G with Levi subgroups as the stabilizers.

Key words: Quantum groups, equivariant quantization, quantum conjugacy classes.

Contents

1	Introduction	2
2	Drinfeld-Jimbo quantum group	4
2.1	Quantized universal enveloping algebra	5
2.2	Defining representations of classical matrix groups	6
2.3	Parabolic subalgebras in $\mathcal{U}_\hbar(\mathfrak{g})$	7
3	Generalized Verma modules over $\mathcal{U}_\hbar(\mathfrak{g})$	8
3.1	Upper and lower (generalized) Verma modules	9
3.2	Pairing between upper and lower generalized Verma modules	11
3.3	Tensor product of finite dimensional and generalized Verma modules	12

*This research is partially supported by the Emmy Noether Research Institute for Mathematics, the Minerva Foundation of Germany, the Excellency Center "Group Theoretic Methods in the study of Algebraic Varieties" of the Israel Science foundation, and by the RFBR grant no. 03-01-00593.

4	Properties of the universal RE matrix	13
4.1	Characteristic polynomial for RE matrix	14
4.2	Minimal polynomial for RE matrix	15
4.3	A construction of central elements	15
5	On quantization of affine algebraic varieties	16
5.1	A flatness criterion	17
6	Quantization of simple algebraic groups	18
6.1	Simple groups as Poisson Lie manifolds	18
6.2	Quantization of the STS bracket on the group	19
6.3	Embedding of $\mathbb{C}_\hbar[G]$ in $\mathcal{U}_\hbar(\mathfrak{g})$	20
7	Center of the algebra $\mathbb{C}_\hbar[G]$	21
7.1	The case of classical series	22
8	Quantization of conjugacy classes	23
8.1	Non-exceptional classes	23
8.2	The quantization theorem	24
8.3	Ideals of quantized non-exceptional conjugacy classes	26
8.3.1	The $\mathcal{U}_\hbar(\mathfrak{gl}(n))$ -case	27
8.3.2	The $\mathcal{U}_\hbar(\mathfrak{so}(2n + 1))$ -case	28
8.3.3	The $\mathcal{U}_\hbar(\mathfrak{sp}(n))$ -case	28
8.3.4	The $\mathcal{U}_\hbar(\mathfrak{so}(2n))$ -case	28
A	More on central characters	29

1 Introduction

Deformation quantization of a Poisson structure on a smooth manifold is a classic problem of mathematical physics. Especially interesting is a quantization that is equivariant with respect to an action of a group and, even more generally, a quantum group. Recently, a significant progress in this field was triggered by a discovered connection between equivariant quantization and the theory of dynamical Yang-Baxter equation, [DM1]. Namely, a star product quantization of semisimple coadjoint orbits and conjugacy classes (with a Levi

subgroup as the stabilizer) of simple Lie groups was constructed in [EE, EEM] in terms of the universal dynamical twist.

On the other hand, semisimple coadjoint orbits and conjugacy classes are affine algebraic varieties. Therefore their quantization may be sought for in terms of generators and relations, as a deformed ring of polynomial functions. This is an alternative approach as compared to star product and it has certain advantages because the solution is formulated by "finite data".

In the present paper, we construct an equivariant deformation quantization of semisimple conjugacy classes of simple classical algebraic groups with Levi subgroups as the stabilizers. More precisely, let G be a complex simple algebraic group from the series A , B , C , and D . Fix the standard, or Drinfeld-Jimbo, factorizable quasitriangular Lie bialgebra structure on $\mathfrak{g} = \text{Lie } G$. Equipped with the corresponding Drinfeld-Sklyanin (DS) bracket, G becomes a Poisson Lie group. Consider the adjoint action of G on itself. The Semenov-Tyan-Shansky (STS) Poisson bracket makes G a Poisson Lie manifold over G . The symplectic leaves of this Poisson structure are exactly the conjugacy classes. Let $\mathcal{U}_\hbar(\mathfrak{g})$ be the Drinfeld-Jimbo quantum group. We construct a $\mathcal{U}_\hbar(\mathfrak{g})$ -equivariant quantization of the ring of polynomial functions along the STS bracket on almost all conjugacy classes with Levi subgroups as the stabilizers. Almost all means all for special linear and symplectic groups. For the orthogonal groups our construction covers the classes of matrices with eigenvalues λ subject to the condition $\lambda^2 = 1 \Rightarrow \lambda = 1$. Those are exactly the classes that are isomorphic to semisimple G -orbits in \mathfrak{g}^* (via the Cayley transformation).

The quantization is given explicitly, in terms of deformed ideals of classes in a quantized ring of polynomial functions on the group (or in the so called reflection equation algebra). Simultaneously, the quantized classes are realized as subalgebras of operators on generalized Verma modules.

The setup of the paper is as follows.

In Section 2 we recall definition of the Drinfeld-Jimbo quantum group and its some important subalgebras.

In Section 3 we study tensor products of finite dimensional and generalized Verma modules over $\mathcal{U}_\hbar(\mathfrak{g})$.

In Section 4 we study properties of a fundamental object of our theory, the universal reflection equation matrix $\mathcal{Q} = \mathcal{R}_{21}\mathcal{R}$ (expressed through the universal R-matrix of $\mathcal{U}_\hbar(\mathfrak{g})$). In particular, we determine its spectrum on tensor products of finite dimensional and generalized Verma modules and compute q-traces of \mathcal{Q}^ℓ .

In Section 5 we develop a method of quantization of affine homogeneous varieties.

In Section 6 we recall some results of [M] on quantization of simple algebraic groups and construct an embedding of the quantized affine coordinate ring $\mathbb{C}_\hbar[G]$ in $\mathcal{U}_\hbar(\mathfrak{g})$.

In Section 7 we recall results of [M] concerning the center of $\mathbb{C}_\hbar[G]$.

In Section 8 we give the quantization of the conjugacy classes.

Appendix contains auxiliary information about invariants of $\mathcal{U}_\hbar(\mathfrak{g})$.

Acknowledgements. The author is grateful to the Max-Planck Institute for Mathematics in Bonn for hospitality and the excellent research conditions. He thanks J. Bernstein for his interest to the present work, helpful discussions and valuable remarks.

2 Drinfeld-Jimbo quantum group

In the present paper, we work over the ring $\mathbb{C}[[\hbar]]$ of formal power series in \hbar . Given a $\mathbb{C}[[\hbar]]$ -module E we denote by E_0 its quotient $E/\hbar E$.

By a deformation of a complex vector space E_0 we mean a free $\mathbb{C}[[\hbar]]$ -module E such that $E/\hbar E \simeq E_0$. Deformation of an associative algebra \mathcal{A}_0 is a $\mathbb{C}[[\hbar]]$ -algebra \mathcal{A} such that $\mathcal{A}/\hbar\mathcal{A} \simeq \mathcal{A}_0$ as associative \mathbb{C} -algebras. The term *deformation quantization* or simply *quantization* is reserved for deformation of commutative algebras.

Quantized universal enveloping algebras are understood as $\mathbb{C}[[\hbar]]$ -algebras, [Dr1]. We will work with the standard or Drinfeld-Jimbo quantization of simple Lie algebras. We will assume \hbar -adic completion of the Cartan subalgebra only. That is possible thanks to the existence of the Poincaré-Birkhoff-Witt base over the Cartan subalgebra. Thus defined, $\mathcal{U}_\hbar(\mathfrak{g})$ is a Hopf algebra in a weaker sense. However that is sufficient for our purposes, because we will deal with \mathfrak{h} -diagonalizable $\mathcal{U}_\hbar(\mathfrak{g})$ -modules.

Although the quasitriangular structure requires completion of tensor products, we will use this structure in the situation when one of the modules is $\mathbb{C}[[\hbar]]$ -finite. This does not lead out of $\mathcal{U}_\hbar(\mathfrak{g})$.

We assume that $\mathcal{U}_\hbar(\mathfrak{g})$ -modules are equipped with \hbar -adic topology and the action of $\mathcal{U}_\hbar(\mathfrak{g})$ is continuous. However we do not require the modules to be complete. We will work with $\mathcal{U}_\hbar(\mathfrak{g})$ -modules that are direct sums of $\mathbb{C}[[\hbar]]$ -finite weight spaces. Each weight space is complete, being $\mathbb{C}[[\hbar]]$ -finite. Thus our point of view is self-consistent, because we assume \hbar -adic completion for the Cartan subalgebra only.

2.1 Quantized universal enveloping algebra

Let \mathfrak{g} be a complex semisimple Lie algebra and \mathfrak{h} its Cartan subalgebra. Let R denote the root system of \mathfrak{g} with a fixed subsystem of positive roots $R_+ \subset R$. By $\Pi \subset R_+$ we denote the subset of simple roots. Let (\cdot, \cdot) denote the Killing form on \mathfrak{g} . We will use the same notation for the invariant scalar product on \mathfrak{h}^* that is induced by the Killing form restricted to \mathfrak{h} . By W we denote the Weyl group of \mathfrak{g} .

For every $\lambda \in \mathfrak{h}^*$ we denote by h_λ its image under the isomorphism $\mathfrak{h}^* \simeq \mathfrak{h}$ implemented by the Killing form. In other words, $\lambda(h) = (h_\lambda, h)$ for all $h \in \mathfrak{h}$. We put $\rho = \frac{1}{2} \sum_{\alpha \in R_+} \alpha$, the half-sum of the positive roots. Its dual $h_\rho \in \mathfrak{h}$ is the unique solution of the system of linear equations $\alpha_i(h_\rho) = \frac{1}{2}(\alpha_i, \alpha_i)$, $i = 1, \dots, \text{rk } \mathfrak{g}$.

Denote by $\mathcal{U}_\hbar(\mathfrak{h})$ the \hbar -adic completion of the algebra $\mathcal{U}(\mathfrak{h}) \otimes \mathbb{C}[[\hbar]]$. Define $\mathcal{U}_\hbar(\mathfrak{g})$ as a $\mathbb{C}[[\hbar]]$ -algebra generated by the elements $e_{\pm\alpha}$, $\alpha \in \Pi$, over $\mathcal{U}_\hbar(\mathfrak{h})$. These generators are subject to the following relations, [Dr1, Ji]:

$$[h_{\alpha_i}, e_{\pm\alpha_j}] = \pm(\alpha_i, \alpha_j)e_{\pm\alpha_j}, \quad [e_{\alpha_i}, e_{-\alpha_j}] = \delta_{ij} \frac{q^{h_{\alpha_i}} - q^{-h_{\alpha_i}}}{q_i - q_i^{-1}},$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1 - a_{ij} \\ k \end{bmatrix}_{q_i} e_{\pm\alpha_i}^{1-a_{ij}-k} e_{\pm\alpha_j} e_{\pm\alpha_i}^k = 0,$$

where $a_{ij} = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$, $i, j = 1, \dots, \text{rk } \mathfrak{g}$, is the Cartan matrix, $q := e^\hbar$, $q_i := e^{\frac{\hbar}{2}(\alpha_i, \alpha_i)}$, and

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}, \quad [n]_q! = [1]_q \cdot [2]_q \cdots [n]_q, \quad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

Strictly speaking, $\mathcal{U}_\hbar(\mathfrak{g})$ is not a Hopf algebra, because Δ takes values in the completed tensor square of $\mathcal{U}_\hbar(\mathfrak{g})$. However, the category of $\mathcal{U}_\hbar(\mathfrak{g})$ -modules that are \mathfrak{h} -diagonalizable is monoidal. So we will understand Hopf algebras in this weaker sense.

The coproduct Δ , antipode γ , and counit ϵ in $\mathcal{U}_\hbar(\mathfrak{g})$ are given by

$$\begin{aligned} \Delta(e_\alpha) &= e_\alpha \otimes 1 + q^{h_\alpha} \otimes e_\alpha, & \Delta(e_{-\alpha}) &= e_{-\alpha} \otimes q^{-h_\alpha} + 1 \otimes e_{-\alpha}, \\ \Delta(h_\alpha) &= h_\alpha \otimes 1 + 1 \otimes h_\alpha, \\ \gamma(e_\alpha) &= -e_\alpha q^{-h_\alpha}, & \gamma(e_{-\alpha}) &= -q^{h_\alpha} e_{-\alpha}, & \gamma(h_\alpha) &= -h_\alpha. \\ \epsilon(h) &= \epsilon(e_{\pm\alpha}) = 0, \end{aligned}$$

for all $\alpha \in \Pi$. The correspondence $e_{-\alpha} \mapsto e_\alpha q^{h_\alpha}$, $e_\alpha \mapsto q^{-h_\alpha} e_{-\alpha}$, $h_\alpha \mapsto h_\alpha$, for all $\alpha \in \Pi$, extends to an involutive anti-algebra and coalgebra transformation of $\mathcal{U}_\hbar(\mathfrak{g})$ denoted by ω .

The $\mathbb{C}[[\hbar]]$ -adic completion of $\mathcal{U}_\hbar(\mathfrak{g})$ is isomorphic to $\mathcal{U}(\mathfrak{g})[[\hbar]]$ as an associative algebra, [Dr3].

The subalgebras in $\mathcal{U}_\hbar(\mathfrak{g})$ generated over $\mathcal{U}_\hbar(\mathfrak{h})$ by $\{e_{+\alpha}\}_{\alpha \in \Pi}$ and by $\{e_{-\alpha}\}_{\alpha \in \Pi}$, respectively, are Hopf algebras. They are quantized universal enveloping algebras of the positive and negative Borel subalgebras \mathfrak{b}^\pm and denoted further by $\mathcal{U}_\hbar(\mathfrak{b}^\pm)$.

The elements $\{e_{\pm\alpha}\}_{\alpha \in \Pi} \subset \mathcal{U}_\hbar(\mathfrak{g})$ are called quantum Chevalley generators. The Chevalley generators can be extended to a system of quantum Cartan-Weyl generators $\{e_{\pm\alpha}\}_{\alpha \in \mathbb{R}_+}$ via the so called q -commutators, see [KhT] and references therein. The Cartan-Weyl base admits an ordering, with respect to which the monomials $\prod_{\alpha \in \mathbb{R}_+}^> e_{-\alpha}^{r_\alpha} \prod_{\alpha \in \mathbb{R}_+}^< e_\alpha^{s_\alpha}$ form a Poincaré-Birkhoff-Witt (PBW) base of $\mathcal{U}_\hbar(\mathfrak{g})$ as a $\mathcal{U}_\hbar(\mathfrak{h})$ -module, [KhT]. Denote by $\mathcal{U}'_\hbar(\mathfrak{b}^\pm)$ the ideals in $\mathcal{U}_\hbar(\mathfrak{b}^\pm)$ generated by $\{e_{\pm\alpha}\}_{\alpha \in \Pi}$. The PBW monomials of positive and, respectively, negative weights with respect to \mathfrak{h} form bases for $\mathcal{U}'_\hbar(\mathfrak{b}^\pm)$ over $\mathcal{U}_\hbar(\mathfrak{h})$.

The universal R-matrix belongs to the completed tensor square of $\mathcal{U}_\hbar(\mathfrak{g})$. It has the structure

$$\mathcal{R} = q^{\Omega_\mathfrak{h}} \text{ mod } \mathcal{U}'_\hbar(\mathfrak{b}^-) \hat{\otimes} \mathcal{U}'_\hbar(\mathfrak{b}^+), \quad (1)$$

where $\Omega_\mathfrak{h} \in \mathfrak{h} \otimes \mathfrak{h}$ is the inverse to the Killing form (the canonical element) restricted to \mathfrak{h} . More precisely, the R-matrix can be represented as the product of the Cartan factor $q^{\Omega_\mathfrak{h}}$ and a series in the Cartan-Weyl generators, [KhT]. The universal R-matrix (1) is a "quantization" of the classical r-matrix

$$r_- = \frac{1}{2} \sum_{\alpha \in \mathbb{R}^+} (\alpha, \alpha) (e_{-\alpha} \otimes e_\alpha - e_\alpha \otimes e_{-\alpha}) \quad (2)$$

called the standard or Drinfeld-Jimbo r-matrix. Here $e_\alpha \in \mathfrak{g}$ are the root vectors normalized to $(e_{-\alpha}, e_\alpha) = \frac{2}{(\alpha, \alpha)}$ with respect to the Killing form. For $\alpha \in \Pi$, they are the classical limits of the quantum Chevalley generators.

2.2 Defining representations of classical matrix groups

By classical Lie algebras (resp. algebraic groups) we mean the simple complex Lie algebras of the types $\mathfrak{g} = sl(n)$, $so(2n+1)$, $sp(n)$, and $so(2n)$, for $n > 1$. We reserve the notation V_0 for the simple (defining) \mathfrak{g} -module of dimension $N = n, 2n+1, 2n$, and $2n$, respectively. We chose the following realization of orthogonal algebras and symplectic algebras. The algebra $so(N)$ leaves invariant the skew-diagonal unit matrix; the symplectic algebra preserves the skew-diagonal matrix with $+1$ above the center and -1 below. We assume that \mathfrak{h} , \mathfrak{n}^+ , and \mathfrak{n}^- are realized in $\text{End}(V_0)$ by, respectively, diagonal, upper- and lower triangular matrices.

Bellow we collect some facts about the defining representation which we will use in our exposition. Let e_{ij} is the standard matrix base in $\text{End}(V_0)$. Define linear functionals $\{\varepsilon_i\}_{i=1}^n$ on \mathfrak{h} setting $\varepsilon_i(e_{ii}) = 1$ for $\mathfrak{g} = sl(n)$ and $\varepsilon_i(e_{ii} - e_{i'i'}) = 1$ otherwise. Here $i' = N + 1 - i$.

The sets of positive and simple roots of \mathfrak{g} are expressed through ε_i by

$$\begin{aligned} R_+ &= \{\varepsilon_i - \varepsilon_j\}_{i < j}, & \Pi &= \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{n-1} - \varepsilon_n\}, & \mathfrak{g} &= sl(n), \\ R_+ &= \{\varepsilon_i \pm \varepsilon_j, \varepsilon_i\}_{i < j}, & \Pi &= \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{n-1} - \varepsilon_n, \varepsilon_n\}, & \mathfrak{g} &= so(2n + 1), \\ R_+ &= \{\varepsilon_i \pm \varepsilon_j, 2\varepsilon_i\}_{i < j}, & \Pi &= \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{n-1} - \varepsilon_n, 2\varepsilon_n\}, & \mathfrak{g} &= sp(n), \\ R_+ &= \{\varepsilon_i \pm \varepsilon_j\}_{i < j}, & \Pi &= \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{n-1} - \varepsilon_n, \varepsilon_{n-1} + \varepsilon_n\}, & \mathfrak{g} &= so(2n). \end{aligned}$$

Our choice of functionals $\{\varepsilon_i\}$ coincides with [VO] excepting $\mathfrak{g} = sl(n)$. In this latter case $\{\varepsilon_i\}$ are linearly dependent and satisfy the condition $\sum_{i=1}^n \varepsilon_i = 0$.

The half-sum of positive roots is expressed in terms of $\{\varepsilon_i\}$ by

$$\rho = \sum_{i=1}^n \rho_i \varepsilon_i, \quad \rho_i = \rho_1 - (i - 1), \quad \rho_1 = \begin{cases} \frac{n-1}{2} & \text{for } \mathfrak{g} = sl(n), \\ n - \frac{1}{2} & \text{for } \mathfrak{g} = so(2n + 1), \\ n & \text{for } \mathfrak{g} = sp(n), \\ n - 1 & \text{for } \mathfrak{g} = so(2n). \end{cases}$$

The set of weights of the defining representation is $\{\varepsilon_i\}_{i=1}^n$ for $sl(n)$, $\{0\} \cup \{\pm\varepsilon_i\}_{i=1}^n$ for $so(2n + 1)$, and $\{\pm\varepsilon_i\}_{i=1}^n$ for $sp(n)$ and $so(2n)$.

2.3 Parabolic subalgebras in $\mathcal{U}_{\hbar}(\mathfrak{g})$

An element $\xi \in \mathfrak{g}$ is called semisimple if $\text{ad}\xi$ is diagonalizable. A semisimple element belongs to a Cartan subalgebra, and all the Cartan subalgebras are conjugated; so one can assume that $\xi \in \mathfrak{h}$.

Let $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ be the triangular decomposition relative to R_+ . A Levi subalgebra in \mathfrak{g} is defined as the centralizer of a semisimple element. It is a reductive Lie algebra of the rank $\text{rk } \mathfrak{g}$. Let \mathfrak{l} be a Levi subalgebra in \mathfrak{g} and $\mathfrak{p}^\pm = \mathfrak{l} + \mathfrak{n}^\pm \subset \mathfrak{g}$ be the parabolic subalgebras. Denote by \mathfrak{n}_\mp^\pm the nilradicals in \mathfrak{p}^\pm . The triangular decomposition $\mathfrak{g} = \mathfrak{n}_\mp^- \oplus \mathfrak{l} \oplus \mathfrak{n}_\mp^+$ induces decomposition $\mathcal{U}(\mathfrak{g}) = \mathcal{U}(\mathfrak{n}_\mp^-)\mathcal{U}(\mathfrak{l})\mathcal{U}(\mathfrak{n}_\mp^+)$, which has a quantum analog, [JT].

The elements $\{e_\alpha, e_{-\alpha}\}_{\alpha \in \Pi_\mathfrak{l}}$ generate over $\mathcal{U}_{\hbar}(\mathfrak{h})$ a Hopf subalgebra $\mathcal{U}_{\hbar}(\mathfrak{l})$ in $\mathcal{U}_{\hbar}(\mathfrak{g})$. This subalgebra is a quantized universal enveloping algebra of the Levi subalgebra $\mathfrak{l} \subset \mathfrak{g}$. It can be represented as $\mathcal{U}_{\hbar}(\mathfrak{l}_0)\mathcal{U}_{\hbar}(\mathfrak{c})$, where $\mathfrak{l}_0 = [\mathfrak{l}, \mathfrak{l}]$ is the semisimple part of \mathfrak{l} and $\mathfrak{c} \subset \mathfrak{l}$ is the center. Also, $\mathcal{U}_{\hbar}(\mathfrak{p}^\pm) := \mathcal{U}_{\hbar}(\mathfrak{l})\mathcal{U}_{\hbar}(\mathfrak{b}^\pm)$ are Hopf subalgebras. They are quantized universal enveloping algebras of the parabolic subalgebras in \mathfrak{g} . This fact follows from the existence of the PBW $\mathcal{U}_{\hbar}(\mathfrak{h})$ -base.

Let \mathbb{Z}_+ denote the set of non-negative integers. Consider in $\mathcal{U}_\hbar(\mathfrak{b}^+)$ the sum of weight spaces with weights from $\mathbb{Z}_+(\Pi_{\mathfrak{g}} - \Pi_{\mathfrak{l}})$. It is an algebra and a deformation of $\mathcal{U}(\mathfrak{h} + \mathfrak{n}_1^+)$, due to the existence of the PBW base in $\mathcal{U}(\mathfrak{h} + \mathfrak{n}_1^+)$. Let us denote this algebra by $\mathcal{U}_\hbar(\mathfrak{h} + \mathfrak{n}_1^+)$. According to [Ke] (see also [JT]), there is a subalgebra in $\mathcal{U}_\hbar(\mathfrak{h} + \mathfrak{n}_1^+)$, denoted further by $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$, such that $\mathcal{U}_\hbar(\mathfrak{b}_1^+)\mathcal{U}_\hbar(\mathfrak{n}_1^+) = \mathcal{U}_\hbar(\mathfrak{b}^+)$. Here $\mathfrak{b}_1^+ = \mathfrak{l} \cap \mathfrak{b}_+$ is the positive Borel subalgebra in \mathfrak{l} . The algebra $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$ is $\mathcal{U}_\hbar(\mathfrak{l})$ -invariant with respect to the adjoint action, and there exists a smash product decomposition

$$\mathcal{U}_\hbar(\mathfrak{p}^+) = \mathcal{U}_\hbar(\mathfrak{l}) \rtimes \mathcal{U}_\hbar(\mathfrak{n}_1^+). \quad (3)$$

Proposition 2.1. *The algebra $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$ is a deformation of $\mathcal{U}(\mathfrak{n}_1^+)$.*

Proof. Decomposition (3) induces the decomposition $\mathcal{U}_\hbar(\mathfrak{h} + \mathfrak{n}_1^+) = \mathcal{U}_\hbar(\mathfrak{h}) \rtimes \mathcal{U}_\hbar(\mathfrak{n}_1^+) \subset \mathcal{U}_\hbar(\mathfrak{p})$. Since $\mathcal{U}_\hbar(\mathfrak{h} + \mathfrak{n}_1^+)$ is a free $\mathcal{U}_\hbar(\mathfrak{h})$ -module generated by the PBW base, $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$ is isomorphic to $\mathcal{U}(\mathfrak{n}_1^+) \otimes \mathbb{C}[[\hbar]]$ as a $\mathbb{C}[[\hbar]]$ -module. By construction, the algebra $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$ is generated by $\text{ad}(u)(e_\alpha)$, where $\alpha \in \Pi_{\mathfrak{g}} - \Pi_{\mathfrak{l}}$ and $u \in \mathcal{U}_\hbar(\mathfrak{l})$. This implies the proposition. \square

Denote by $\mathcal{U}_\hbar(\mathfrak{n}_1^-)$ the ω -image of $\mathcal{U}_\hbar(\mathfrak{n}_1^+)$, where ω is the quantum Chevalley involution. Since $\omega(\mathcal{U}_\hbar(\mathfrak{p}^+)) = \mathcal{U}_\hbar(\mathfrak{p}^-)$ and $\mathcal{U}_\hbar(\mathfrak{l})$ is ω -stable, (3) induces decomposition $\mathcal{U}_\hbar(\mathfrak{p}^-) = \mathcal{U}_\hbar(\mathfrak{n}_1^-) \rtimes \mathcal{U}_\hbar(\mathfrak{l})$, through ω . The algebra $\mathcal{U}_\hbar(\mathfrak{g})$ admits the triangular decomposition

$$\mathcal{U}_\hbar(\mathfrak{g}) = \mathcal{U}_\hbar(\mathfrak{n}_1^-)\mathcal{U}_\hbar(\mathfrak{l})\mathcal{U}_\hbar(\mathfrak{n}_1^+), \quad (4)$$

which is a deformation of the classical one, [JT]. Note that $\mathcal{U}_\hbar(\mathfrak{n}_1^\pm)$ are not Hopf algebras.

Thanks to the triangular decomposition (4), the algebra $\mathcal{U}_\hbar(\mathfrak{g})$ can be represented as a direct sum

$$\mathcal{U}_\hbar(\mathfrak{l}) \oplus \sum_{\alpha \in \Pi_{\mathfrak{g}} - \Pi_{\mathfrak{l}}} (e_{-\alpha}\mathcal{U}_\hbar(\mathfrak{g}) + \mathcal{U}_\hbar(\mathfrak{g})e_\alpha).$$

By $\mathcal{P}_1: \mathcal{U}_\hbar(\mathfrak{g}) \rightarrow \mathcal{U}_\hbar(\mathfrak{l})$ we denote the projection along the second summand.

3 Generalized Verma modules over $\mathcal{U}_\hbar(\mathfrak{g})$

In the present section we study tensor product of finite dimensional and generalized Verma modules over $\mathcal{U}_\hbar(\mathfrak{g})$. This will be the basis for our further considerations.

3.1 Upper and lower (generalized) Verma modules

Let $\mathcal{U}_q(\mathfrak{g})$ be the quantum group in the sense of Lusztig, [L]. It is a $\mathbb{C}(q)$ -Hopf algebra generated by $\{e_{\pm\alpha_i}, q_i^{\pm\hbar\alpha_i}\}_{\alpha_i \in \Pi}$. The algebra $\mathcal{U}_q(\mathfrak{g})$ contains a $\mathbb{C}[q, q^{-1}]$ -Hopf subalgebra $\check{\mathcal{U}}_q(\mathfrak{g})$ such that $\mathcal{U}_q(\mathfrak{g}) \simeq \check{\mathcal{U}}_q(\mathfrak{g}) \otimes_{\mathbb{C}[q, q^{-1}]} \mathbb{C}(q)$. It is generated by $\{e_{\pm\alpha_i}, q_i^{\pm\hbar\alpha_i}, [e_{\alpha_i}, e_{-\alpha_i}]\}_{\alpha_i \in \Pi}$, see e.g. [DCK]. Clearly the algebra $\check{\mathcal{U}}_{\hbar}(\mathfrak{g}) := \check{\mathcal{U}}_q(\mathfrak{g}) \otimes_{\mathbb{C}[q, q^{-1}]} \mathbb{C}[[\hbar]]$, where $\mathbb{C}[q, q^{-1}]$ is embedded in $\mathbb{C}[[\hbar]]$ via $q \mapsto e^{\hbar}$, is dense in $\mathcal{U}_{\hbar}(\mathfrak{g})$ in the \hbar -adic topology. Remark that $\check{\mathcal{U}}_{\hbar}(\mathfrak{g})$ is $\mathcal{U}_{\hbar}(\mathfrak{g})$ -invariant with respect to the adjoint action.

There is a one-to-one correspondence between finite-dimensional \mathfrak{g} -modules and finite-dimensional $\mathcal{U}_q(\mathfrak{g})$ -modules with $q^{\mathbb{Z}}$ -valued weights. Each such module is isomorphic to $\check{W} \otimes_{\mathbb{C}[q, q^{-1}]} \mathbb{C}(q)$, where \check{W} is a $\check{\mathcal{U}}_q(\mathfrak{g})$ -module, free and finite over $\mathbb{C}[q, q^{-1}]$, [Jan2]. The specialization $\check{W} \bmod (q-1)$ gives a finite dimensional \mathfrak{g} -module. Therefore \check{W} extends to a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module, free and finite over $\mathbb{C}[[\hbar]]$. We will call such $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules **finite dimensional**. They are deformations of \mathfrak{g} -modules, diagonalizable over $\mathcal{U}_{\hbar}(\mathfrak{h})$, and have the same weight structure. This correspondence between finite dimensional \mathfrak{g} -modules and $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules is additive. That is, every finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module is a direct sum of "simple" modules, i. e. those whose classical limit is simple.

Every submodule of a finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module is \mathfrak{h} -diagonalizable with $q^{\mathbb{Z}}$ -valued weights; hence it is again finite dimensional. We will also deal with $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules that are finitely generated over $\mathbb{C}[[\hbar]]$ but not free. We always assume that such modules are quotients of finite dimensional and called them just $\mathbb{C}[[\hbar]]$ -finite.

A highest weight $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module is generated by a weight vector annihilated by $\mathcal{U}'_{\hbar}(\mathfrak{n}^+)$. Similarly, a lowest weight module is generated by a weight vector annihilated by $\mathcal{U}'_{\hbar}(\mathfrak{n}^-)$. Finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules have highest and lowest weights simultaneously. Their highest weights are integral dominant. Finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules with highest weights are deformations of irreducible finite dimensional \mathfrak{g} -module. They are substitutes for irreducibles, since a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module is almost never irreducible in the usual sense (multiplication by \hbar is a morphism).

For reductive \mathfrak{l} , the highest weight $\mathcal{U}_{\hbar}(\mathfrak{l})$ -modules are defined similarly to the semisimple case. Let A be a representation of $\mathcal{U}_{\hbar}(\mathfrak{l})$ with highest weight. It extends to a representation of $\mathcal{U}_{\hbar}(\mathfrak{p})$ for $\mathfrak{p} = \mathfrak{p}^+$, by setting it zero on $\mathcal{U}'_{\hbar}(\mathfrak{n}_l^+)$. A generalized Verma module over $\mathcal{U}_{\hbar}(\mathfrak{g})$ is the induced module $M_{\mathfrak{p}, A} = \mathcal{U}_{\hbar}(\mathfrak{g}) \otimes_{\mathcal{U}_{\hbar}(\mathfrak{p})} A =: \text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} A$, cf. [J1]. By \mathbb{C}_{λ} we will denote the one dimensional $\mathcal{U}_{\hbar}(\mathfrak{l})$ -module defined by a character $\lambda: \mathfrak{l} \rightarrow \mathbb{C}$. The module $M_{\mathfrak{p}, \mathbb{C}_{\lambda}}$ will be denoted simply by $M_{\mathfrak{p}, \lambda}$ and the Verma module $M_{\mathfrak{b}, \lambda}$ by M_{λ} .

Every $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module with highest weight λ is a quotient of M_{λ} . The generalized Verma

module $M_{\mathfrak{p},A}$ induced from a $\mathcal{U}_{\hbar}(\mathfrak{l})$ -module A with highest weight λ is a highest weight module, so it is a quotient of M_λ as well.

Lemma 3.1. *Let A be a finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{l})$ -module. Then the $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module $M_{\mathfrak{p},A}$ is a deformation of the classical generalized Verma module over $\mathcal{U}(\mathfrak{g})$.*

Proof. It follows from (4) that $M_{\mathfrak{p},A}$ is a free $\mathcal{U}_{\hbar}(\mathfrak{n}_1^-)$ -module generated by $1 \otimes_{\mathcal{U}_{\hbar}(\mathfrak{p})} A$ and hence it is $\mathbb{C}[[\hbar]]$ -free. Since the decomposition (4) is a deformation of the classical triangular decomposition, $M_{\mathfrak{p},A}$ is a deformation of the corresponding generalized Verma module over $\mathcal{U}(\mathfrak{g})$. \square

A highest weight representation of $\mathcal{U}_{\hbar}(\mathfrak{l})$ can be extended to a representation of $\mathcal{U}_{\hbar}(\mathfrak{p}^-)$ by setting it trivial on $\mathcal{U}_{\hbar}(\mathfrak{n}_1^-)$. This is possible, due to (3). Similarly to $M_{\mathfrak{p}^+,A}$, the module $M_{\mathfrak{p}^-,A} := \mathcal{U}_{\hbar}(\mathfrak{g}) \otimes_{\mathcal{U}_{\hbar}(\mathfrak{p}^-)} A$ is introduced, where A is taken to be a lowest weight $\mathcal{U}_{\hbar}(\mathfrak{l})$ -module.

Proposition 3.2. *Let A be a lowest and B a highest weight $\mathcal{U}_{\hbar}(\mathfrak{l})$ -modules. Then the $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module $M_{\mathfrak{p}^-,A} \otimes M_{\mathfrak{p}^+,B}$ is isomorphic to $\text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(A \otimes B) := \mathcal{U}_{\hbar}(\mathfrak{g}) \otimes_{\mathcal{U}_{\hbar}(\mathfrak{l})} (A \otimes B)$.*

Proof. Consider the map $\mathcal{U}_{\hbar}(\mathfrak{g}) \otimes A \otimes B \mapsto (\mathcal{U}_{\hbar}(\mathfrak{g}) \otimes A) \otimes (\mathcal{U}_{\hbar}(\mathfrak{g}) \otimes B)$ defined by $u \otimes a \otimes b \mapsto u^{(1)} \otimes a \otimes u^{(2)} \otimes b$, where $u^{(1)} \otimes u^{(2)}$ is the standard symbolic notation for the coproduct $\Delta(u)$. This map induces a homomorphism

$$\Delta_{\mathfrak{l},A,B}^{\mathfrak{g}}: \mathcal{U}_{\hbar}(\mathfrak{g}) \otimes_{\mathcal{U}_{\hbar}(\mathfrak{l})} (A \otimes B) \rightarrow M_{\mathfrak{p}^-,A} \otimes M_{\mathfrak{p}^+,B} \quad (5)$$

of $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules. We claim that this map is an isomorphism.

First let us prove the statement assuming $\mathfrak{l} = \mathfrak{h}$, $A = \mathbb{C}_{\mu}$, and $B = \mathbb{C}_{\nu}$. Introduce the grading in $\mathcal{U}_{\hbar}(\mathfrak{n}^{\pm})$ by weight height setting $\deg e_{\pm\alpha} = 1$ for $\alpha \in \Pi$. The gradings in $\mathcal{U}_{\hbar}(\mathfrak{n}^{\pm})$ induce a double grading in $\text{Ind}_{\mathfrak{h}}^{\mathfrak{g}}(\mathbb{C}_{\mu} \otimes \mathbb{C}_{\nu})$ and $M_{\mathfrak{p}^-, \mu} \otimes M_{\mathfrak{p}^+, \nu}$, which can be identified as graded spaces. Each homogeneous component has finite rank over $\mathbb{C}[[\hbar]]$. Let v_{μ} and v_{ν} be the generators of $M_{\mathfrak{p}^-, \mu}$ and $M_{\mathfrak{p}^+, \nu}$, respectively. Take $u^{\pm} \in \mathcal{U}_{\hbar}(\mathfrak{n}^{\pm})$ to be monomials in $\{e_{\pm\alpha}\}_{\alpha \in \Pi}$ and compute the map $\Delta_{\mathfrak{l}, \mathbb{C}_{\mu}, \mathbb{C}_{\nu}}^{\mathfrak{g}}$:

$$\begin{aligned} u^- u^+(v_{\mu} \otimes v_{\nu}) &\mapsto (u^-)^{(1)}(u^+)^{(1)}v_{\mu} \otimes (u^-)^{(2)}(u^+)^{(2)}v_{\nu} = (u^-)^{(1)}u^+v_{\mu} \otimes (u^-)^{(2)}v_{\nu} \\ &= c_q u^+v_{\mu} \otimes u^-v_{\nu} + w^+v_{\mu} \otimes w^-v_{\nu}. \end{aligned}$$

Here $c_q \in \mathbb{C}[[\hbar]]$ is invertible and the elements $w^{\pm} \in \mathcal{U}_{\hbar}(\mathfrak{n}^{\pm})$ belong to subspaces of degree $< \deg u^{\pm}$. This computation shows that $\Delta_{\mathfrak{l}, \mathbb{C}_{\mu}, \mathbb{C}_{\nu}}^{\mathfrak{g}}$ is a triangular operator (relative to the double grading) with invertible diagonal. Therefore it is an isomorphism.

The above consideration also proves that the map (5) is an epimorphism in the general situation, as A and B are quotients of $M_{p^-, \mu}$ and $M_{p^+, \nu}$, respectively. We must check that (5) is injective. The map $\Delta_{\mathfrak{l}, A, B}^{\mathfrak{g}}$ is surjective modulo \hbar as a $\mathcal{U}(\mathfrak{g})$ -morphism. By dimensional arguments based on the bi-grading, we conclude that this $\mathcal{U}(\mathfrak{g})$ -morphism is an isomorphism. Therefore $\Delta_{\mathfrak{l}, A, B}^{\mathfrak{g}}$ is an isomorphism by the obvious deformation arguments. \square

Remark 3.3. In conclusion of this section we remark that the triangular decomposition and the generalized Verma modules can be naturally defined for the algebra $\check{\mathcal{U}}_{\hbar}(\mathfrak{g})$. We will use this observation in Section 8, where the induction is made from the character $\lambda/2\hbar$, $\lambda \in \mathfrak{c}^*$. The $\check{\mathcal{U}}_{\hbar}(\mathfrak{g})$ -action on $M_{\mathfrak{p}, \lambda/2\hbar}$ does not extend to an action of $\mathcal{U}_{\hbar}(\mathfrak{g})$. However the $\check{\mathcal{U}}_{\hbar}(\mathfrak{g})$ -action on $\text{End}(M_{\mathfrak{p}, \lambda/2\hbar})$ extends to an action of $\mathcal{U}_{\hbar}(\mathfrak{h})$, and that is what we need for our construction.

3.2 Pairing between upper and lower generalized Verma modules

Denote by \mathfrak{c} the center of \mathfrak{l} and by $\mathfrak{c}_{reg} \subset \mathfrak{c}$ the subset of elements whose centralizer is exactly \mathfrak{l} . Clearly \mathfrak{c}_{reg} is a dense open set in \mathfrak{c} . The coadjoint \mathfrak{g} -module \mathfrak{g}^* is canonically identified with \mathfrak{g} via the Killing form. Then the dual space \mathfrak{c}^* is identified with the orthogonal complement to the annihilator of \mathfrak{c} in \mathfrak{h}^* , so $\mathfrak{c}^* \subset \mathfrak{h}^*$ under this convention.

Given a root $\alpha \in R$, let α^\vee denote the dual root $\alpha^\vee = \frac{2}{(\alpha, \alpha)}\alpha$. We call a weight $\lambda \in \mathfrak{c}^*$ generic if $(\lambda, \alpha^\vee) \notin \mathbb{Z}$ for all $\alpha \in \Pi_{\mathfrak{g}} - \Pi_{\mathfrak{l}}$. Clearly the set \mathfrak{c}_{gen}^* of generic weights is a dense open subset in $\mathfrak{c}_{reg}^* = \{\lambda \in \mathfrak{c}^* \mid (\lambda, \alpha^\vee) \neq 0, \forall \alpha \in \Pi_{\mathfrak{g}} - \Pi_{\mathfrak{l}}\}$.

With every finite dimensional irreducible \mathfrak{l} -module A one can associate a weight $\lambda_A \in \mathfrak{c}^*$ such that $ha = \lambda_A(h)a$ for all $h \in \mathfrak{c}$ and $a \in A$. We call A generic if $\lambda_A \in \mathfrak{c}_{gen}^*$. An arbitrary finite dimensional \mathfrak{l} -module is called generic if its every irreducible submodule is generic. This terminology extends to the corresponding $\mathcal{U}_{\hbar}(\mathfrak{l})$ -modules.

There exists a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -equivariant pairing between M_{p^-, A^*} and $M_{p^+, A}$. The construction goes as follows. Consider a bilinear $\mathcal{U}_{\hbar}(\mathfrak{g})$ -equivariant map $(\mathcal{U}_{\hbar}(\mathfrak{g}) \otimes A^*) \otimes (\mathcal{U}_{\hbar}(\mathfrak{g}) \otimes A) \rightarrow \mathbb{C}[[\hbar]]$ defined by $u_1 \otimes \xi \otimes u_2 \otimes x \mapsto \xi(\mathcal{P}_{\mathfrak{l}}(\gamma(u_1)u_2)x)$, where $\mathcal{P}_{\mathfrak{l}}$ is introduced in Subsection 2.3. This map is equivariant by construction and factors through a bilinear equivariant map $M_{p^-, A^*} \otimes M_{p^+, A} \rightarrow \mathbb{C}[[\hbar]]$, as required.

Proposition 3.4. *Let A be a generic finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{l})$ -module. Then the equivariant pairing between $M_{p^+, A}$ and M_{p^-, A^*} is nondegenerate.*

Proof. Without loss of generality, we may assume that $A_0 = A/\hbar A$ is irreducible.

Since $-\mathfrak{c}_{gen}^* = \mathfrak{c}_{gen}^*$, the modules A and A^* are generic simultaneously. It follows from [Jan1], Satz 3, that the $\mathcal{U}(\mathfrak{l})$ -module $M_{\mathfrak{p}^+, A}/\hbar M_{\mathfrak{p}^+, A}$ is irreducible for $\lambda \in \mathfrak{c}_{gen}^*$. Clearly the same is true for $M_{\mathfrak{p}^-, A^*}/\hbar M_{\mathfrak{p}^-, A^*}$. Therefore the pairing in question is nondegenerate modulo \hbar (being $\mathcal{U}(\mathfrak{l})$ -equivariant and not identically zero).

With respect to the pairing, the weight spaces of weights μ and ν are orthogonal unless $\mu + \nu = 0$. Since the weight spaces in $M_{\mathfrak{p}^+, A}$ and $M_{\mathfrak{p}^-, A^*}$ are $\mathbb{C}[[\hbar]]$ -finite, non-degeneracy of the pairing follows from non-degeneracy modulo \hbar . \square

3.3 Tensor product of finite dimensional and generalized Verma modules

We call a $\mathcal{U}_\hbar(\mathfrak{g})$ -module a weight module if it is \mathfrak{h} -diagonalizable and its weight spaces are finite and free over $\mathbb{C}[[\hbar]]$.

For any weight module let $\Lambda'(U)$ denote the set of weights of U and $U[\mu]$ the weight space for $\mu \in \Lambda'(U)$. The dual module U^* of linear functionals on U consists of infinite formal sums $f = \sum_{\mu} f_{\mu}$, where $f_{\mu} \in U[\mu]^*$. The action of $\mathcal{U}_\hbar(\mathfrak{g})$ on U^* is defined to be $(xf)(u) = f(\gamma(x)u)$, for $f \in U^*$, $u \in U$, and $x \in \mathcal{U}_\hbar(\mathfrak{g})$. Define the restricted dual U° as a natural $\mathcal{U}_\hbar(\mathfrak{g})$ -submodule in U^* by setting $U^\circ = \bigoplus_{\mu \in \Lambda'(U)} U[\mu]^* \subset U^*$ (only finite sums admitted).

Lemma 3.5. *Let U_1, U_2 be weight $\mathcal{U}_\hbar(\mathfrak{g})$ -modules and W_1, W_2 finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -modules. Then $\text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(W_1 \otimes U_1, W_2 \otimes U_2) \simeq \text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(U_2^\circ \otimes U_1, W_2 \otimes W_1^*)$.*

Proof. Clear. \square

Proposition 3.4 asserts that $M_{\mathfrak{p}^-, A^*} \simeq M_{\mathfrak{p}, A}^\circ$ for generic A .

Lemma 3.6. *Let W be a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module, A and B finite dimensional $\mathcal{U}_\hbar(\mathfrak{l})$ -modules, and $\lambda \in \mathfrak{c}_{gen}^*$. Then the following $\mathbb{C}[[\hbar]]$ -linear isomorphisms take place:*

$$\begin{aligned}
\text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}, M_{\mathfrak{p}, B \otimes \mathbb{C}_\lambda}) &\simeq \text{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(A, B), \\
\text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}, W \otimes M_{\mathfrak{p}, \lambda}) &\simeq \text{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(A, W), \\
\text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(W \otimes M_{\mathfrak{p}, \lambda}, M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}) &\simeq \text{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(W, A), \\
\text{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(W \otimes M_{\mathfrak{p}, \lambda}, W \otimes M_{\mathfrak{p}, \lambda}) &\simeq \text{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(W, W).
\end{aligned} \tag{6}$$

Proof. The proof is based on Propositions 3.2 and 3.4 and can be conducted similarly as in [DM1] for the classical case of $\mathcal{U}(\mathfrak{g})$ -modules. For instance, let us check the first isomorphism.

By Proposition 3.4, the module $M_{\mathfrak{p}^-, \mathbb{C}_\lambda^* \otimes B^*}$ is isomorphic to the restricted dual $M_{\mathfrak{p}, B \otimes \mathbb{C}_\lambda}^\circ$. Therefore

$$\mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}, M_{\mathfrak{p}, B \otimes \mathbb{C}_\lambda}) \simeq \mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{g})}(M_{\mathfrak{p}^-, \mathbb{C}_\lambda^* \otimes B^*} \otimes M_{\mathfrak{p}^+, A \otimes \mathbb{C}_\lambda}, \mathbb{C}[[\hbar]]), \quad (7)$$

by Lemma 3.5. Here $\mathbb{C}[[\hbar]]$ is the trivial $\mathcal{U}_\hbar(\mathfrak{g})$ -module. According to Lemma 3.2, the tensor product of lower and upper generalized Verma modules is induced from the $\mathcal{U}_\hbar(\mathfrak{l})$ -module $\mathbb{C}_\lambda^* \otimes B^* \otimes A \otimes \mathbb{C}_\lambda \simeq B^* \otimes A$. Applying the Frobenius reciprocity, we continue (7) with $\mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(B^* \otimes A, \mathbb{C}[[\hbar]]) \simeq \mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(A, B)$, as required. \square

Proposition 3.7. *Let W be a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module and $\lambda \in \mathfrak{c}_{gen}^*$. Then $W \otimes M_{\mathfrak{p}, \lambda}$ admits the direct sum decomposition*

$$W \otimes M_{\mathfrak{p}, \lambda} = \bigoplus_{A_0} M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda}, \quad (8)$$

where summation is taken over the simple \mathfrak{l} -modules with multiplicities entering W_0 .

Proof. The isomorphisms (6) hold modulo \hbar . Moreover, they commute with taking quotients mod \hbar . First let us prove the classical mod \hbar analog of decomposition (8), retaining the same notation for the Verma modules over $\mathcal{U}(\mathfrak{g})$. We assume in (8) a fixed decomposition within each isotypic A_0 -component of W_0 . Let $j^{A_0}: A_0 \hookrightarrow W_0$ be the \mathfrak{l} -equivariant injections such that $\sum_A j^{A_0} = \mathrm{id}_{W_0}$. Let \hat{j}^{A_0} be their lifts $M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda} \rightarrow W \otimes M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda}$. For generic $\lambda \in \mathfrak{c}_{gen}^*$ all the modules $M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda}$ are irreducible. Therefore all \hat{j}^{A_0} are linearly independent, and the \mathfrak{g} -equivariant map $\bigoplus_{A_0} M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda} \xrightarrow{\sum_{A_0} \hat{j}^{A_0}} W \otimes M_{\mathfrak{p}, \lambda}$ is an embedding. Then it is an embedding of \mathfrak{h} -modules $\mathcal{U}(\mathfrak{n}_1^-) \otimes W_0 \simeq \bigoplus_{A_0} \mathcal{U}(\mathfrak{n}_1^-) \otimes A_0 \xrightarrow{\sum_{A_0} \hat{j}^{A_0}} W \otimes \mathcal{U}(\mathfrak{n}_1^-)$ and therefore an isomorphism. This proves the statement modulo \hbar .

We can chose $j^A \in \mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(A, W)$ to be deformations of morphisms j^{A_0} splitting the $\mathcal{U}_\hbar(\mathfrak{l})$ -module W into the direct sum of highest weight submodules. Take the $\mathcal{U}_\hbar(\mathfrak{g})$ -morphisms $\hat{j}^A: M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda} \rightarrow W \otimes M_{\mathfrak{p}, \lambda}$ corresponding to $j^A \in \mathrm{Hom}_{\mathcal{U}_\hbar(\mathfrak{l})}(A, W)$ under the second isomorphism from (6). Since isomorphisms (6) commute with taking quotients mod \hbar , we have $\hat{j}^A = \hat{j}^{A_0} \pmod{\hbar}$. Consider the morphism $\bigoplus_{A_0} M_{\mathfrak{p}, A_0 \otimes \mathbb{C}_\lambda} \xrightarrow{\sum_{A_0} \hat{j}^A} W \otimes M_{\mathfrak{p}, \lambda}$ of $\mathcal{U}_\hbar(\mathfrak{g})$ -modules. Restricting consideration to weight spaces we conclude that $\sum_{A_0} \hat{j}^A$ is an isomorphism because it is so modulo \hbar . \square

4 Properties of the universal RE matrix

In the present section we recall general properties of the universal reflection equation (RE) matrix $\mathcal{Q} = \mathcal{R}_{21} \mathcal{R}$ and study its action on tensor products of finite dimensional and (gener-

alized) Verma modules.

Recall from [Dr2] that the element $\mathcal{Q} = \mathcal{R}_{21}\mathcal{R}$ can be represented as $\mathcal{Q} = \Delta(v)(v^{-1} \otimes v^{-1})$, where $v = \gamma(\mathcal{R}_1^{-1})\mathcal{R}_2^{-1} = \gamma^2(\mathcal{R}_1)\mathcal{R}_2$. Conjugation with v implements the squared antipode $v x v^{-1} = \gamma^2(x)$ for all $x \in \mathcal{U}_{\hbar}(\mathfrak{g})$. On the other hand, the squared antipode can be written as the conjugation $\gamma^2(x) = q^{-2h_\rho} x q^{2h_\rho}$, $x \in \mathcal{U}_{\hbar}(\mathfrak{g})$. It follows from here that $v = q^{-2h_\rho} z$, where z is some invertible element from the center of $\mathcal{U}_{\hbar}(\mathfrak{g})$. Therefore we can write

$$\mathcal{Q} = \Delta(z)(z^{-1} \otimes z^{-1}), \quad (9)$$

as q^{-2h_ρ} is a group-like element. Let $\lambda \in \mathfrak{h}^*$ and χ^λ be the corresponding central character of $\mathcal{U}_{\hbar}(\mathfrak{g})$. It is easy to compute the value $\chi^\lambda(z)$ via a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module W with highest weight, using the structure of R-matrix (1). Let w_λ be the highest weight vector in W . Since $\mathcal{U}'_{\hbar}(\mathfrak{n}^+)w_\lambda = 0$, we have $v w_\lambda = q^{(\lambda, \lambda)} w_\lambda = \chi^\lambda(z) q^{-2(\rho, \lambda)} w_\lambda$. From this we find

$$\chi^\lambda(z) = q^{(\lambda, \lambda) + 2(\rho, \lambda)}. \quad (10)$$

The element \mathcal{Q} satisfies the identities

$$\mathcal{R}_{21} \mathcal{Q}_{13} \mathcal{R}_{12} \mathcal{Q}_{23} = \mathcal{Q}_{23} \mathcal{R}_{21} \mathcal{Q}_{13} \mathcal{R}_{12}, \quad (11)$$

$$(\Delta \otimes \text{id})(\mathcal{Q}) = \mathcal{R}_{12}^{-1} \mathcal{Q}_{13} \mathcal{R}_{12} \mathcal{Q}_{23}, \quad (12)$$

of which the first may be called the universal reflection equation, cf. [DKM]. Equation (12) is the key identity of the fusion procedure for solutions to the RE, [DKM].

4.1 Characteristic polynomial for RE matrix

Given a finite dimensional representation (W, π_W) let \mathcal{Q}_W denote the element $(\pi_W \otimes \text{id})(\mathcal{Q}) \in \text{End}(W) \otimes \mathcal{U}_{\hbar}(\mathfrak{g})$.

Proposition 4.1. *Let W be a highest weight $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module with the multiset of weights $\Lambda(W)$. Then i) there exists a polynomial p of degree $\#\Lambda(W)$ in one variable with coefficients in the center of $\mathcal{U}_{\hbar}(\mathfrak{g})$ such that $p(\mathcal{Q}_W) = 0$, ii) the spectrum of the operator \mathcal{Q}_W on $W \otimes M_\lambda$ is*

$$\left\{ q^{2(\lambda + \rho, \nu_i) - 2(\rho, \nu) + (\nu_i, \nu_i) - (\nu, \nu)} \right\}_{\nu_i \in \Lambda(W)}, \quad (13)$$

where ν is the highest weight of W .

Proof. Notice that a symmetric function in the eigenvalues (13) is invariant under the action of the Weyl group. Then i) follows from ii) through the Harish-Chandra homomorphism, [Jan2]. So let us check ii).

As a $\mathbb{C}[[\hbar]]$ -module, $W \otimes M_\lambda$ is isomorphic to $W \otimes \mathcal{U}_\hbar(\mathfrak{n}^-)$. Under this isomorphism, the subspace of weight $\beta \in \mathfrak{h}^*$ in $W \otimes \mathcal{U}_\hbar(\mathfrak{n}^-)$ goes to the subspace of weight $\beta + \lambda$ in $W \otimes M_\lambda$. Since \mathcal{Q} is invariant, it preserves the weight spaces in $W \otimes M_\lambda$, which have finite rank over $\mathbb{C}[[\hbar]]$. Now observe that for generic λ ii) follows from Proposition 3.7. This implies ii) for arbitrary λ , since restriction of \mathcal{Q}_W to any weight space has polynomial dependence on λ in every order in \hbar . \square

4.2 Minimal polynomial for RE matrix

In this subsection we determine the spectrum of \mathcal{Q} on the tensor product of finite dimensional and generalized Verma modules over $\mathcal{U}_\hbar(\mathfrak{g})$.

Theorem 4.2. *Let $\lambda \in \mathfrak{c}_{gen}^*$ be a generic character of a Levi subalgebra in $\mathfrak{l} \subset \mathfrak{g}$. Let W be a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module and let $\Lambda_{\mathfrak{l}}(W) = \{v_l = \nu_{i_l}\} \subset \Lambda(W)$ denote the multiset of highest weights of simple \mathfrak{l} -modules entering W_0 . The operator \mathcal{Q} is diagonalizable on $W \otimes M_{\mathfrak{p},\lambda}$ and has eigenvalues*

$$\left\{ q^{2(\lambda + \rho, v_l) - 2(\rho, \nu) + (v_l, v_l) - (\nu, \nu)} \right\}_{v_l \in \Lambda_{\mathfrak{l}}(W)}. \quad (14)$$

Proof. For generic λ the module $W \otimes M_{\mathfrak{p},\lambda}$ splits into the direct sum of highest weight modules, by Proposition 3.7. The operator \mathcal{Q} is proportional to $\Delta(z)$ on $W \otimes M_{\mathfrak{p},\lambda}$. Thus we conclude that \mathcal{Q} is diagonalizable because $\Delta(z)$ is diagonalizable.

For every $\mathcal{U}_\hbar(\mathfrak{l})$ -module $A \subset W$ with highest weight μ the summand $M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}$ in (8) is a $\mathcal{U}_\hbar(\mathfrak{g})$ -module of highest weight $\lambda + \mu$. Hence $\Delta(z)$ and therefore \mathcal{Q} act as scalar multipliers on $M_{\mathfrak{p}, A \otimes \mathbb{C}_\lambda}$. Their eigenvalues are computed using (10). \square

4.3 A construction of central elements

Assume that \mathfrak{g} is an arbitrary complex simple Lie algebra. Let (W, π_W) be a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module. Let $X \in \text{End}(W) \otimes \mathcal{U}_\hbar(\mathfrak{g})$ be an invariant matrix, i. e. commuting with $\pi_W(x^{(1)}) \otimes x^{(2)}$ for all $x \in \mathcal{U}_\hbar(\mathfrak{g})$. It is known that the q-trace defined by

$$\text{Tr}_q(X) := \text{Tr}(\pi_W(q^{2h\rho})X)$$

is ad-invariant and hence belongs to the center of $\mathcal{U}_\hbar(\mathfrak{g})$.

The annihilator of the Verma module with the highest weight λ is generated by the kernel of a central character χ^λ . Let us compute the values $\chi^\lambda(\tau^\ell)$.

Define a map $d: \mathfrak{h}^* \rightarrow \mathbb{C}[[\hbar]]$ setting

$$d(\lambda) := \prod_{\alpha \in \mathbb{R}_+} \frac{q^{(\lambda+\rho, \alpha)} - q^{-(\lambda+\rho, \alpha)}}{q^{(\rho, \alpha)} - q^{-(\rho, \alpha)}}, \quad q = e^\hbar. \quad (15)$$

For a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module W with the highest weight λ , the Weyl character formula [N] gives $d(\lambda) = \text{Tr}_q(\text{id}_W)$, the q -dimension of W .

Proposition 4.3. *Let W be a finite dimensional $\mathcal{U}_\hbar(\mathfrak{g})$ -module with the multiset of weights $\Lambda(W)$. Then for any $\lambda \in \mathfrak{h}^*$*

$$\chi^\lambda\left(\text{Tr}_q(\mathcal{Q}_W^\ell)\right) = \sum_{\nu_i \in \Lambda(W)} x_{\nu_i}^\ell \frac{d(\lambda + \nu_i)}{d(\lambda)}, \quad (16)$$

where x_{ν_i} are the eigenvalues of \mathcal{Q}_W given by (13).

Proof. We adapt a proof from [GZB], because it is suitable for any $\mathcal{U}_\hbar(\mathfrak{g})$ -invariant operator.

Let us check the statement for special λ first. Namely, we suppose that λ is integral dominant and ν , the highest weight of W , is subordinate to λ . The weight ν is called subordinate to λ if $\lambda + \nu_i$ are dominant integral for all $\nu_i \in \Lambda(W)$. For ν fixed we denote by \mathfrak{D}_ν^+ the set of such λ that ν is subordinate to λ . According to [K], a polynomial function on \mathfrak{h}^* is determined by its values on \mathfrak{D}_ν^+ . Both sides of (16) are polynomials in λ in every order in \hbar , thus it suffices to compare them on \mathfrak{D}_ν^+ only.

For $\lambda \in \mathfrak{D}_\nu^+$ let us compute the central character in question on the finite dimensional module U with the highest weight λ . In this case, the module $W \otimes U$ splits into the direct sum of $\mathcal{U}_\hbar(\mathfrak{g})$ -modules of highest weights $\nu_i + \lambda$ for $\nu_i \in \Lambda(W)$. An invariant operator on $W \otimes U$ decomposes into over invariant projectors $\{P_i\}_{\nu_i \in \Lambda(W)}$ to the highest weight submodules, so it suffices to compute the q -trace of these projectors. The operator $(\text{Tr}_q \otimes \text{id})(P_i)$ is constant on U . Taking q -trace over U gives $(\text{Tr}_q \otimes \text{id})(P_i) = (\text{Tr}_q \otimes \text{Tr}_q)(P_i) / (\text{id} \otimes \text{Tr}_q)(P_i) = d(\lambda + \nu_i) / d(\lambda)$, since the q -trace is multiplicative with respect to the tensor product. This proves the statement for $\lambda \in \mathfrak{D}_\nu^+$ and therefore for all $\lambda \in \mathfrak{h}^*$. \square

5 On quantization of affine algebraic varieties

In this section we develop a machinery for equivariant quantization. Throughout the section we assume that \mathfrak{g} is semisimple. Moreover, here we admit an *arbitrary*, i. e. triangular or even trivial, quantization of $\mathcal{U}(\mathfrak{g})$.

In what follows, we use some standard facts from commutative algebra, such as the Nakayama lemma. The reader can find the "list" of facts we rely on in [M].

5.1 A flatness criterion

Recall that an associative algebra and (left) $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module \mathcal{A} is called a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module algebra if its multiplication is compatible with the $\mathcal{U}_{\hbar}(\mathfrak{g})$ -action. That is, for all $h \in \mathcal{U}_{\hbar}(\mathfrak{g})$ and all $a, b \in \mathcal{A}$

$$h \triangleright (ab) = (h^{(1)} \triangleright a)(h^{(2)} \triangleright b), \quad \text{where} \quad h^{(1)} \otimes h^{(2)} = \Delta(h).$$

Definition 5.1. A deformation \mathcal{A} of a $\mathcal{U}(\mathfrak{g})$ -module algebra \mathcal{A}_0 is called equivariant if \mathcal{A} is a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module algebra and the action of $\mathcal{U}_{\hbar}(\mathfrak{g})$ on \mathcal{A} coincides modulo \hbar with the action of $\mathcal{U}(\mathfrak{g})$ on \mathcal{A}_0 .

For every $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module E and a finite dimensional highest weight $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module W there exists a natural morphism $W \otimes \text{Hom}_{\mathcal{U}_{\hbar}(\mathfrak{g})}(W, E) \rightarrow E$ of $\mathcal{U}_{\hbar}(\mathfrak{g})$ -modules. We call the image of this morphism the isotypic W -component of E .

We call a $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module **admissible** if it is a direct sum of its isotypic components and each of them is finitely generated over $\mathbb{C}[[\hbar]]$. It can be shown that submodules and quotient modules of an admissible module are admissible.

The following proposition gives a flatness criterion for quotient algebras. Informally, if one constructs an ideal \mathfrak{J} that lies in the kernel of a certain homomorphism $\mathfrak{S} \rightarrow \mathfrak{T}$ and has the "right classical limit", then \mathfrak{J} equals the entire kernel.

Proposition 5.2 (deformation method). *Let \mathfrak{S} be an admissible and \mathfrak{T} torsion free $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module algebras; let $\Psi: \mathfrak{S} \rightarrow \mathfrak{T}$ be a non-zero equivariant homomorphism. Suppose $\ker \Psi$ contains an invariant ideal \mathfrak{J} such that the image \mathfrak{J}_0^b of \mathfrak{J} in \mathfrak{S}_0 is a maximal \mathfrak{g} -invariant ideal. Then i) $\text{im } \Psi$ is a $\mathbb{C}[[\hbar]]$ -free deformation of $\mathfrak{S}_0/\mathfrak{J}_0^b$, ii) $\ker \Psi = \mathfrak{J}$.*

Proof. The $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module $\text{im } \Psi$ is admissible and torsion free; hence it is $\mathbb{C}[[\hbar]]$ -free.

Since $\text{im } \Psi$ is free, we have the direct sum decomposition $\mathfrak{S} \simeq \ker \Psi \oplus \text{im } \Psi$ of $\mathbb{C}[[\hbar]]$ -modules. Hence $\mathfrak{S}_0 \simeq (\ker \Psi)_0 \oplus (\text{im } \Psi)_0$ and $(\ker \Psi)_0 \supset \mathfrak{J}_0^b$. By assumption, \mathfrak{J}_0^b is a maximal \mathfrak{g} -invariant ideal, hence either $(\ker \Psi)_0 = \mathfrak{S}_0$ or $(\ker \Psi)_0 = \mathfrak{J}_0^b$. The first option is impossible. Indeed, then $(\text{im } \Psi)_0 = 0$ and $\text{im } \Psi = 0$ since $\text{im } \Psi$ is $\mathbb{C}[[\hbar]]$ -free. In this case $\ker \Psi = \mathfrak{S}$, and the map Ψ would have been zero. Therefore $(\text{im } \Psi)_0 = \mathfrak{S}_0/\mathfrak{J}_0^b$ and i) is proven.

We have an embedding $\mathfrak{J} \hookrightarrow \ker \Psi$ and an epimorphism $\mathfrak{J}_0 \rightarrow (\ker \Psi)_0 = \mathfrak{J}_0^b$. Applying the Nakayama lemma to each isotypic component, we prove ii). \square

Remark 5.3. 1. We emphasize that \mathfrak{J}_0^b is assumed to be not just $\mathfrak{J}_0 = \mathfrak{J}/\hbar\mathfrak{J}$ but its image in \mathfrak{S}_0 . It is essential to distinguish between \mathfrak{J}_0 and \mathfrak{J}_0^b because the functor $\text{mod } \hbar$ is not left exact. Eventually, \mathfrak{J}_0 and \mathfrak{J}_0^b coincide in our situation. However that is not *a priori* obvious and follows from the proof. In practice, \mathfrak{J} is often defined via a system of generators. Then \mathfrak{J}_0^b is generated by their images in \mathfrak{S}_0 , so it is even easier to control \mathfrak{J}_0^b than \mathfrak{J}_0 .

6 Quantization of simple algebraic groups

In the present section, we construct quantization of a special Poisson bracket on a simple algebraic group. The quantized ring of polynomial functions is realized as a quotient of the so called reflection equation algebra and simultaneously as a subalgebra in the quantized universal enveloping algebra.

6.1 Simple groups as Poisson Lie manifolds

Let \mathfrak{g} be a complex simple Lie algebra and G be a connected Lie group corresponding to \mathfrak{g} . An element $g \in G$ is called semisimple if it belongs to a maximal torus in G .

Let r be a classical r -matrix defining a factorizable Lie bialgebra structure on \mathfrak{g} . Denote by r_- and Ω its skew and symmetric parts, respectively. We assume r to be normalized so that Ω is the inverse (canonical element) of the Killing form on \mathfrak{g} .

Given an element $\xi \in \mathfrak{g}$ let ξ^l and ξ^r denote, respectively the left- and right invariant vector fields on G generated by ξ :

$$(\xi^l f)(g) = \frac{d}{dt} f(g e^{t\xi})|_{t=0}, \quad (\xi^r f)(g) = \frac{d}{dt} f(e^{t\xi} g)|_{t=0}$$

for every smooth function f on G .

The Semenov-Tyan-Shansky (STS) Poisson structure [STS] on the group G is defined by the bivector field

$$r_-^{l,l} + r_-^{r,r} - r_-^{r,l} - r_-^{l,r} + \Omega^{l,l} - \Omega^{r,r} + \Omega^{r,l} - \Omega^{l,r} = r_-^{\text{ad,ad}} + (\Omega^{r,l} - \Omega^{l,r}). \quad (17)$$

Here $\xi^{\text{ad}} := \xi^l - \xi^r$, $\xi \in \mathfrak{g}$.

Consider the group G as a G -space with respect to the adjoint action. The STS Poisson structure makes G a Poisson-Lie manifold over the Poisson Lie group G endowed with the bracket $r^{l,l} - r^{r,r}$. In fact, G is a Poisson Lie manifold not only over G , but over a Poisson Lie group corresponding to the double Lie bialgebra $\mathfrak{D}\mathfrak{g}$.

Theorem 6.1 ([AM]). *Symplectic leaves of the STS Poisson structure coincide with conjugacy classes in G .*

Let us compute the restriction of the STS bracket to the class C_g of a semisimple element $g \in G$. The Lie algebra \mathfrak{g} splits into the direct sum $\mathfrak{g} = \mathfrak{l} \oplus \mathfrak{m}$ of vector spaces, where \mathfrak{l} is the eigenspace of $\text{Ad}(g)$ corresponding to the eigenvalue 1 and \mathfrak{m} is the $\text{Ad}(g)$ -invariant subspace where $\text{Ad}(g) - \text{id}$ is invertible. This decomposition is orthogonal with respect to the Killing form. The tangent space to C_g at the point g is identified with \mathfrak{m} , while \mathfrak{l} is the Lie algebra of the centralizer of g .

Let $\{\xi_\mu\} \subset \mathfrak{m}$ be an orthonormal base of eigenvectors of $\text{Ad}(g)$ labelled by the eigenvalues of $\text{Ad}(g)$. We have $(\xi_\mu, \xi_\nu) = 0$ unless $\mu\nu = 1$ and assume the normalization $(\xi_\mu, \xi_{\mu^{-1}}) = 1$. One can check that the restriction of the STS bracket to the tangent spaces at the point g is the bivector

$$r_{\mathfrak{m} \wedge \mathfrak{m}} + \sum_{\mu} \frac{\mu + 1}{\mu - 1} \xi_{\mu} \otimes \xi_{\mu^{-1}} \in \mathfrak{m} \wedge \mathfrak{m},$$

where the first term is the projection of r to $\mathfrak{m} \wedge \mathfrak{m}$. The second term is correctly defined since $\text{Ad}(g) - \text{id}$ is invertible on \mathfrak{m} .

6.2 Quantization of the STS bracket on the group

In this subsection we describe quantization of the STS bracket on classical matrix groups in terms of generators and relations.

By $\mathbb{C}_{\hbar}[G]$ we denote the $\mathcal{U}_{\hbar}(\mathfrak{D}\mathfrak{g})$ -equivariant deformation (quantization) of the affine ring $\mathbb{C}[G]$. This quantization in the form of star product was constructed in [DM3]. Below we give a description of $\mathbb{C}_{\hbar}[G]$ in terms of generators and relations, using the so called reflection equation algebra.

Let (V, π) be the defining representation of $\mathcal{U}_{\hbar}(\mathfrak{g})$ and R the image of \mathcal{R} in $\text{End}(V^{\otimes 2})$. Denote by \mathcal{K} the quotient of the tensor algebra of $\text{End}(V^*)$ by the quadratic relations

$$R_{21}K_1R_{12}K_2 = K_2R_{21}K_1R_{12}. \quad (18)$$

Here $K = \|K_j^i\|$ is the matrix of the generators forming the standard matrix base in $\text{End}(V^*)$. The algebra \mathcal{K} is called reflection equation (RE) algebra, [KSk1]. It is a $\mathbb{C}[[\hbar]]$ -flat deformation of the polynomial ring $\mathbb{C}[\text{End}(V)]$ only for $\mathfrak{g} = \mathfrak{sl}(n)$.

The $\mathcal{U}_{\hbar}(\mathfrak{g})$ -equivariant quantization of $\mathbb{C}[G]$ can be realized as a quotient of \mathcal{K} . Below we describe the corresponding ideals only for G orthogonal and symplectic. That can also be

done for the case $G = SL(n)$, but then the algebra \mathcal{K} is good enough for our purposes, so it is even more convenient to work with \mathcal{K} rather than $\mathbb{C}_\hbar[SL(n)]$.

Assume \mathfrak{g} to be orthogonal or symplectic. Let B be the $\mathcal{U}_\hbar(\mathfrak{g})$ -invariant element from $V \otimes V$, [FRT].

Proposition 6.2. *The algebra $\mathbb{C}_\hbar[G]$ is isomorphic to the quotient of \mathcal{K} by the $\mathcal{U}_\hbar(\mathfrak{g})$ -invariant ideal of relations*

$$R_1^t K^t ((R_{1'})^{-1} B^{-1} (R_2)^{-1}) R_2 K = B^{-1}, \quad K R_1 B K^t R_2^t = R_{1'} B R_{2'}^t. \quad (19)$$

A proof that this quotient is $\mathbb{C}[[\hbar]]$ -free is given in [M].

6.3 Embedding of $\mathbb{C}_\hbar[G]$ in $\mathcal{U}_\hbar(\mathfrak{g})$

Recall that a (left) Yetter-Drinfeld (YD) module over $\mathcal{U}_\hbar(\mathfrak{g})$ is simultaneously a left $\mathcal{U}_\hbar(\mathfrak{g})$ -module, a left $\mathcal{U}_\hbar(\mathfrak{g})$ -comodule, and these two structures are compatible in a certain way, [Y]. A YD algebra over $\mathcal{U}_\hbar(\mathfrak{g})$ is a $\mathcal{U}_\hbar(\mathfrak{g})$ -module algebra, $\mathcal{U}_\hbar(\mathfrak{g})$ -comodule algebra and a YD module with respect to these structures. An example of YD module algebra is $\mathcal{U}_\hbar(\mathfrak{g})$ considered as the adjoint module and comodule via the coproduct.

Let us describe a method of producing YD modules (module algebras) out of $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ -modules, where $\mathfrak{D}\mathfrak{g}$ is the double of \mathfrak{g} . For a factorizable semisimple Lie bialgebra \mathfrak{g} the algebra $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ is isomorphic to the twisted tensor square $\mathcal{U}_\hbar(\mathfrak{g}) \overset{\mathcal{R}}{\otimes} \mathcal{U}_\hbar(\mathfrak{g})$. The universal R-matrix $\hat{\mathcal{R}}$ of $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ is expressed through the universal R-matrix \mathcal{R} of $\mathcal{U}_\hbar(\mathfrak{g})$ by the formula $\hat{\mathcal{R}} = \mathcal{R}_{41}^{-1} \mathcal{R}_{31}^{-1} \mathcal{R}_{24} \mathcal{R}_{23}$. It is easy to see that the right component of $\hat{\mathcal{R}}$ lies, in fact, in $\mathcal{U}_\hbar(\mathfrak{g})$ (via diagonal embedding). Then any $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ -module (module algebra) becomes a YD module (YD module algebra) when equipped with the $\mathcal{U}_\hbar(\mathfrak{g})$ -coaction $\delta(a) = \hat{\mathcal{R}}_2 \otimes \hat{\mathcal{R}}_1 \triangleright a$. The $\mathcal{U}_\hbar(\mathfrak{g})$ -action is induced through the embedding of $\Delta: \mathcal{U}_\hbar(\mathfrak{g}) \hookrightarrow \mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$.

The algebra $\mathbb{C}_\hbar[G]$ is a $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ -algebra and hence YD module algebra, by the above construction. It can be realized in as a $\mathbb{C}[[\hbar]]$ -submodule in the FRT dual to $\mathcal{U}_\hbar(\mathfrak{g})$. The Hopf pairing between the FRT dual and $\mathcal{U}_\hbar(\mathfrak{g})$ induces a pairing between $\mathbb{C}_\hbar[G]$ and $\mathcal{U}_\hbar(\mathfrak{g})$. By means of this pairing, the universal RE matrix \mathcal{Q} implements a $\mathcal{U}_\hbar(\mathfrak{g})$ -algebra homomorphism

$$\mathbb{C}_\hbar[G] \rightarrow \mathcal{U}_\hbar(\mathfrak{g}), \quad a \mapsto \langle a, \mathcal{Q}_1 \rangle \mathcal{Q}_2. \quad (20)$$

It is easy to check using the explicit form of the \mathcal{R} matrix that $\mathbb{C}_\hbar[G]$ lies in $\check{\mathcal{U}}_\hbar(\mathfrak{g}) \subset \mathcal{U}_\hbar(\mathfrak{g})$. The map (20) is, in fact, a homomorphism of YD algebras.

Proposition 6.3. *The map (20) is embedding.*

Proof. The proof easily follows from Proposition 5.2 after a slight adaptation to YD module algebras. The algebra $\mathbb{C}_\hbar[G]$ decomposes into the direct sum $\bigoplus_{W_0} W \otimes W^*$ taken over the simple finite dimensional \mathfrak{g} -modules. Each summand is a YD module and its quotient mod \hbar is $\mathfrak{D}\mathfrak{g} = \mathfrak{g} \oplus \mathfrak{g}$ -irreducible with multiplicity one.

We have a pair of YD module algebras $\mathfrak{S} = \mathbb{C}_\hbar[G]$ and $\mathfrak{T} = \mathcal{U}_\hbar(\mathfrak{g})$. The former is admissible (as a YD-module), while the latter has no torsion. Set Ψ to be the map (20). Its image and kernel are YD modules and are free over $\mathbb{C}[[\hbar]]$. Therefore \mathfrak{S} decomposes into the direct sum $\ker \Psi \oplus \text{im } \Psi$. The rest of the proof is readily adapted from the proof of Proposition 5.2 if one observes that YD modules become $\mathfrak{D}\mathfrak{g}$ -modules in the quasi-classical limit (the first order in \hbar). Put $\mathfrak{J} = 0$. The algebra $\mathfrak{S}_0 = \mathbb{C}[G]$ has no non-zero $\mathfrak{D}\mathfrak{g}$ -invariant ideals. Thus we conclude $\ker \Psi = \mathfrak{J} = 0$. \square

7 Center of the algebra $\mathbb{C}_\hbar[G]$

Let G be a simple complex algebraic group. If G is simply connected, then $\mathbb{C}[G]$ is a free module over the subalgebra of invariants $I(G)$, $[\mathbb{R}]$. More precisely, there exists a submodule $\mathcal{E}_0 \subset \mathbb{C}[G]$ such that the multiplication map $I(G) \otimes \mathcal{E}_0 \rightarrow \mathbb{C}[G]$ is an isomorphism of vector spaces. Each isotypic component in \mathcal{E}_0 enters with finite multiplicity. This fact has a quantum analog.

Theorem 7.1 ([M]). *Let G be a simple complex algebraic group and let $\mathbb{C}_\hbar[G]$ be the $\mathcal{U}_\hbar(\mathfrak{D}\mathfrak{g})$ -equivariant quantization of $\mathbb{C}[G]$ along the STS bracket. Then*

- i) the subalgebra $I_\hbar(G)$ of $\mathcal{U}_\hbar(\mathfrak{g})$ -invariants coincides with the center of $\mathbb{C}_\hbar[G]$,*
- ii) $I_\hbar(G) \simeq I(G) \otimes \mathbb{C}[[\hbar]]$ as \mathbb{C} -algebras.*

Let \hat{G} be the simply connected covering of G . Then

- iii) $\mathbb{C}_\hbar[\hat{G}]$ is a free $I_\hbar(\hat{G})$ -module generated by a $\mathcal{U}_\hbar(\mathfrak{g})$ -submodule $\mathcal{E} \subset \mathbb{C}_\hbar[\hat{G}]$. Each isotypic component in \mathcal{E} is $\mathbb{C}[[\hbar]]$ -finite.*

Theorem 7.1 i) implies, in particular, that the center of $\mathbb{C}_\hbar[G]$ is the intersection of $\mathbb{C}_\hbar[G]$ with the center of $\mathcal{U}_\hbar(\mathfrak{g})$.

If G is not simply connected, the Theorem 7.1 iii) will be true only if the classical algebra of invariants $I(G)$ is polynomial. That is the case, e.g. for $G = SO(2n+1)$, however not so for $G = SO(2n)$.

We will use Theorem 7.1 for quantization of conjugacy classes in Subsection 8.2. Important for us will be the following fact.

Proposition 7.2. *Let G be a simple complex algebraic group. Suppose λ is a character of $I_{\hbar}(G)$, i. e. a unital homomorphism to $\mathbb{C}[[\hbar]]$. Denote by J_λ the ideal in $\mathbb{C}_\hbar[G]$ generated by $\ker \lambda$. Then the quotient $\mathbb{C}_\hbar[G]/J_\lambda$ is an admissible $\mathcal{U}_\hbar(\mathfrak{g})$ -module.*

Proof. The group G is a quotient of its universal covering group \hat{G} over a finite central subgroup $Z \subset \hat{G}$. The affine ring $\mathbb{C}[G]$ is embedded in $\mathbb{C}[\hat{G}]$ as a subalgebra of Z -invariants with respect to the regular action. The group Z naturally acts on the quantized algebra $\mathbb{C}_\hbar[\hat{G}]$, and the subalgebra of Z -invariants is exactly $\mathbb{C}_\hbar[G]$. Accordingly, $I_\hbar(G)$ is the subalgebra of Z -invariants in $I_\hbar(\hat{G})$. The latter is finitely generated over $I_\hbar(G)$, since Z is finite.

Let λ be a character of $I_\hbar(G)$. Denote by J_λ and \hat{J}_λ the ideals generated by $\ker \lambda$ in $\mathbb{C}_\hbar[G]$ and $\mathbb{C}_\hbar[\hat{G}]$, respectively. We have $J_\lambda = \hat{J}_\lambda \cap \mathbb{C}_\hbar[G]$, as follows from Z -invariance. Therefore the quotient $\mathbb{C}_\hbar[G]/J_\lambda$ is embedded in $\mathbb{C}_\hbar[\hat{G}]/\hat{J}_\lambda$. The quotient $\hat{I}_\hbar(G)/\hat{J}_\lambda$ is $\mathbb{C}[[\hbar]]$ -finite and therefore the module $\mathbb{C}_\hbar[\hat{G}]/\hat{J}_\lambda$ has finite isotypic components. Its submodule $\mathbb{C}_\hbar[G]/J_\lambda$ has also finite components. It is admissible, being a submodule of an admissible module. \square

Let us emphasize that the quotient $\mathbb{C}_\hbar[G]/J_\lambda$ is $\mathbb{C}[[\hbar]]$ -free if the classical subalgebra of invariants is a polynomial algebra, cf. remark after Theorem 7.1. Otherwise it may have \hbar -torsion.

7.1 The case of classical series

The case of $G = SO(2n)$ differs from other classical matrix groups. Let us focus on a more simple case of the A, B, C series first.

Proposition 7.3. *The elements τ^ℓ , $\ell = 1, \dots, N$, generate the \hbar -adic completion of the center in the algebras $\mathbb{C}_\hbar[SL(n)]$, $\mathbb{C}_\hbar[SO(2n+1)]$, and $\mathbb{C}_\hbar[Sp(n)]$.*

Proof. The center in $\mathbb{C}_\hbar[G]$ is isomorphic to classical subalgebra of invariants extended by $\mathbb{C}[[\hbar]]$, as stated in Theorem 7.1. Modulo \hbar it is generated by the classical limits of τ^ℓ , i.e. the traces of matrix powers. This implies the statement. \square

In the rest of the subsection we assume $G = SO(2n)$. Let V be the defining $\mathcal{U}_\hbar(\mathfrak{so}(2n))$ -module and $\{\pm \varepsilon_i\}_{i=1}^n = \Lambda(V)$ be the set of weights.

In the classical limit, the n -th exterior power of V_0 splits into two irreducible submodules with highest weights $\pm \varepsilon_n + \sum_{i=1}^{n-1} \varepsilon_i$, [VO]. Therefore the $\mathcal{U}_\hbar(\mathfrak{so}(2n))$ -module $\wedge_q^n V$ (q -antisymmetrized) is a direct sum of two modules (W_\pm, π_\pm) with the highest weights $\pm \varepsilon_n + \sum_{i=1}^{n-1} \varepsilon_i$, respectively. Let $p_\pm: V^{\otimes n} \rightarrow W_\pm$ be the intertwining projectors. The element

$(\Delta^n \otimes \text{id})(\mathcal{Q})$ is expressed through \mathcal{Q} and the numerical matrix $R = (\pi \otimes \pi)(\mathcal{R})$ in the defining representation, by virtue of (12). Hence \mathcal{Q}_{W_\pm} can be explicitly expressed through the matrix \mathcal{Q}_V whose entries generate $\mathbb{C}_\hbar[G]$ within $\mathcal{U}_\hbar(\mathfrak{g})$. In the same fashion, we can define the invariant matrices $K_\pm \in \text{End}(W_\pm) \otimes \mathcal{K}$ expressing them through K , R , and p_\pm by the same formulas. The matrices \mathcal{Q}_{W_\pm} are obtained from K_\pm via the projection $\mathcal{K} \rightarrow \mathbb{C}_\hbar[G]$.

Define a central element τ^- of $\mathcal{U}_\hbar(\mathfrak{so}(2n))$ by setting

$$\tau^- := \text{Tr}_q(\mathcal{Q}_{W_+}) - \text{Tr}_q(\mathcal{Q}_{W_-}). \quad (21)$$

Similarly we introduce the central elements $\text{Tr}_q(K_+) - \text{Tr}_q(K_-) \in \mathcal{K}$.

Proposition 7.4. *Let $\lambda \in \mathfrak{h}^*$ be a weight. Then $\chi^\lambda(\tau^-) = \prod_{i=1}^n (q^{2(\lambda+\rho, \varepsilon_i)} - q^{-2(\lambda+\rho, \varepsilon_i)})$.*

Proof. By Corollary A.2, the central element $\text{Tr}_q(\mathcal{Q}_{W_\pm})$ acts on a module with highest weight λ as multiplication by the scalar $\text{Tr}(\pi_\pm(q^{2(h\lambda+h\rho)}))$. This scalar becomes the group character $\text{ch}_\pm(t)$ (a class function) associated with W_\pm upon the substitution $q^{2(\lambda+\rho, \varepsilon_i)} \rightarrow t_i$, where $\{t_i\}_{i=1}^n$ are the coordinate functions on the maximal torus in $SO(2n)$.

It is known [We] that the ring of characters of the group $SO(2n)$ is isomorphic to $\mathbb{C}[t_1, \dots, t_n, t_1^{-1}, \dots, t_n^{-1}]^{W_{SO(2n)}}$. The group $W_{SO(2n)}$ acts by permutations of the pairs (t_i, t_i^{-1}) and even number of inversions $t_i \leftrightarrow t_i^{-1}$.

The difference $\text{ch}_+(t) - \text{ch}_-(t)$ changes sign under every inversion $t_i \leftrightarrow t_i^{-1}$, hence it is divided by $\prod_{i=1}^n (t_i - t_i^{-1})$. In fact, $\text{ch}_+(t) - \text{ch}_-(t)$ equals $\prod_{i=1}^n (t_i - t_i^{-1})$. This is verified in a standard way by comparing the highest and lowest terms with respect to a natural lexicographic ordering in $\mathbb{C}[t_1, \dots, t_n, t_1^{-1}, \dots, t_n^{-1}]$ (and elementary analysis of the weight structure of $\Lambda^n V_0$). \square

As a corollary, we obtain the following.

Proposition 7.5. *The elements τ^- and τ^ℓ , $\ell = 1, \dots, N$, generate the \hbar -adic completion of the center of $\mathbb{C}_\hbar[SO(2n)]$.*

Proof. Similarly to Proposition 7.3. \square

8 Quantization of conjugacy classes

8.1 Non-exceptional classes

Semisimple conjugacy classes are the only closed classes of a simple algebraic group, [S]. Among semisimple classes of a classical matrix group G we select those which are isomorphic

to semisimple orbits in \mathfrak{g}^* as affine algebraic varieties; we call them "non-exceptional". The isomorphism is implemented by the Cayley transformation $X \mapsto (1 \mp X)(1 \pm X)^{-1}$ for G orthogonal and symplectic. Thus all such classes have Levi subgroups as stabilizers. Non-exceptional covers all semisimple classes for $SL(n)$ and all classes with Levi stabilizers for $Sp(n)$. For orthogonal groups non-exceptional are classes of matrices with no eigenvalues $+1$ and -1 simultaneously.

Let $g \in G$ be a semisimple element. It satisfies a matrix polynomial equation $p(g) = 0$ and defines a character χ^g of the subalgebra of invariants in $\mathbb{C}[G]$.

Theorem 8.1. *Defining ideal $\mathcal{N}(C_g)$ of a non-exceptional conjugacy class C_g is generated by the kernel of χ^g and by the entries of the minimal matrix polynomial for g .*

A proof of this theorem will be given elsewhere. It is based on the following facts. a) Non-exceptional classes are isomorphic to semisimple coadjoint orbits via the Cayley transformation. b) The defining ideals of semisimple coadjoint orbits can be obtained as classical limits of annihilators of generalized Verma modules, [DGS]. c) It was shown in [G] (see also [J2]) that for certain weights the annihilator of the generalized Verma module is generated by a copy of adjoint module in $\mathcal{U}(\mathfrak{g})$ and the kernel of the central character. This is sufficient to describe the defining ideals for all semisimple coadjoint orbits and hence for non-exceptional conjugacy classes.

8.2 The quantization theorem

In this subsection G is a simple complex algebraic group from the classical series and \mathfrak{g} its Lie algebra. Fix a non-exceptional semisimple element g in the maximal torus $T \subset G$. Let \mathfrak{l} be the Lie algebra of the centralizer of g , which is a Levi subgroup. Take $h_\lambda \in \mathfrak{h}$ such that $e^{h_\lambda} = g$ and $\lambda \in \mathfrak{c}_{reg}^*$ is a regular character of \mathfrak{l} .

As was stated in Remark 3.3, the generalized Verma modules can be naturally considered over the ring of polynomials $\mathbb{C}[\mathfrak{c}^*]$, where \mathfrak{c} is the center of \mathfrak{l} . In this case, the parabolic induction is performed from the representation of $\check{\mathcal{U}}_\hbar(\mathfrak{l})$ in $\mathbb{C}[\mathfrak{c}^*][[\hbar]]$. Propositions 4.1 and 4.2 are valid for such modules, since they are valid for generic element of \mathfrak{c}^* . Let (c_i) be coordinate functions generating $\mathbb{C}[\mathfrak{c}^*]$ and (λ_i) the coordinates of a regular element $\lambda \in \mathfrak{c}_{reg}^*$. The homomorphism map $c_i \mapsto \lambda_i/(2\hbar)$ defines a character of $\check{\mathcal{U}}_\hbar(\mathfrak{l})$ (but not of $\mathcal{U}_\hbar(\mathfrak{l})$). Let $\check{M}_{\lambda/2\hbar}$ denote the corresponding Verma module over $\check{\mathcal{U}}_\hbar(\mathfrak{g})$. It is easy to see that the $\check{\mathcal{U}}_\hbar(\mathfrak{g})$ -action on $\text{End}(\check{M}_{\mathfrak{p},\lambda/2\hbar})$ extends to an action of $\mathcal{U}_\hbar(\mathfrak{g})$.

Recall that (V, π) stands for the defining representation of $\mathcal{U}_\hbar(\mathfrak{g})$. Also recall from Subsection 6.2 that the entries of the matrix $\mathcal{Q}_V = \pi(\mathcal{Q}_1) \otimes \mathcal{Q}_2 \in \text{End}(V) \otimes \mathcal{U}_\hbar(\mathfrak{g})$ generate the subalgebra $\mathbb{C}_\hbar[G] \subset \mathcal{U}_\hbar(\mathfrak{g})$, which is the equivariant quantization of $\mathbb{C}[G]$.

Theorem 8.2. *For non-exceptional $\lambda \in \mathfrak{c}^* \subset \mathfrak{h}^*$ the image of $\mathbb{C}_\hbar[G]$ in $\text{End}(\check{M}_{\mathfrak{p}, \lambda/2\hbar})$ is a quantization of the ring of regular functions on the conjugacy class $C_{\exp(\hbar\lambda)} \subset G$. The ideal of the quantized class is generated by the entries of the minimal polynomial in \mathcal{Q}_V over the kernel of a character of $I_\hbar(G)$.*

Proof. The composite map

$$\mathbb{C}_\hbar[G] \hookrightarrow \check{\mathcal{U}}_\hbar(\mathfrak{g}) \rightarrow \text{End}(\check{M}_{\mathfrak{p}, \lambda/2\hbar}) \quad (22)$$

is a $\mathcal{U}_\hbar(\mathfrak{g})$ -equivariant algebra homomorphism. When restricted to the center in $\mathbb{C}_\hbar[G]$, this map defines a character of $I_\hbar(G)$. Indeed, a central element from $\mathbb{C}_\hbar[G]$ acts on $\check{M}_{\mathfrak{p}, \lambda/2\hbar}$ as multiplication by a scalar. That scalar is a polynomial in the eigenvalues of the matrix \mathcal{Q}_V , as follows from Proposition A.1. The eigenvalues are given by (13), where λ should be replaced by $\lambda/2\hbar$. This proves that the representation in $\check{M}_{\mathfrak{p}, \lambda/2\hbar}$ defines a character of the center of $\check{\mathcal{U}}_\hbar(\mathfrak{g})$ and therefore a character of $I_\hbar(G)$. This character is a deformation of the $I(G)$ -character χ^g , and will be denoted by χ_\hbar^g .

The matrix \mathcal{Q}_V satisfies a polynomial equation for a polynomial p with simple roots. The roots of p are given by Theorem 4.2, where again λ should be replaced by $\lambda/2\hbar$. These roots are regular in \hbar and go over to the eigenvalues $e^{(\lambda, \varepsilon_i)}$ of the matrix $g = e^{\hbar\lambda} \in G$.

Next we are going to apply Proposition 5.2.

Put \mathfrak{S} to be the quotient of $\mathbb{C}_\hbar[G]$ over the ideal generated by $\ker \chi_\hbar^g$. The algebra \mathfrak{S} is admissible, due to Proposition 7.2. Put $\mathfrak{T} = \text{End}(\check{M}_{\mathfrak{p}, \lambda/2\hbar})$. The $\mathbb{C}[[\hbar]]$ -module \mathfrak{T} has no torsion, as a subspace of endomorphisms of a torsion free module. By construction, the map (22) factors through an equivariant algebra map $\Psi: \mathfrak{S} \rightarrow \mathfrak{T}$.

Put $\mathfrak{J} \subset \mathbb{C}_\hbar[G]$ to be the ideal generated by the entries of $p(\mathcal{Q}_V)$ projected to \mathfrak{S} . By construction, $\mathfrak{J} \subset \ker \Psi$. The image \mathfrak{J}_0^b of \mathfrak{J}_0 in $\mathfrak{S}_0 = \mathbb{C}[G]/(I(G) - \chi^g)$ is a maximal \mathfrak{g} -invariant ideal, because it is the image of the maximal \mathfrak{g} -invariant ideal $\mathcal{N}(C_g) \subset \mathbb{C}[G]$.

Thus the conditions of Proposition 5.2 are satisfied. Therefore $\mathfrak{S}/\mathfrak{J}$ is the quantization of $\mathbb{C}[C_g]$ and $\ker \Psi = \mathfrak{J}$. In terms of $\mathbb{C}_\hbar[G]$, the defining ideal of $\mathbb{C}[C_g]$ is generated by the kernel of χ_\hbar^g and the entries of $p(\mathcal{Q}_V)$. \square

Next we compute the quantized ideals of non-exception conjugacy classes.

8.3 Ideals of quantized non-exceptional conjugacy classes

First of all, we specialize the formula (16) for \mathfrak{g} being the simple matrix Lie algebra and V the defining representation of $\mathcal{U}_\hbar(\mathfrak{g})$. Since every weight has multiplicity one, we rewrite (16) as

$$\chi^\lambda(\tau^\ell) = \sum_{\nu_i \in \Lambda(V)} x_{\nu_i}^\ell \prod_{\alpha \in \mathbb{R}_+} \frac{q^{(\lambda+\rho+\nu_i, \alpha)} - q^{-(\lambda+\rho+\nu_i, \alpha)}}{q^{(\lambda+\rho, \alpha)} - q^{-(\lambda+\rho, \alpha)}}. \quad (23)$$

Here x_{ν_i} , $\nu_i \in \Lambda(V)$, are the roots of the characteristic polynomial for \mathcal{Q} considered as an operator on $V \otimes M_\lambda$. They are related to the highest weight λ by $x_{\nu_i} = q^{2(\lambda+\rho, \nu_i) - 2(\rho, \nu)}$, $\nu_i \neq 0$ and $x_0 = q^{-2(\rho, \nu) - (\nu, \nu)}$.

Perform the following substitution in (23). Set $x_0 := q^{-2n}$ for $\mathfrak{g} = so(2n+1)$, $x_i = x_{\nu_i}$ for all \mathfrak{g} , and $x_{i'} = x_{-\nu_i} = x_i^{-1} q^{-4(\rho, \nu)}$ for \mathfrak{g} orthogonal and symplectic. Here i ranges from 1 to $n = \text{rk } \mathfrak{g}$. The eigenvalue x_0 is present only for $\mathfrak{g} = so(2n+1)$. Below we use the convention $i' = N+1-i$, cf. Subsection 2.2. Under the adopted enumeration of weights of V the eigenvalues x_i and $x_{i'}$, $i = 1, \dots, N$, correspond to the weights of the opposite sings. As a result of this substitution, we obtain the functions

$$\begin{aligned} \vartheta_{sl(n)}^{(\ell)}(x) &= \sum_{i=1}^n x_i^\ell \prod_{\substack{j=1 \\ j \neq i}}^n \frac{qx_i - x_j \bar{q}}{x_i - x_j}, \\ \vartheta_{so(2n+1)}^{(\ell)}(x) &= \sum_{i=1}^n x_i^\ell \frac{qx_i - x_0}{x_i - x_0 q} \prod_{\substack{j=1 \\ j \neq i}}^n \frac{qx_i - x_j \bar{q}}{x_i - x_j} \frac{qx_i - x_{j'} \bar{q}}{x_i - x_{j'}} + i \leftrightarrow i' + x_0^\ell, \\ \vartheta_{sp(n)}^{(\ell)}(x) &= \sum_{i=1}^n x_i^\ell \frac{q^2 x_i - x_{i'} \bar{q}^2}{x_i - x_{i'}} \prod_{\substack{j=1 \\ j \neq i}}^n \frac{qx_i - x_j \bar{q}}{x_i - x_j} \frac{qx_i - x_{j'} \bar{q}}{x_i - x_{j'}} + i \leftrightarrow i', \\ \vartheta_{so(2n)}^{(\ell)}(x) &= \sum_{i=1}^n x_i^\ell \prod_{\substack{j=1 \\ j \neq i}}^n \frac{qx_i - x_j \bar{q}}{x_i - x_j} \frac{qx_i - x_{j'} \bar{q}}{x_i - x_{j'}} + i \leftrightarrow i', \\ \vartheta_{so(2n)}^-(x) &= q^{2n(\rho, \nu)} \prod_{i=1}^n (x_i - x_{i'}), \end{aligned}$$

where $\bar{q} := q^{-1}$. Remark that $\vartheta_{\mathfrak{g}}^{(\ell)}$ are in fact polynomial in $x_i, x_{i'}, x_0$.

Fix a pair $(\boldsymbol{\mu}, \mathbf{n}) \in \mathbb{C}^k \times \mathbb{Z}_+^k$ such that $\sum_{i=1}^k n_i = n$. Define the vector

$$x(\boldsymbol{\mu}) := \underbrace{(\mu_1, \dots, \mu_1)}_{n_1}, \dots, \underbrace{(\mu_k, \dots, \mu_k)}_{n_k} \in \mathbb{C}^n \quad (24)$$

and $x_q(\boldsymbol{\mu})$ obtained from $x(\boldsymbol{\mu})$ through replacing the "constant" string (μ_i, \dots, μ_i) by "quantum" $(\mu_i, \mu_i q^{-2}, \dots, \mu_i q^{-2(n_i-1)})$ for each $i = 1, \dots, k$. Define functions $\vartheta_{\mathfrak{g}}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu})$ via the substitution $x = x_q(\boldsymbol{\mu})$, $x_0 = q^{-2n}$, to $\vartheta_{\mathfrak{g}}^{(\ell)}(x)$.

Recall from Subsection 2.2 that weights of the defining representation can be expressed in terms of the orthogonal set $\{\varepsilon_i\}$. Next we specialize different types of Levi subalgebras in \mathfrak{g} .

It is convenient to consider $gl(n)$ instead of $sl(n)$. The Levi subalgebras in $gl(n)$ have the form $\mathfrak{l} = \bigoplus_{i=1}^k gl(n_i)$. The irreducible \mathfrak{l} -submodules in the defining \mathfrak{g} -module V_0 are labelled with highest weights

$$\{\varepsilon_1, \varepsilon_{n_1+1}, \dots, \varepsilon_{n_1+\dots+n_{k-1}+1}\}, \quad (25)$$

The Levi subalgebras in the orthogonal and symplectic algebras can be represented as $\mathfrak{l} = \bigoplus_{i=1}^k \mathfrak{l}_i$, where $\mathfrak{l}_i = gl(n_i)$ for $i = 1, \dots, k-1$ and \mathfrak{l}_k equals either $gl(n_k)$ or $so(2n_k+1)$, $sp(n_k)$, $so(2n_k)$ for \mathfrak{g} being $so(2n+1)$, $sp(n)$, $so(2n)$, respectively. That is, the latter option corresponds to \mathfrak{l}_k of the same type as \mathfrak{g} .

In the case $\mathfrak{l}_k = gl(n_k)$ the irreducible \mathfrak{l} -submodules in V_0 are labelled with the highest weights

$$\{\varepsilon_1, \varepsilon_{n_1+1}, \dots, \varepsilon_{n_1+\dots+n_{k-1}+1}; -\varepsilon_{n_1+\dots+n_k}, \dots, -\varepsilon_{n_1+n_2}, -\varepsilon_{n_1}\}, \quad (26)$$

and the zero weight for $so(2n+1)$. Each simple \mathfrak{l} -modules enters with its dual.

For \mathfrak{l}_k of the same type as \mathfrak{g} the irreducible \mathfrak{l} -modules in the vector representation V_0 are labelled with the highest weights

$$\{\varepsilon_1, \varepsilon_{n_1+1}, \dots, \varepsilon_{n_1+\dots+n_{k-2}+1}; \varepsilon_{n-n_k+1}; -\varepsilon_{n_1+\dots+n_{k-1}}, \dots, -\varepsilon_{n_1+n_2}, -\varepsilon_{n_1}\}, \quad (27)$$

We assume that the components μ_i of $\boldsymbol{\mu}$ are non zero and pairwise distinct. For \mathfrak{g} orthogonal and symplectic we also assume $\mu_i \neq \mu_j^{-1}$ for $i \neq j$ and $\mu_i^2 \neq 1$ unless otherwise stated.

Below $Q = Q_V$ is the matrix whose entries generate $\mathbb{C}_\hbar[G] \subset \mathcal{U}_\hbar(\mathfrak{g})$.

8.3.1 The $\mathcal{U}_\hbar(gl(n))$ -case

Put $g = \sum_{i=1}^n g_i e_{ii} \in GL(n)$ to be the diagonal matrix with $g_i = x(\boldsymbol{\mu})_i$. The quantized ideal $\mathcal{N}_\hbar(C_g) \subset \mathbb{C}_\hbar[SL(n)]$ of the class C_g is generated by the relations

$$\prod_{i=1}^k (Q - \mu_i) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{sl(n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}).$$

This reproduces the results of [DM2], where the functions $\vartheta_{sl(n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu})$ are written out in a manifestly polynomial form. In the present form, the central characters of the RE algebras are calculated for the general case of Hecke symmetries in [GS].

8.3.2 The $\mathcal{U}_{\hbar}(so(2n+1))$ -case

1. The case $\mathfrak{l}_k = gl(n_k)$. Set $g := g = \sum_{i=1}^{2n+1} g_i e_{ii} \in SO(2n+1)$ with $g_i = x(\boldsymbol{\mu})_i$ for $i = 1, \dots, n$, $g_{n+1} = 1$, and $g_i = x(\boldsymbol{\mu})_{i'}^{-1}$ for $i = n+2, \dots, 2n+1$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[SO(2n+1)]$ of the class C_g is generated by the relations

$$(Q - q^{-2n}) \prod_{i=1}^k (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2n_i}) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{so(2n+1)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}).$$

2. The case $\mathfrak{l}_k = so(2n_k+1)$. Set g as in the case $\mathfrak{l}_k = gl(n_k)$ with $\mu_k = 1$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[SO(2n+1)]$ of the class C_g is generated by the relations

$$(Q - \mu_k) \prod_{i=1}^{k-1} (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2n_i}) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{so(2n+1)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}),$$

where $\mu_k = q^{2(n_k-n)}$.

8.3.3 The $\mathcal{U}_{\hbar}(sp(n))$ -case

1. The case $\mathfrak{l}_k = gl(n_k)$. Set $g = \sum_{i=1}^{2n} g_i e_{ii} \in Sp(n)$ with $g_i = x(\boldsymbol{\mu})_i$ for $i = 1, \dots, n$, and $g_i = x(\boldsymbol{\mu})_{i'}^{-1}$ for $i = n+1, \dots, 2n$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[Sp(n)]$ of the class C_g is generated by the relations

$$\prod_{i=1}^k (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2(n_i-1)}) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{sp(n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}).$$

2. The case $\mathfrak{l}_k = sp(n_k)$. Set g as in the case $\mathfrak{l}_k = gl(n_k)$ with $\mu_k = 1$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[Sp(n)]$ of the class C_g is generated by the relations

$$(Q - \mu_k) \prod_{i=1}^{k-1} (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2(n_i-1)}) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{sp(n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}),$$

where $\mu_k = q^{2(n_k-n)}$.

8.3.4 The $\mathcal{U}_{\hbar}(so(2n))$ -case

1. The case $\mathfrak{l}_k = gl(n_k)$. Set $g = \sum_{i=1}^{2n} g_i e_{ii} \in SO(2n)$ with $g_i = x(\boldsymbol{\mu})_i$ for $i = 1, \dots, n$, and $g_i = x(\boldsymbol{\mu})_{i'}^{-1}$ for $i = n+1, \dots, 2n$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[SO(2n)]$ of the class C_g is generated by the relations

$$\prod_{i=1}^k (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2(n_i+1)}) = 0, \quad (28)$$

$$\text{Tr}_q(Q_{W_+}) - \text{Tr}_q(Q_{W_-}) = \vartheta^-(\mathbf{n}, \boldsymbol{\mu}), \quad \text{Tr}_q(Q^\ell) = \vartheta_{so(2n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}).$$

Without specializing the value of $\text{Tr}_q(Q_{W_+}) - \text{Tr}_q(Q_{W_-})$ we get a quantization of the intersection of a $SL(2n)$ -class with the group $SO(2n)$. That intersection is an $O(2n)$ -class consisting of two isomorphic (as Poisson Lie manifolds) $SO(2n)$ -classes.

2. The case $\mathfrak{l}_k = \mathfrak{so}(2n_k)$. Set g as in the case $\mathfrak{l}_k = \mathfrak{gl}(n_k)$ with $\mu_k = 1$. The quantized ideal $\mathcal{N}_{\hbar}(C_g) \subset \mathbb{C}_{\hbar}[SO(2n)]$ of the class C_g is generated by the relations

$$(Q - \mu_k) \prod_{i=1}^{k-1} (Q - \mu_i)(Q - \mu_i^{-1} q^{-4n+2(n_i+1)}) = 0, \quad \text{Tr}_q(Q^\ell) = \vartheta_{\mathfrak{so}(2n)}^{(\ell)}(\mathbf{n}, \boldsymbol{\mu}),$$

where $\mu_k = q^{2(n_k-n)}$.

Remark 8.3. To express the quantized ideals in terms of the generators $\{K_j^i\} \subset \mathcal{K}$ one should replace Q by K in the formulas above and impose additional relations (19) in case \mathfrak{g} is orthogonal or symplectic. If $\mathfrak{g} = \mathfrak{sl}(n)$, one can consider relations of Subsection 8.3.1 as those in \mathcal{K} . There are no additional relations needed, and the eigenvalues may take arbitrary pairwise distinct values. This case has been studied in a two parameter setting in [DM2].

A More on central characters

In this appendix we derive some formulas for central characters of Drinfeld-Jimbo quantum groups.

The coproduct on the Cartan-Weyl generators reads

$$\Delta(e_\alpha) = e_\alpha \otimes 1 + q^{h_\alpha} \otimes e_\alpha + \dots, \quad \Delta(e_{-\alpha}) = e_{-\alpha} \otimes q^{-h_\alpha} + 1 \otimes e_{-\alpha} + \dots \quad (29)$$

The omitted terms have lower root vectors in each tensor component.

$$\mathcal{R} = q^{\Omega_{\mathfrak{h}}} \mathcal{R}'(e_{-\alpha} \otimes e_\alpha) = q^{\Omega_{\mathfrak{h}}} \text{ mod } \mathcal{U}'_{\hbar}(\mathfrak{n}^-) \otimes \mathcal{U}'_{\hbar}(\mathfrak{n}^+), \quad (30)$$

where $\mathcal{R}'(e_{-\alpha} \otimes e_\alpha)$ is a series in $e_{-\alpha} \otimes e_\alpha$, $\alpha \in \mathbb{R}_+$ and $\Omega_{\mathfrak{h}} \in \mathfrak{h} \otimes \mathfrak{h}$ is the inverse to the Killing form (the canonical element) restricted to \mathfrak{h} .

Let W be a finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module. Let $X \in \text{End}(W) \otimes \mathcal{U}_{\hbar}(\mathfrak{g})$ be an invariant matrix, i. e. commuting with $\pi(x^{(1)}) \otimes x^{(2)}$ for all $x \in \mathcal{U}_{\hbar}(\mathfrak{g})$. It is known that the q -trace defined by

$$\text{Tr}_q(X^n) := \text{Tr}(\pi_W(q^{2h_\rho})X^n)$$

belongs to the center of $\mathcal{U}_{\hbar}(\mathfrak{g})$. Below we will derive formula for $\chi^\lambda(\text{Tr}_q(Q_W^\ell))$, where χ^λ is a central character $\mathcal{U}_{\hbar}(\mathfrak{g})$ corresponding to λ .

For every weight $\lambda \in \mathfrak{h}^*$ define a linear endomorphism θ_λ of $\mathcal{U}_{\hbar}(\mathfrak{g})$ setting

$$\theta_\lambda(x) := q^{2h_\lambda + 2h_\rho - 2(\rho, \nu)} \mathcal{R}_1 x \mathcal{R}_2.$$

This endomorphism restricts to the subspace of \mathfrak{h} -invariants in $\mathcal{U}_{\hbar}(\mathfrak{g})$, since \mathcal{R} is of zero weight.

Proposition A.1. *Let (W, π_W) be a finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module with the highest weight ν . Then*

$$\chi^\lambda(\mathrm{Tr}_q(\mathcal{Q}_W^\ell)) = q^{-\ell(\nu, \nu)} (\mathrm{Tr}_W \circ \pi_W)(\theta_\lambda^\ell(q^{2h_\rho})) v_\lambda. \quad (31)$$

Proof. We have

$$\chi^\lambda(\mathrm{Tr}_q(\mathcal{Q}_W^\ell)) = \mathrm{Tr}_q\left((\pi \otimes \mathrm{id})(\Delta(z^\ell))\right) q^{-\ell((\lambda, \lambda) + 2(\rho, \lambda) + (\nu, \nu) + 2(\rho, \nu))}.$$

Using induction in ℓ , we find

$$z^\ell = (q^{2h_\rho} \nu)^\ell = q^{2\ell h_\rho} \nu^\ell = q^{2\ell h_\rho} \underbrace{\gamma^{2\ell}(\mathcal{R}_{1'})}_{\ell} \left(\dots \underbrace{(\gamma^2(\mathcal{R}_1) \mathcal{R}_2)}_{\ell} \dots \right) \mathcal{R}_{2'}.$$

Let v_λ be the highest weight vector of M_λ . Consider a linear map $\psi_\lambda: \mathcal{U}_{\hbar}(\mathfrak{g}) \rightarrow \mathbb{C}[[\hbar]]$ defined by $xv_\lambda = \psi_\lambda(x)v_\lambda + \text{lower weight terms}$. Then

$$\chi^\lambda(\mathcal{Q}_W^\ell) = q^{-\ell(\nu, \nu)} (\mathrm{Tr} \circ \pi)(\Upsilon).$$

where we set $\Upsilon = q^{-\ell(\lambda, \lambda) - 2\ell(\rho, \lambda) - 2\ell(\rho, \nu)} q^{2\ell h_\rho} (\mathrm{id} \otimes \psi_\lambda)(\Delta(z^\ell)) \in \mathcal{U}_{\hbar}(\mathfrak{g})$. Using the expression (30) for the universal R-matrix and formulas for comultiplication on generators, we compute Υ to be

$$q^{-2\ell(\rho, \nu)} q^{2\ell h_\rho} q^{2\ell h_\rho} \underbrace{\{q^{h_\lambda} q^{h_\lambda} \gamma^{2\ell}(\mathcal{R}_{1'}) q^{-h_\lambda}\}}_{\ell} \dots \underbrace{\{q^{h_\lambda} q^{h_\lambda} \gamma^2(\mathcal{R}_1) q^{-h_\lambda}\}}_{\ell} \underbrace{\{q^{h_\lambda} \mathcal{R}_2\}}_{\ell} \dots \underbrace{\{q^{h_\lambda} \mathcal{R}_{2'}\}}_{\ell}.$$

We can push the factor q^{h_λ} from before each copy \mathcal{R}_2 to the left till it meets the factor q^{-h_λ} following the corresponding component $\gamma^{2j}(\mathcal{R}_1)$. Since \mathcal{R} is of zero weight, q^{h_λ} commutes with the factors in between. Now recall that γ^2 is given by conjugation with q^{-2h_ρ} . Taking this into account, we obtain

$$\Upsilon = q^{-2\ell(\rho, \nu)} \underbrace{\{q^{2h_\lambda + 2h_\rho} \mathcal{R}_{1'}\}}_{\ell} \dots \underbrace{\{q^{2h_\lambda + 2h_\rho} \mathcal{R}_1\}}_{\ell} q^{2\ell h_\rho} \underbrace{\{\mathcal{R}_2\}}_{\ell} \dots \underbrace{\{\mathcal{R}_{2'}\}}_{\ell} = \theta^{\circ\ell}(q^{2h_\rho}).$$

This makes the proof immediate. □

Corollary A.2. *Let W be a finite dimensional $\mathcal{U}_{\hbar}(\mathfrak{g})$ -module. Then*

$$\chi^\lambda(\mathrm{Tr}_q(\mathcal{Q}_W)) = \mathrm{Tr}(\pi_W(q^{2(h_\lambda + h_\rho)})).$$

Proof. Notice that $\mathcal{R}_1 q^{2h_\rho} \mathcal{R}_2$ equals z . □

References

- [AM] A. Alekseev and Malkin: *Symplectic structures associated to Lie-Poisson groups*, Comm. Math. Phys. **162** (1994) 147–173.
- [DCK] C. De Concini and V. Kac: *Representation of quantum groups at roots of 1*, Operator algebras, unitary representations, enveloping algebras, and invariant theory, Prog. Math., **92** (1990) 471–506.
- [Dr1] V. Drinfeld: *Quantum Groups*, in Proc. Int. Congress of Mathematicians, Berkeley, 1986, ed. A. V. Gleason, AMS, Providence (1987) 798–820.
- [Dr2] V. Drinfeld: *Almost cocommutative Hopf algebras*, Leningrad Math. J. **1** (1990) # 2, 321–342.
- [Dr3] V. Drinfeld: *Quasi-Hopf algebras*, Leningrad Math. J., **1** (1990) 1419–1457.
- [DGS] J. Donin, D. Gurevich, and S. Shnider: *Quantization of function algebras on semisimple orbits in \mathfrak{g}^** , arXiv:q-alg/9607008.
- [DKM] J. Donin, P. P. Kulish, and A. Mudrov: *On a universal solution to reflection equation*, Lett. Math. Phys, **63** (2003) #3, 179–194.
- [DM1] J. Donin and A. Mudrov: *Dynamical Yang-Baxter equation and quantum vector bundles*, arXiv: math.QA/0306028.
- [DM2] J. Donin and A. Mudrov, *Explicit equivariant quantization on coadjoint orbits of $GL(n, \mathbb{C})$* , Lett. Math. Phys, **62** (2002) 17–32.
- [DM3] J. Donin and A. Mudrov: *Reflection Equation, Twist, and Equivariant Quantization*, Isr. J. Math., **136** (2003) 11–28.
- [EE] B. Enriquez and P. Etingof: *Quantization of classical dynamical r-matrices with non-abelian base*, arXiv: math.QA/0311224.
- [EEM] B. Enriquez, P. Etingof, and I. Marshall: *Quantization of some Poisson-Lie dynamical r-matrices and Poisson homogeneous spaces*, arXiv: math.QA/0403283.
- [FRT] L. Faddeev, N. Reshetikhin, and L. Takhtajan: *Quantization of Lie groups and Lie algebras*, Leningrad Math. J., **1** (1990) 193–226.

- [G] R. K. Gupta: *Copies of the adjoint representation inside induced ideals*, preprint, (1985) Paris.
- [GS] D. Gurevich and P. Saponov: *Geometry of non-commutative orbits related to Hecke symmetries* math.QA/0411579.
- [GZB] M. Gould, R. Zhang, and A. Braken: *Generalized Gel'fand invariants and characteristic identities for quantum groups*, J. Math. Phys., **32** (1991) 2298–2303.
- [Jan1] J. C. Jantzen: *Kontravariante formen und Induzierten Darstellungen halbeinfacher Lie-Algebren* Math. Ann. **226** (1977) 53–65.
- [Jan2] J. C. Jantzen: *Lectures on quantum groups*. Grad. Stud. in Math., **6**. AMS, Providence, RI, 1996.
- [Ji] M. Jimbo: *A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985) 63–69.
- [J1] A. Joseph: *Quantum groups and their primitive ideals*. Springer-Verlag, Berlin, 1995.
- [J2] A. Joseph: *A criterion for an ideal to be induced*, J. Algebra **110** (1987) 480–497.
- [JT] A. Joseph and D. Todorovic: *On the quantum KPRV determinants for semisimple and affine Lie algebras*, Alg. Rep. Theor., **5**, (2002) 57–99.
- [K] B. Kostant: J. Funct. Anal. *On the tensor product of a finite and an infinite dimensional representation*, **20** (1975) 257–285.
- [Ke] M. K  b  : *\mathcal{O} -alg  bres quantiques*, C. R. Acad. Sci. Paris Sr. I Math. **322** (1996) 1–4.
- [KhT] S. Khoroshkin and V. Tolstoy: *Universal R -matrix for quantized (super)algebras*, Comm. Math. Phys. **141** (1991) 599–617.
- [KSk] P. P. Kulish and E. K. Sklyanin, *Algebraic structure related to the reflection equation*, J. Phys. A, **25** (1992) 5963–2389.
- [L] G. Lusztig: *Introduction to quantum groups*, Prog. Math., **110**, Boston, MA, 1993.
- [M] A. Mudrov: *On quantization of Semenov-Tian-Shansky Poisson bracket on simple algebraic groups*, math.QA/0412360.

- [N] M. Naimark: *Teoriya predstavlenii grupp.* (Russian) [Representation theory of groups], Moscow, 1976.
- [R] R. Richardson: *An application of the Serre conjecture to semisimple algebraic groups*, Lect. Notes in Math. **848** (1981) 141–151.
- [S] T. Springer: *Conjugacy classes in algebraic groups*, Lect. Not. Math., **1185** (1984) 175–209.
- [STS] M. Semenov-Tian-Shansky: *Poisson-Lie Groups, Quantum Duality Principle, and the Quantum Double*, Contemp. Math., **175** (1994) 219–248.
- [VO] E. Vinberg and A. Onishchik: *Seminar po gruppam Li i algebraicheskim gruppam.* (Russian) [A seminar on Lie groups and algebraic groups], Moscow, 1988.
- [We] H. Weyl: *The classical groups. Their Invariants and representations*, Princeton, New Jersey, 1966.
- [Y] D. Yetter: *Quantum groups and representations of monoidal categories*, Math. Proc. Cambridge Philos. Soc. **108** (1990) # 2, 261–290.