

**Quasistatic Small-strain Plasticity in the Limit of  
Vanishing Hardening and its Numerical Approximation**

**Sören Bartels, Alexander Mielke, Tomas Roubicek**

**no. 490**

Diese Arbeit ist mit Unterstützung des von der Deutschen Forschungsgemeinschaft getragenen Sonderforschungsbereichs 611 an der Universität Bonn entstanden und als Manuskript vervielfältigt worden.

Bonn, Februar 2011

# QUASISTATIC SMALL-STRAIN PLASTICITY IN THE LIMIT OF VANISHING HARDENING AND ITS NUMERICAL APPROXIMATION

SÖREN BARTELS, ALEXANDER MIELKE, AND TOMÁŠ ROUBÍČEK

ABSTRACT. The quasistatic rate-independent evolution of the Prager-Ziegler-type model of linearized plasticity with hardening is shown to converge to the rate-independent evolution of the Prandtl-Reuss elastic/perfectly plastic model. Based on the concept of energetic solutions we study the convergence of the solutions in the limit for hardening coefficients converging to 0 by using the abstract method of  $\Gamma$ -convergence for rate-independent systems. An unconditionally convergent numerical scheme is devised and 2D and 3D numerical experiments are presented. A two-sided energy inequality is a posteriori verified to document experimental convergence rates.

*Mathematical Subject classification:* 35K65, 35K85, 49S05, 65M60, 74C05.

## 1. INTRODUCTION, PLASTICITY WITH HARDENING

There is an extensive engineering and mathematical literature addressing plasticity with hardening, e.g. [1, 2, 3, 18, 20, 21, 22, 34, 47, 49, 50, 51, 54]. The simplest example involving a quadratic stored energy is *linearized plasticity*. We will deal with the simplest variant of this linearized plasticity with isotropic hardening and possibly also kinematic hardening. The rate-independent plasticity without hardening, i.e. the *Prandtl-Reuss* elastic/perfectly plastic model, received attention already long time ago, see e.g. in [25, 28, 34, 46]. Since models without hardening are degenerate, they are mathematically much more challenging as the displacements lie in  $BD(\Omega)$  only rather than in the much better space  $H^1(\Omega)$ . We refer to [11] for a recent analysis in the context of so-called energetic solutions. This work will be a crucial basic for our analysis.

The aim of this article is to show that starting from a model with hardening we have classical solutions that converge in the limit of vanishing hardening to solutions of the Prandtl-Reuss model. Beside justification of the Prandtl-Reuss model as such a limit, the motivation of considering and implementing a small hardening is also numerically justified, e.g. a-posteriori error estimates and convergence of iterative schemes can be proved [3, 19, 47]. One should nevertheless mention that there are algorithms allowing for a direct treating of the Prandtl-Reuss model without hardening, e.g. Newton's schemes work often very well for the standard models with zero hardening, but there is no convergence analysis for the time-dependent case (see [46, 47] for static and incremental problems). Thus, our paper can be seen as a natural synthesis of the rate-independent evolutionary approach from [11] and the numerical approximation with small hardening used in [46, 47].

Let us still remark that similar convergence analysis has been carried out for gradient-plasticity, using the gradient of the plastic strain both in stored energy and in dissipation potential. Namely, for vanishing coefficients in these gradient terms,

---

*Date:* 21.12.2010.

*Key words and phrases.* Rate-independent plasticity, hardening, Prandtl-Reuss elastic/perfectly plastic model, energetic solution, convergence, finite elements.

in [17, Sect.9.2], this model was shown to converge to the Prandtl-Reuss model in terms of the energetic solutions. A similar justification has been given by vanishing viscosity in the plastic flow rule [53]. The case of vanishing inertia is treated in [36], but only for the case of fixed positive hardening. The peculiarity of the vanishing hardening presented here is the changing structure, because the limit Prandtl-Reuss problem has a different set of internal variables, since the isotropic-hardening parameter disappears. Moreover, the vanishing-hardening limit is especially adapted to analyze the simultaneous limit with temporal and finite-element discretizations, see our Section 5.

The plasticity problem is considered to be *rate independent*, i.e. no inertia is considered (which is usually called *quasistatic* and there are no internal time scales. Thus, the problem is invariant under time rescaling. This allows us to use the so-called *energetic formulation*, which in our convex case is fully equivalent to the commonly used evolutionary variational inequalities, see [35]. However, the energetic formulation gives rise to a useful  $\Gamma$ -convergence theory, which will be employed in Sections 4 and 5.

The elastoplastic evolution is given in terms of a time-dependent *stored energy* functional  $\mathcal{E}$  and a *dissipation potential*  $\mathcal{R}$  being positively homogeneous of degree 1 (i.e.  $\mathcal{R}(\lambda z) = \lambda \mathcal{R}(z)$  for  $\lambda \geq 0$ ), which reflects the *rate-independence* of the process, namely the invariance under any monotone re-scaling of time. Both functionals are defined with respect to a suitable state space  $Q := U \times Z$ , where  $U$  and  $Z$  are Banach spaces. The triple  $(Q, \mathcal{E}, \mathcal{R})$  will be called a *rate-independent system*. Thereby a state  $q = (u, z) \in U \times Z = Q$  is given by the *displacement* field  $u$  and the *internal variable*  $z$  that describes here plastic strain and possibly hardening, in some other applications it may also be damage or some phase-transformation variables. We assume that  $\mathcal{R}$  involves only  $z$ , which distinguishes it as a “slow” variable while  $u$  is a “fast” variable. We choose a further Banach space  $X \supset Z$  on which  $\mathcal{R}$  is coercive.

Formally, the evolution of a rate-independent system  $(Q, \mathcal{E}, \mathcal{R})$  is given through solutions  $q : [0, T] \rightarrow Q$  of the following system of *doubly nonlinear* degenerate *parabolic/elliptic variational inclusions*:

$$\partial_u \mathcal{E}(t, u(t), z(t)) \ni 0 \quad \text{and} \quad \partial \mathcal{R}(\dot{z}(t)) + \partial_z \mathcal{E}(t, u(t), z(t)) \ni 0 \quad (1.1)$$

for a.a.  $t \in (0, T)$ , where “ $\partial$ ” refers to a (partial) subdifferential, using that  $\mathcal{R}(\cdot)$ ,  $\mathcal{E}(t, \cdot, z)$ , and  $\mathcal{E}(t, u, \cdot)$  are always here convex functionals. The first relation is the balance of forces and the second is the plastic flow rule.

For our subsequent work it turns out that the notion of *energetic solutions* is better suited. The following precise definition is the basis for the rest of this paper. A function  $q : [0, T] \rightarrow Q$  is called an energetic solution for  $(Q, \mathcal{E}, \mathcal{R})$ , if  $t \mapsto \mathcal{E}(t, q(t))$  lies in  $L^\infty([0, T])$ , if  $t \mapsto \mathcal{E}'_t(t, q(t))$  lies in  $L^1([0, T])$ , and if for all  $t \in [0, T]$  the stability condition (S) and the energy balance (E) hold:

$$(S) \quad \mathcal{E}(t, q(t)) \leq \mathcal{E}(t, \hat{q}) + \mathcal{R}(\hat{z} - z(t)) \quad \text{for all } \hat{q} = (\hat{u}, \hat{z}) \in Q, \quad (1.2a)$$

$$(E) \quad \mathcal{E}(t, q(t)) + \text{Diss}_{\mathcal{R}}(z, [0, t]) = \mathcal{E}(0, q_0) + \int_0^t \mathcal{E}'_s(s, q(s)) \, ds, \quad (1.2b)$$

with  $\mathcal{E}'_t(t, q) = \frac{\partial}{\partial t} \mathcal{E}(t, q)$  and the dissipation functional  $\text{Diss}_{\mathcal{R}}(z, [0, t]) := \sup \sum_{j=1}^N \mathcal{R}(z(t_j) - z(t_{j-1}))$ , where the supremum is taken over all partitions of  $[0, t]$ .

The major advantage of the concept of energetic solutions is that the formulation (1.2) is derivative free: it does not contain  $\dot{q}$ ,  $\partial \mathcal{R}$ , nor  $\mathcal{E}'_q$ . Moreover, it shows immediately the basic energetic a priori estimates. If  $\mathcal{E}$  is coercive in  $Z$  we immediately

obtain

$$u \in \mathbf{B}([0, T]; U) \quad \text{and} \quad z \in \mathbf{B}([0, T]; Z) \cap \mathbf{BV}([0, T]; X),$$

where we used  $\text{Diss}_{\mathcal{R}}(z, [0, T]) < \infty$  and the assumed coercivity of  $\mathcal{R}$  in  $X$ ; here “ $\mathbf{B}([0, T]; \cdot)$ ” and “ $\mathbf{BV}([0, T]; \cdot)$ ” stands for bounded measurable and bounded variation functions on  $[0, T]$ , respectively. The equivalence of (1.1) and the energetic solution is standard for the case with hardening (see e.g. [35, Sect. 2]), while for the case without hardening the more subtle equivalence is established in [11].

We are now in a position to explain the main results and ideas of the paper. In Section 2 we describe the plasticity model with hardening. In that situation  $\mathcal{E}$  is uniformly convex (implying a quadratic coercivity) and we choose as state space  $Q$  a closed subspaces of the Hilbert space  $\mathbf{H}^1(\Omega; \mathbb{R}^d) \times \mathbf{L}^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \times \mathbf{L}^2(\Omega)$ . The classical existence result for solutions (cf. e.g. [20, 24]) is stated in Proposition 2.1 for completeness.

In Section 3 we present the Prandtl-Reuss model for perfect elastoplasticity without hardening. We still have convexity, but the coercivity only holds with a linear lower bound. Hence, the underlying state space  $Q_{\text{PR}}$  becomes a closed subspace of  $\mathbf{BD}(\Omega; \mathbb{R}^d) \times \mathcal{M}(\Omega \cup \Gamma_{\text{D}}; \mathbb{R}_{\text{dev}}^{d \times d})$ . One major problem in our limit passage is that  $Q$  for the hardening case is only weakly\* dense in  $Q_{\text{PR}}$  but not strongly. In Proposition 3.2 we provide a slight generalization of the existence result in [11] for energetic solutions for the Prandtl-Reuss model written a rate-independent system  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$ . Using different techniques similar existence results were obtained much earlier in [25, 53]. For simplicity, we have restricted all our work to the case without volume and surface forces, and thus we do not need the classical *safe-load condition*. We restricted the loading to time-dependent Dirichlet boundary data  $u_{\text{D}}$ , but everything presented here can be generalized to included volume and surface loading, when adding a suitable the safe-load condition.

In Section 4 we consider a family  $(Q, \mathcal{E}_{\varepsilon}, \mathcal{R})$  of rate-independent systems and study the limit of its energetic solutions  $q_{\varepsilon} : [0, T] \rightarrow Q$  for  $\varepsilon \rightarrow 0$ . To study the case of vanishing hardening, we assume that  $\mathcal{E}_{\varepsilon}$  has the form

$$\mathcal{E}_{\varepsilon}(t, u, \pi, \eta) := \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u + 2u_{\text{D}}(t)) - \pi) : (e(u) - \pi) + \frac{\varepsilon}{2} \mathbb{H} \pi : \pi + \frac{\varepsilon}{2} b \eta^2 \, dx,$$

where  $\varepsilon > 0$  measures the size of the hardening. Here  $\pi(t, x) \in \mathbb{R}_{\text{dev}}^{d \times d}$  is the plastic strain tensor and  $\eta$  the scalar isotropic hardening parameter. The dissipation potential (or equivalently on the elastic domain) is assumed to be independent of  $\varepsilon$  and to satisfy the quite general structure condition (4.1). Then, based on suitable a priori bounds and the abstract theory of  $\Gamma$ -convergence for energetic solutions of rate-independent systems (see [38]), Theorem 4.1 states that the solutions  $q_{\varepsilon} = (u_{\varepsilon}, \pi_{\varepsilon}, \eta_{\varepsilon})$  converge (in terms of subsequences of  $u$ - and  $\pi$ -components) to energetic solutions  $(u, \pi)$  of the Prandtl-Reuss system  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$ .

In Section 5 we study the joint convergence of vanishing hardening and space-time discretization. Choosing the time step  $\tau = T/N$  we define discrete solutions via the incremental problem

$$\text{Minimize } \mathcal{E}_{\varepsilon}(k\tau, u, \pi, \eta) + \mathcal{R}(\pi - \pi_{\varepsilon\tau h}^{k-1}, \eta - \eta_{\varepsilon\tau h}^{k-1}) \quad \text{subject to } (u, \pi, \eta) \in Q_h,$$

where  $h$  is the mesh parameter, i.e. the maximal mesh size in the finite-element discretization for the functions in  $Q$  (for conformal P1 elements for  $u$  and P0 elements for  $(\pi, \eta)$ ). Again using the abstract  $\Gamma$ -convergence theory we establish convergence of (a subsequence of) the associated solutions  $\bar{q}_{\varepsilon\tau h}$  (piecewise constant interpolant of the incremental solutions) to energetic solutions of the Prandtl-Reuss system

$(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$ . We cannot establish convergence of the whole family  $\bar{q}_{\varepsilon\tau h}$ , as it is not known that the solutions of the Prandtl-Reuss system are unique. However, since it is shown in [11, Thm. 5.9] that the stresses

$$\sigma = \mathbb{C}(e(u+u_{\text{D}}(t))-\pi) \quad (1.3)$$

are uniquely defined, and conclude that the stresses  $\bar{\sigma}_{\varepsilon\tau h}$  converge without choosing a subsequence. This also explains why in [46] only convergence of the stresses (even with explicit error rates) could be shown.

Section 6 reports on numerical experiments where one two-dimensional and one three-dimensional situation are studied, which are set up in such a way that a shear band must form. We study the influence of varying the hardening parameter  $\varepsilon$  and the discretization parameters  $h$  and  $\tau$ .

## 2. PLASTICITY WITH HARDENING

Let us define

$$\mathbb{R}_{\text{sym}}^{d \times d} := \left\{ A \in \mathbb{R}^{d \times d} \mid A^{\text{T}} = A \right\} \quad \text{and} \quad \mathbb{R}_{\text{dev}}^{d \times d} := \left\{ A \in \mathbb{R}_{\text{sym}}^{d \times d} \mid \text{tr } A = 0 \right\}. \quad (2.1)$$

We consider the classical formulation of linearized elastoplasticity with a trace-free plastic strain tensor  $\pi \in \mathbb{R}_{\text{dev}}^{d \times d}$  and a scalar, isotropic hardening parameter  $\eta$ . The system consists of the mechanical equilibrium  $\text{div } \sigma + f = 0$  for the elastic stress  $\sigma = \mathbb{C}(e(u) - \pi)$ , i.e.

$$-\text{div}(\mathbb{C}(e(u) - \pi)) = f, \quad \text{where} \quad e(u) = \frac{1}{2}(\nabla u + (\nabla u)^{\text{T}}) \in \mathbb{R}_{\text{sym}}^{d \times d}. \quad (2.2a)$$

and the evolution law (flow rule) for the internal, plastic variable  $z = (\pi, \eta)$  in the form

$$\partial R \begin{pmatrix} \dot{\pi} \\ \dot{\eta} \end{pmatrix} + \begin{pmatrix} \text{dev } \mathbb{C}(\pi - e(u)) \\ b\eta \end{pmatrix} + \mathbb{H}\pi \ni \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (2.2b)$$

Here  $\mathbb{C}$  is the symmetric positive semidefinite 4th-order elastic moduli tensor, i.e.  $\mathbb{C}_{ijkl} = \mathbb{C}_{jikl} = \mathbb{C}_{klij}$  for all  $i, j, k, l = 1, \dots, d$ . The *kinematic hardening tensor*  $\mathbb{H}$  is a symmetric and positive semidefinite operator on  $\mathbb{R}_{\text{dev}}^{d \times d}$  (of Prager/Ziegler type [44, 55]), whereas the scalar parameter  $b \geq 0$  determines the *isotropic hardening*.

The dissipation potential  $R$  is assumed to be lower semicontinuous, convex, and positively homogeneous of degree 1. The latter condition reflects rate independence, since the subdifferential  $\partial R(\cdot)$  is homogeneous of degree 0. We define the elastic domain via  $S = \partial R(0) \subset \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R}$ , which is closed and convex, and denote by  $\delta_S$  the convex indicator function, i.e.  $\delta_S(\zeta, \xi) = 0$  for  $(\zeta, \xi) \in S$  and  $+\infty$  otherwise. Then, we have  $R = \delta_S^*$ , where  $\delta_S^*$  the Legendre-Fenchel conjugate functional to  $\delta_S$ . Throughout we assume that  $R$  satisfies the coercivity estimate

$$\exists c > 0 \forall (\dot{\pi}, \dot{\eta}) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R} : \quad R(\dot{\pi}, \dot{\eta}) \geq c(|\dot{\pi}| + |\dot{\eta}|).$$

This condition is equivalent to assuming that  $S$  contains an open neighborhood of  $(0, 0) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R}$ .

Then, by the classical duality relation of convex analysis we have

$$[\partial R]^{-1} = \partial R^* = \partial \delta_S^* = \partial \delta_S = N_S = \text{the normal-cone mapping for } S.$$

The problem (2.2) can thus equivalently be written in a form which is more standard in the engineering literature, namely

$$\operatorname{div} \sigma + f = 0 \quad \text{with } \sigma = \mathbb{C}(e(u) - \pi), \quad (2.3a)$$

$$\frac{\partial}{\partial t} \begin{pmatrix} \pi \\ \eta \end{pmatrix} \in N_S \left( \begin{array}{c} \operatorname{dev} \sigma - \mathbb{H}\pi \\ -b\eta \end{array} \right), \quad (2.3b)$$

which reveals that  $\mathbb{H}\pi$  is in the position of the *back stress* to the *elastic stress*  $\sigma$ .

We assume that such elasto-plastic body occupies a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^d$ . The system is to be completed by boundary conditions. For this, we divide the boundary  $\partial\Omega$  into two disjoint parts,  $\Gamma_D$  and  $\Gamma_N$ , and consider

$$u|_{\Gamma_D} = u_D(t) \quad \text{on } \Gamma_D, \quad (\mathbb{C}(e(u) - \pi))\nu = g \quad \text{on } \Gamma_N = \partial\Omega \setminus \Gamma_D. \quad (2.4)$$

Throughout this work we assume that  $u_D(t)$  may depend on time and is defined in all of  $\Omega$ , and that

$$\Gamma_D \text{ has a nonempty relative interior and} \quad (2.5a)$$

$$\text{a } (d-2)\text{-dimensional Lipschitz boundary.} \quad (2.5b)$$

Further, we make a transformation to the homogeneous Dirichlet boundary conditions, i.e. replace  $u$  by  $u + u_D$ . Neglecting an (unimportant) additive constant thus created in the shifted functional  $\mathcal{E}$ , the function-space setting and the energetics can be defined by

$$U := \left\{ u \in H^1(\Omega; \mathbb{R}^d) \mid u|_{\Gamma_D} = 0 \right\}, \quad Z := L^2(\Omega; \mathbb{R}_{\operatorname{dev}}^{d \times d}) \times L^2(\Omega), \quad (2.6a)$$

$$Q := U \times Z, \quad X := L^1(\Omega; \mathbb{R}_{\operatorname{dev}}^{d \times d}) \times L^1(\Omega), \quad (2.6b)$$

$$\mathcal{E}(t, u, \pi, \eta) := \frac{1}{2} \int_{\Omega} \mathbb{C}(e(u) - \pi) : (e(u) - \pi) + \mathbb{H}\pi : \pi + b\eta^2 \, dx - \langle f_{\text{ext}}(t), (u, \pi) \rangle, \quad (2.6c)$$

$$\mathcal{R}(\dot{\pi}, \dot{\eta}) := \int_{\Omega} \delta_S^*(\dot{\pi}, \dot{\eta}) \, dx \quad (2.6d)$$

with the extended force  $f_{\text{ext}}(t) \in U^* \times Z^*$  in (2.6c) being defined by

$$\langle f_{\text{ext}}(t), (u, \pi) \rangle := \int_{\Omega} f(t) \cdot u - \mathbb{C}e(u_D(t)) : (e(u) - \pi) \, dx + \int_{\Gamma_N} g(t) \cdot u \, d\mathcal{H}^{d-1}. \quad (2.6e)$$

As the stability condition (S) in (1.2a) is an intrinsic part of the definition of energetic solutions, we introduce the set of stable states at time  $t$  via

$$\mathcal{S}(t) := \left\{ q \in Q \mid \mathcal{E}(t, q) < \infty, \forall \hat{q} \in Q: \mathcal{E}(t, q) \leq \mathcal{E}(t, \hat{q}) + \mathcal{R}(\hat{q} - q) \right\}. \quad (2.7)$$

Then, assuming

$$u_D \in W^{1,1}(I; H^1(\Omega; \mathbb{R}^d)), \quad f \in W^{1,1}(I; L^{p_f}(\Omega; \mathbb{R}^d)), \quad g \in W^{1,1}(I; L^{p_g}(\Gamma; \mathbb{R}^d)), \quad (2.8)$$

with  $p_f = 2d/(d+2)$  and  $p_g = 2-2/d$  if  $d > 2$ , or  $p_f > 1$  and  $p_g > 1$  for  $d = 2$ , or  $p_f = 1 = p_g$  for  $d = 1$ , we have  $f_{\text{ext}} \in W^{1,1}(I; U^* \times Z^*)$  and the proof of the following proposition is standard, see [1, 20, 21, 22, 24, 35].

**Proposition 2.1** (Existence of energetic solutions). *Let  $\mathbb{C}$ ,  $\mathbb{H}$ ,  $b \geq 0$ ,  $S$ ,  $U$ , and  $Z$  be as described above, and (2.8) hold. Moreover assume that  $q_0 = (u_0, \pi_0, \eta_0) \in$*

$\mathcal{S}(0)$ , i.e.  $q_0$  is stable at time  $t = 0$ . Furthermore, let the hardening form be coercive on  $\text{dom}(\delta_S^*)$ , i.e.

$$\inf \left\{ \frac{\mathbb{H}\pi:\pi + b\eta^2}{|\pi|^2 + \eta^2} \mid 0 \neq (\pi, \eta) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R}, \delta_S^*(\pi, \eta) < \infty \right\} > 0. \quad (2.9)$$

Then, there is a unique energetic solution  $q = (u, \pi, \eta)$  with  $q(0) = q_0$  for the energetic rate-independent system  $(U \times Z, \mathcal{E}, \mathcal{R})$  defined in (2.6). Moreover, we have  $q \in W^{1,1}(I; U \times Z)$ .

### 3. PLASTICITY WITHOUT HARDENING

In the case without hardening, i.e.  $\mathbb{H} = 0$  and  $b = 0$ , one obtains the *Prandtl-Reuss system* of perfect elastoplasticity. The peculiarity here is that the displacement no longer lives in the conventional Sobolev  $H^1$ -space but rather in the space of *functions with bounded deformations* introduced in [52], defined as

$$\text{BD}(\Omega; \mathbb{R}^d) := \left\{ u \in L^1(\Omega; \mathbb{R}^d) \mid e(u) \in \mathcal{M}(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}) \right\}, \quad (3.1)$$

where  $e(u)$  is the symmetric part of distributional gradient of  $u$ . For general bounded Borel sets  $S \subset \mathbb{R}^d$  we set  $\mathcal{M}(S, V) = C_0(S; V)^*$  denotes  $V$ -valued Radon measures on  $\Omega$  (here  $C_0(S; V)$  denotes continuous function with compact support on  $S$ ). The space  $\text{BD}(\Omega; \mathbb{R}^d)$  has a predual space, and thus a weak\* topology. The weak\* convergence means weak convergence in  $L^1(\Omega; \mathbb{R}^d)$  together with weak\* convergence of  $e(u)$  in  $\mathcal{M}(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$ . See [54, 11] for many details about BD-functions.

The rate-independent system  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$  for the Prandtl-Reuss system is given by

$$U_{\text{PR}} := \text{BD}(\Omega; \mathbb{R}^d), \quad Z_{\text{PR}} = X_{\text{PR}} := \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}), \quad (3.2a)$$

$$Q_{\text{PR}} := \left\{ (u, \pi) \in U_{\text{PR}} \times Z_{\text{PR}} \mid e(u) - \pi|_{\Omega} \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}), \right. \\ \left. u \otimes \nu d\mathcal{H}^{d-1} + \pi|_{\Gamma_D} = 0 \text{ on } \Gamma_D \right\}, \quad (3.2b)$$

$$\mathcal{E}_{\text{PR}}(t, u, \pi) := \frac{1}{2} \int_{\Omega} \mathbb{C}(e(u + 2(u_D(t)) - \pi)) : (e(u) - \pi) dx, \quad (3.2c)$$

$$\mathcal{R}_{\text{PR}}(\dot{\pi}) := \int_{\Omega \cup \Gamma_D} \delta_P^*(\cdot) d\dot{\pi}(x) \quad \text{for } \dot{\pi} \in \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}), \quad (3.2d)$$

where  $a \otimes b$  means the symmetrized tensorial product  $\frac{1}{2}(a \otimes b + b \otimes a)$  and where  $P \subset \mathbb{R}_{\text{dev}}^{d \times d}$  is the elasticity domain, which is assumed to be a bounded, closed convex neighborhood of 0. Equivalently, (3.2d) can also be written as  $\mathcal{R}_{\text{PR}}(\dot{\pi}) = \int_{\Omega \cup \Gamma_D} \delta_P^*\left(\frac{d\dot{\pi}}{d|\dot{\pi}|}\right) d|\dot{\pi}|$  where  $|\dot{\pi}|$  is the total variation of  $\dot{\pi}$  and  $\frac{d\dot{\pi}}{d|\dot{\pi}|}$  is the Radon-Nykodym derivative of  $d\dot{\pi}$  with respect to  $|\dot{\pi}|$ . See [11] for further details about functions on  $\mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d})$ .

Functions from  $\text{BD}(\Omega; \mathbb{R}^d)$  have traces in  $L^1(\partial\Omega; \mathbb{R}^d)$ . However, one has to be aware of jumps that can occur at the boundary, i.e. the measure  $e(u)$  may concentrate on the boundary  $\partial\Omega$ . Thus, the classical boundary condition  $u = 0$  on  $\Gamma_D$  used in the definition of  $Q$  in (2.6b) is replaced by the more involved relation  $u \otimes \nu d\mathcal{H}^{d-1} + \pi|_{\Gamma_D} = 0$  on  $\Gamma_D$  for  $Q_{\text{PR}}$  in (3.2b). This relation has to be understood as an equality of measures on  $\Gamma_D$ , viz.

$$\forall \text{measurable } A \subset \Gamma_D : \int_A u \otimes \nu d\mathcal{H}^{d-1} = \int_A d\pi = \pi(A).$$

The relation simply means that any jump of  $u$  on the boundary with respect to the Dirichlet condition  $u = 0$  is due to a localized plastic deformation.

In fact, to define traces properly, one use suitable extensions of functions into a neighborhood of  $\Omega$ . For  $w \in H^1(\mathbb{R}^d; \mathbb{R}^d)$  representing prescribed Dirichlet condition  $w|_{\Gamma_D}$ , we define

$$A(w) := \left\{ (u, \pi) \in \text{BD}(\Omega; \mathbb{R}^d) \times \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}) \mid e(u) - \pi|_{\Omega} \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}), \right. \\ \left. \pi|_{\Gamma_D} = (w - u) \otimes \nu \, d\mathcal{H}^{d-1} \text{ on } \Gamma_D \right\}. \quad (3.3)$$

Lemma 2.1 in [11] shows that  $A : w \mapsto A(w)$  has a weak  $\times$  weak\*-closed graph in  $H^1(\mathbb{R}^d; \mathbb{R}^d) \times (\text{BD}(\Omega; \mathbb{R}^d) \times \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}))$ . This shows that  $Q_{\text{PR}}$  from (3.2b), being equal to  $A(0)$ , is a closed subset of the Banach space of

$$\tilde{Q}_{\text{PR}} := \left\{ (u, \pi) \in \text{BD}(\Omega; \mathbb{R}^d) \times \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}) \mid e(u) - \pi|_{\Omega} \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}) \right\}$$

which is equipped with the norm

$$\|(u, \pi)\|_{\tilde{Q}_{\text{PR}}} := \|u\|_{L^1(\Omega)} + \|e(u)\|_{\mathcal{M}(\Omega)} + \|\pi\|_{\mathcal{M}(\Omega \cup \Gamma_D)} + \|e(u) - \pi\|_{L^2(\Omega)}.$$

Thus,  $Q_{\text{PR}}$  is itself a Banach space with the norm  $\|\cdot\|_{\tilde{Q}_{\text{PR}}}$ .

In view of (3.2b), for any energetic solution  $q \in L^\infty(I; Q)$  the elastic stress  $\sigma = \mathbb{C}(e(u + u_D) - \pi)$  lies in  $L^\infty(I; L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}))$  even though its particular components  $\mathbb{C}e(u)$  and  $\mathbb{C}\pi$  may exhibit spatial concentration. Note also that, comparing to (2.6e), we now consider  $f = 0$  and  $g = 0$  to prevent uncontrolled slip and thus blow-up of a-priori estimates. In fact, certain qualified nonvanishing  $f = 0$  or  $g = 0$  could be admitted if a suitable ‘‘safe load condition’’ is additionally imposed, cf. [11, Formulas (2.17)-(2.18)].

Existence of solution to the Prandtl-Reuss system has been proved in [25, 53] by using the method of vanishing viscosity (or Yosida regularization). In [14], the limit passage from the rate-independent incremental problem was done while using weaker notion of solutions. Only recently, in [11, Thm.4.5], Dal Maso et al. executed this limit procedure in the framework of energetic solutions. In [17, Sect.9], it was shown the Prandtl-Reuss limit to also be reached from a rate-independent model of gradient plasticity.

We give a slight variant of the existence proof from [11] which is based on quite different arguments based on abstract BV relaxations. This variant follows [48] but still relies on the technical results from [11]. However, we hope that it can be generalized more easily to other plasticity models. We introduce an intermediate notion of convergence in  $Q_{\text{PR}}$  denoted by ‘‘ $\xrightarrow{\text{strict}}$ ’’ and defined via

$$(u_n, \pi_n) \xrightarrow{\text{strict}} (u, \pi) \iff \begin{cases} (u_n, \pi_n) \xrightarrow{*} (u, \pi) & \text{in } \text{BV}(\Omega; \mathbb{R}^d) \times \mathcal{M}(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}), \\ |\pi_n|(\Omega \cup \Gamma_D) \rightarrow |\pi|(\Omega \cup \Gamma_D) & \text{in } \mathbb{R}, \\ e(u_n) - \pi_n|_{\Omega} \rightarrow e(u) - \pi|_{\Omega} & \text{in } L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}). \end{cases} \quad (3.4)$$

Moreover, we define the subspace

$$Q_{\text{PR}}^0 := \left\{ (u, \pi) \in W^{1,1}(\Omega; \mathbb{R}^d) \times L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d}) \mid u = 0 \text{ on } \Gamma_D, \right. \\ \left. e(u) - \pi \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}) \right\}. \quad (3.5)$$

Obviously,  $Q_{\text{PR}}^0$  is a strongly closed subspace of  $Q_{\text{PR}}$ ; in fact it is the strong closure of  $\mathbb{P}Q$  for  $Q$  from (2.6b), where  $\mathbb{P}(u, \pi, \eta) = (u, \pi)$  is the projection dropping the isotropic hardening parameter.



Our different approach is based on the following central conjecture, which states that a weakened version weak stability is sufficient to imply the desired stability. We believe that the implication (3.6) is true under quite general conditions. We will show that it holds at least in the case considered in [11], see Remark 3.3.

**Conjecture 3.1.** *For all  $t \in [0, T]$  and all  $(u, \pi) \in Q_{\text{PR}}$  the following holds:*

$$\left. \begin{array}{l} \text{If } \mathcal{E}_{\text{PR}}(t, u, \pi) \leq \mathcal{E}_{\text{PR}}(t, u + \hat{u}, \pi + \hat{\pi}) + \mathcal{R}_{\text{PR}}(\hat{\pi}) \text{ for all } (\hat{u}, \hat{\pi}) \in Q_{\text{PR}}^0, \\ \text{then } \mathcal{E}_{\text{PR}}(t, u, \pi) \leq \mathcal{E}_{\text{PR}}(t, u + \tilde{u}, \pi + \tilde{\pi}) + \mathcal{R}_{\text{PR}}(\tilde{\pi}) \text{ for all } (\tilde{u}, \tilde{\pi}) \in Q_{\text{PR}}. \end{array} \right\} \quad (3.6)$$

The following arguments (from [48]) support the validity of (3.6). First one knows by [4, Thm. 2.38] that

$$\mathcal{J} : (\tilde{u}, \tilde{\pi}) \mapsto \mathcal{E}_{\text{PR}}(t, u + \tilde{u}, \pi + \tilde{\pi}) + \mathcal{R}_{\text{PR}}(\tilde{\pi})$$

is sequentially lower semicontinuous on  $Q_{\text{PR}}$  with respect to the weak\* convergence. Thus, the minimum over  $(\tilde{u}, \tilde{\pi}) \in Q_{\text{PR}}$  on the right-hand side of the lower line in (3.6) is achieved. Moreover, [4, Thm. 2.39] guarantees that  $\mathcal{J}$  is even sequentially continuous with respect to the strict convergence (3.4) on  $Q_{\text{PR}}$ . Under suitable conditions on  $\partial\Omega$  and  $\Gamma_{\text{D}}$  it should be possible to show that  $Q_{\text{PR}}^0$  is dense in  $Q_{\text{PR}}$  with respect to the strict convergence, see [48]. Then, (3.6) follows by continuity and density.

As a side effect of the vanishing-hardening convergence below, we obtain the following existence result.

**Proposition 3.2** (Energetic solution for the Prandtl-Reuss model). *Let the rate-independent system  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$  be as described in (3.2) such that (3.6) holds. Moreover, assume  $u_{\text{D}} \in W^{1,1}(I; W^{1,2}(\Omega; \mathbb{R}^d))$  and that  $(u_0, \pi_0) \in Q_{\text{PR}}$  is stable at  $t = 0$ . Then, there exists an energetic solution  $(u, \pi)$  with  $(u(0), \pi(0)) = (u_0, \pi_0)$ . Moreover,  $\mathbb{C}(e(u) - \pi)$  and thus also the “true” elastic stress  $\sigma$  from (1.3) is determined uniquely.*

*Proof.* It suffices to merge Proposition 2.1 with Theorem 4.1 below for  $\eta_0 \in L^2(\Omega)$  such that  $\delta_S(\pi_0, \eta_0) = \delta_P(\pi_0)$  and realize that (4.1) can always be satisfied for a suitable  $S$ , e.g. for that one from Example 4.2.  $\square$

**Remark 3.3.** Conjecture 3.1 holds in the case considered in [11]. The equivalence proved in [11, Prop. 3.5] was used for exactly the same purpose as we use (3.6), namely to show that the set of stable states is closed under the weak\* convergence, see [11, Thm. 3.7]. There the following additional conditions were used:

$$\Omega \text{ has a } (d-1) \text{ dimensional } C^2 \text{ boundary,} \quad (3.7a)$$

$$\Gamma_{\text{D}} \text{ has a } (d-2) \text{ dimensional } C^2 \text{ boundary,} \quad (3.7b)$$

$$\mathbb{C}e = \mathbb{C}_{\text{D}} \left( e - \frac{\text{tr } e}{d} \mathbb{I} \right) + \kappa (\text{tr } e) \mathbb{I} \quad \text{with}$$

$$\mathbb{C}_{\text{D}} : \mathbb{R}_{\text{dev}}^{d \times d} \rightarrow \mathbb{R}_{\text{dev}}^{d \times d} \text{ positive definite and } \kappa > 0. \quad (3.7c)$$

The proof of (3.6) for this case is obtained in [11, Sect. 3.2] as a result of a subtle regularity theory for the stress  $\sigma = \mathbb{C}(e(u) - \pi) \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$  (namely  $\text{dev } \sigma \in L^\infty(\Omega)$  and  $\text{div } \sigma = 0$  in  $\Omega$  and  $\sigma \nu = 0$  on  $\Gamma_{\text{N}} = \partial\Omega \setminus \Gamma_{\text{D}}$ ) and a careful analysis of the stress-strain duality, see e.g. [11, Prop. 2.4].

In fact, in [11], the stronger hypotheses  $f_{\text{ext}} \in W^{1,1}(I; L^d(\Omega; \mathbb{R}^d))$  are assumed, which would lead here to the qualification  $u_{\text{D}} \in W^{1,1}(I; W^{2-1/d, d}(\Gamma_{\text{D}}; \mathbb{R}^d))$  that allows the extension of  $u_{\text{D}}$  in  $W^{1,1}(I; W^{2, d}(\Omega; \mathbb{R}^d))$  so that  $\text{div } \mathbb{C}e(u_{\text{D}})$  is in  $W^{1,1}(I; L^d(\Omega; \mathbb{R}^d))$ . Yet, one should realize a special character of the Dirichlet part

of the load  $f_{\text{ext}}$  from (2.6e) that needs only  $e(u_{\text{D}}) \in W^{1,1}(I; L^2(\Omega; \mathbb{R}^{d \times d}))$  when tested by  $e(u) - \pi \in L^2(\Omega; \mathbb{R}^{d \times d})$ .

#### 4. THE LIMIT OF VANISHING HARDENING

Now we present and prove the main result, namely that the time-dependent Prandtl-Reuss model can be obtained as a limit from the classical elastoplastic model with hardening, if the hardening approaches zero. Most previous existence results for the Prandtl-Reuss system are based on viscous regularizations of the flow rule, see e.g. [5, 14, 53]. The vanishing-hardening limit was investigated only in the context of the dynamical problem in [9, 10], where the inertial effects can be used to prevent instantaneous formation of shear bands. To the best of the authors knowledge, the limit of vanishing hardening in the strict rate-independent setting was not investigated so far. And it is exactly this limit, which can be combined easily with numerical space-time discretizations, see the following section.

We naturally need some moderate qualifications for the relation between the dissipative potentials  $\delta_S^*$  and  $\delta_P^*$  occurring in (2.6d) and (3.2d), respectively. We suppose

$$\forall \pi \in \mathbb{R}^{d \times d}_{\text{dev}} : \quad \delta_P^*(\pi) = \min_{\eta \in \mathbb{R}} \delta_S^*(\pi, \eta), \quad (4.1a)$$

$$\exists C \in \mathbb{R} \quad \forall \pi \in \mathbb{R}^{d \times d}_{\text{dev}} \quad \exists \eta \in \mathbb{R} : \quad \delta_S^*(\pi, \eta) = \delta_P^*(\pi) \quad \text{and} \quad |\eta| \leq C|\pi|. \quad (4.1b)$$

We show in Example 4.2 that these conditions hold in the standard cases used for modeling hardening.

**Theorem 4.1** (Convergence of vanishing hardening). *Let all the assumptions of Proposition 2.1 and (3.6) hold. Define*

$$\mathcal{E}_\varepsilon(t, u, \pi, \eta) := \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u + 2u_{\text{D}}(t)) - \pi) : (e(u) - \pi) + \frac{\varepsilon}{2} \mathbb{H}\pi : \pi + \frac{\varepsilon}{2} b\eta^2 \, dx, \quad (4.2)$$

assume that  $q_0 = (u_0, \pi_0, \eta_0) \in Q$  is stable at  $t = 0$  for each  $\varepsilon > 0$ , and denote by  $q_\varepsilon = (u_\varepsilon, \pi_\varepsilon, \eta_\varepsilon)$  be the unique energetic solution to  $(Q, \mathcal{E}_\varepsilon, \mathcal{R})$ ; cf. (2.6). Then the following holds:

(i) *There is a constant  $C$  independent of  $\varepsilon$  giving the a-priori estimates*

$$\|\pi_\varepsilon\|_{\text{BV}(\bar{I}; L^1(\Omega; \mathbb{R}^{d \times d}_{\text{dev}}))} \leq C, \quad (4.3a)$$

$$\|e(u_\varepsilon) - \pi_\varepsilon\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}_{\text{sym}}))} \leq C, \quad (4.3b)$$

$$\|e(u_\varepsilon)\|_{L^\infty(I; L^1(\Omega; \mathbb{R}^{d \times d}_{\text{sym}}))} \leq C, \quad (4.3c)$$

$$\|\pi_\varepsilon\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}_{\text{dev}}))} \leq C/\sqrt{\varepsilon}, \quad (4.3d)$$

$$\|e(u_\varepsilon)\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}_{\text{sym}}))} \leq C/\sqrt{\varepsilon}, \quad (4.3e)$$

$$\|\eta_\varepsilon\|_{L^\infty(I; L^2(\Omega))} \leq C/\sqrt{\varepsilon}. \quad (4.3f)$$

(ii) *For  $\varepsilon \rightarrow 0$ , there is a subsequence of  $\{(u_\varepsilon, \pi_\varepsilon)\}_{\varepsilon > 0}$  converging weakly\* in  $L^\infty(I; \text{BD}(\Omega; \mathbb{R}^d)) \times \text{BV}(\bar{I}; \mathcal{M}(\Omega \cup \Gamma_{\text{D}}; \mathbb{R}^{d \times d}_{\text{dev}}))$  to some  $(u, \pi)$  and also  $\pi_\varepsilon(t) \overset{*}{\rightharpoonup} \pi(t)$  in  $\mathcal{M}(\Omega \cup \Gamma_{\text{D}}; \mathbb{R}^{d \times d}_{\text{dev}})$  for any  $t \in \bar{I}$ . Every such couple  $(u, \pi)$  is an energetic solution to  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}})$  with  $(u(0), \pi(0)) = (u_0, \pi_0)$ ; see (1.2) and (3.2).*

(iii) *Moreover, the whole sequence of stresses  $\{\mathbb{C}(e(u_\varepsilon) - \pi_\varepsilon)\}_{\varepsilon > 0}$  converges weakly\* in  $L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}_{\text{sym}}))$  to  $\mathbb{C}(e(u) - \pi)$  and even strongly pointwise on  $[0, T]$ , i.e.*

$$\forall t \in [0, T] : \quad \mathbb{C}(e(u_\varepsilon(t)) - \pi_\varepsilon(t)) \rightarrow \mathbb{C}(e(u(t)) - \pi(t)|_{\Omega}) \quad \text{in } L^2(\Omega; \mathbb{R}^{d \times d}_{\text{sym}}), \quad (4.4)$$

and analogous assertion holds for “true” stresses from (1.3).

Let us note that (4.3c) cannot yield any uniform  $L^\infty(I; W^{1,1}(\Omega; \mathbb{R}^d))$ -estimate for  $u_\varepsilon$  because Korn’s inequality does not hold for this limit case, cf. [43].

An example for stable initial conditions that are independent of  $\varepsilon$  is  $\pi_0 = 0$  and  $\eta_0 = 0$  and  $u_0$  minimizing  $\mathcal{E}_\varepsilon(t, \cdot, 0, 0)$  which obviously does not depend on  $\varepsilon$ ; note that  $\delta_S(0, 0) = \delta_P(0)$ . In principle, one may weaken the requirement of stability of initial conditions independently of  $\varepsilon$  and another approximation would allow for  $u_0 \in W^{1,1}(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  and  $\pi_0 \in L^1(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$ . For this we would need to approximate both  $u_0$  and  $\pi_0$  by some  $u_{0\varepsilon}$  and  $\pi_{0\varepsilon} \in L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  so that  $e(u_{0\varepsilon}) - \pi_{0\varepsilon} \rightarrow e(u_0) - \pi_0$  in  $L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})$  and  $\|\pi_{0\varepsilon}\|_{L^2(\Omega; \mathbb{R}_{\text{dev}}^{d \times d})} = o(\varepsilon^{-1/2})$  so that  $\int_\Omega \varepsilon \mathbb{H} \pi_{0\varepsilon} : \pi_{0\varepsilon} dx \rightarrow 0$ .

*Proof of Theorem 4.1.* We divide the proof to the six steps.

*Step 1 (a-priori estimates):* The coercivity of  $\mathcal{R}$  and of the “ $\mathbb{C}$ -part” of  $\mathcal{E}_\varepsilon$  then gives the estimates (4.3a,b). By the obvious inequality  $\|e(u)\|_{L^1(\Omega; \mathbb{R}^{d \times d})} \leq (\text{meas}^d(\Omega))^{1/2} \|e(u) - \pi\|_{L^2(\Omega; \mathbb{R}^{d \times d})} + \|\pi\|_{L^1(\Omega; \mathbb{R}^{d \times d})}$ , we obtain also (4.3c). From the “ $\mathbb{H}$ -part” of  $\mathcal{E}_\varepsilon$  and from  $\frac{1}{2}\varepsilon\eta^2$ , we further obtain (4.3d) and (4.3f), respectively. Then, by the triangle inequality  $\|e(u)\|_{L^2(\Omega; \mathbb{R}^{d \times d})} \leq \|e(u) - \pi\|_{L^2(\Omega; \mathbb{R}^{d \times d})} + \|\pi\|_{L^2(\Omega; \mathbb{R}^{d \times d})}$ , we eventually obtain (4.3e).

*Step 2 (selection of convergent subsequences):* Let us take the energetic solution  $(u_\varepsilon, \pi_\varepsilon, \eta_\varepsilon)$  to the problem  $(U \times Z, \mathcal{E}_\varepsilon, \mathcal{R}, u_0, \pi_0, \eta_0)$ , which exists due to Proposition 2.1. For the convergence we use the estimates (4.3a-c) and select the subsequence by Banach’s selection principle. Moreover, by Helly’s selection principle and metrizable of the weak\* topology on the balls in  $\mathcal{M}(\Omega \cup \Gamma_D)$ , we can also claim  $\pi_\varepsilon(t) \xrightarrow{*} \pi(t)$  in  $\mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d})$  for all  $t \in \bar{I}$ . Then also

$$\sigma_\varepsilon(t) = \mathbb{C}(e(u_\varepsilon(t)) - \pi_\varepsilon(t)) \rightharpoonup \sigma(t) = \mathbb{C}(e(u(t)) - \pi(t)|_\Omega) \quad \text{in } L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}) \quad (4.5)$$

for all  $t \in \bar{I}$ ; it follows from the a-priori estimate (4.3b) and the uniqueness of the stress, cf. the arguments in [34, Sect.4.2.3] or [11, Thm.5.9]. Also, for any  $t \in \bar{I}$ ,  $u_\varepsilon(t)$  converges weakly\* in  $\text{BD}(\Omega; \mathbb{R}^d)$  to a limit which can be uniquely identified with  $u(t)$  because its strain  $e(u(t)) = \mathbb{C}^{-1}\sigma(t) + \pi(t)|_\Omega \in \mathcal{M}(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$  has already been determined uniquely when choosing  $\pi_\varepsilon(t) \xrightarrow{*} \pi(t)$  in  $\mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d})$  and because of the prescribed Dirichlet boundary conditions.

*Step 3 (stability):* The convergence in the stability condition (1.2a) requires essentially an explicit construction of *mutual recovery sequences* in the spirit of [38]. Here, because of the disappearance of the isotropic hardening in the limit, we must consider a modification of this concept. For each  $t$  the solutions  $q_\varepsilon(t) = (u_\varepsilon(t), \pi_\varepsilon(t), \eta_\varepsilon(t))$  form a stable sequence, i.e.  $\sup_{\varepsilon > 0} \mathcal{E}_\varepsilon(t, q_\varepsilon(t)) < \infty$  and  $q_\varepsilon(t) \in \mathcal{S}_\varepsilon(t)$  with  $\mathcal{S}_\varepsilon(t)$  from (2.7) with  $\mathcal{E}_\varepsilon$  in place of  $\mathcal{E}$ , such that  $(u_\varepsilon, \pi_\varepsilon) \xrightarrow{*} (u(t), \pi(t)) \in Q_{\text{PR}}$ , i.e.

$$u_\varepsilon(t) \xrightarrow{*} u(t) \quad \text{in } \text{BD}(\Omega; \mathbb{R}^d), \quad (4.6a)$$

$$\pi_\varepsilon(t) \xrightarrow{*} \pi(t) \quad \text{in } \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}), \quad (4.6b)$$

$$e(u_\varepsilon(t)) - \pi_\varepsilon(t) \rightharpoonup e(u(t)) - \pi(t) \quad \text{in } L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}). \quad (4.6c)$$

In particular, this implies  $\|\pi_\varepsilon(t)\|_{L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})} = \mathcal{O}(\varepsilon^{-1/2})$  and  $\|\eta_\varepsilon(t)\|_{L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})} = \mathcal{O}(\varepsilon^{-1/2})$ .

Now, for fixed  $t$  and a given test function  $(\tilde{u}, \tilde{\pi}) \in Q_{\text{PR}}$  we have to construct a so-called mutual recovery sequence  $(\tilde{q}_\varepsilon)_{\varepsilon>0}$  in the spirit of [38] such that

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(t, \tilde{q}_\varepsilon) + \mathcal{R}(\tilde{q}_\varepsilon - q_\varepsilon(t)) - \mathcal{E}_\varepsilon(t, q_\varepsilon(t)) \\ \leq \mathcal{E}_{\text{PR}}(\tilde{u}, \tilde{\pi}) + \mathcal{R}_{\text{PR}}(\tilde{\pi} - \pi(t)) - \mathcal{E}_{\text{PR}}(t, u, \pi(t)). \end{aligned} \quad (4.7)$$

The main point is that we first restrict ourselves to special test functions  $(\tilde{u}, \tilde{\pi}) \in Q_{\text{PR}}$ , namely those given in the form (where we temporarily drop the time dependence)

$$\begin{aligned} (\tilde{u}, \tilde{\pi}, \tilde{\eta}) &= (u, \pi, 0) + (\hat{u}, \hat{\pi}, \hat{\eta}) \\ \text{with } (\hat{u}, \hat{\pi}) &\in Q_{\text{PR}}^1 := \text{H}_{\Gamma_D}^1(\Omega; \mathbb{R}^d) \times \text{L}^2(\Omega; \mathbb{R}^{d \times d}), \\ \text{and } \hat{\eta} &\in \text{L}^2(\Omega) \text{ such that } \delta_S^*(\hat{\pi}, \hat{\eta}) = \delta_P^*(\hat{\pi}), \end{aligned} \quad (4.8)$$

where  $\text{H}_{\Gamma_D}^1(\Omega; \mathbb{R}^d) := U$  from (2.6a). Then we put

$$\tilde{q}_\varepsilon = (\tilde{u}_\varepsilon, \tilde{\pi}_\varepsilon, \tilde{\eta}_\varepsilon) = (u_\varepsilon + \tilde{u} - u, \pi_\varepsilon + \tilde{\pi} - \pi, \eta_\varepsilon + \tilde{\eta} - 0); \quad (4.9)$$

note that the difference  $\tilde{q}_\varepsilon - q_\varepsilon = (\tilde{u} - u, \tilde{\pi} - \pi, \tilde{\eta}) = \hat{q}$  is independent of  $\varepsilon$ . Hence,

$$\begin{aligned} \mathcal{R}(\tilde{\pi}_\varepsilon - \pi_\varepsilon, \tilde{\eta}_\varepsilon - \eta_\varepsilon) &= \int_{\Omega} \delta_S^*(\tilde{\pi}_\varepsilon - \pi_\varepsilon, \tilde{\eta}_\varepsilon - \eta_\varepsilon) \, dx = \int_{\Omega} \delta_S^*(\tilde{\pi} - \pi, \tilde{\eta}) \, dx \\ &= \int_{\Omega} \delta_S^*(\hat{\pi}, \hat{\eta}) \, dx = \int_{\Omega} \delta_P^*(\hat{\pi}) \, dx = \mathcal{R}_{\text{PR}}(\tilde{\pi} - \pi). \end{aligned} \quad (4.10)$$

Thus, we have the middle term in the limsup in (4.7) under control.

To control the limit of the two energy terms in the limsup (4.7) we use a cancellation property of the quadratic functionals  $\mathcal{E}_\varepsilon$  and  $\mathcal{E}_{\text{PR}}$ : because of the quadratic structure of both  $\mathcal{E}_\varepsilon$  and  $\mathcal{E}_{\text{PR}}$ , for  $f_{\text{ext}}$  defined again by (2.6e) now with  $f = 0$  and  $g = 0$ , we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(t, \tilde{u}_\varepsilon, \tilde{\pi}_\varepsilon, \tilde{\eta}_\varepsilon) - \mathcal{E}_\varepsilon(t, u_\varepsilon, \pi_\varepsilon, \eta_\varepsilon) \\ = \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\tilde{u}_\varepsilon - u_\varepsilon) - \tilde{\pi}_\varepsilon + \pi_\varepsilon) : (e(\tilde{u}_\varepsilon + u_\varepsilon) - \tilde{\pi}_\varepsilon - \pi_\varepsilon) \\ + \frac{\varepsilon}{2} \mathbb{H}(\tilde{\pi}_\varepsilon - \pi_\varepsilon) : (\tilde{\pi}_\varepsilon + \pi_\varepsilon) + \frac{\varepsilon}{2} (\tilde{\eta}_\varepsilon - \eta_\varepsilon)(\tilde{\eta}_\varepsilon + \eta_\varepsilon) \, dx - \langle f_{\text{ext}}(t), (\tilde{u}_\varepsilon - u_\varepsilon, \tilde{\pi}_\varepsilon - \pi_\varepsilon) \rangle \\ = \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\hat{u}) - \hat{\pi}) : (e(2u_\varepsilon + \hat{u}) - 2\pi_\varepsilon - \hat{\pi}) + \frac{\varepsilon}{2} \mathbb{H}\hat{\pi} : (2\pi_\varepsilon + \hat{\pi}) \\ + \frac{\varepsilon}{2} \tilde{\eta}(\tilde{\eta}_\varepsilon + \eta_\varepsilon) \, dx - \langle f_{\text{ext}}(t), (\hat{u}, \hat{\pi}) \rangle \\ = \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\hat{u}) - \hat{\pi}) : (e(2u + \hat{u}) - 2\pi - \hat{\pi}) + \, dx - \langle f_{\text{ext}}(t), (\hat{u}, \hat{\pi}) \rangle \\ = \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\tilde{u} - u) - \tilde{\pi} + \pi) : (e(\tilde{u} + u) - \tilde{\pi} - \pi) \, dx - \langle f_{\text{ext}}(t), (\tilde{u} - u, \tilde{\pi} - \pi) \rangle \\ = \mathcal{E}_{\text{PR}}(t, \tilde{u}, \tilde{\pi}) - \mathcal{E}_{\text{PR}}(t, u, \pi), \end{aligned} \quad (4.11)$$

where we used (4.6c)

$$\begin{aligned} \left| \int_{\Omega} \frac{\varepsilon}{2} \mathbb{H}\hat{\pi} : (2\pi_\varepsilon + \hat{\pi}) \, dx \right| &\leq \frac{\varepsilon}{2} |\mathbb{H}| \|\hat{\pi}\|_{\text{L}^2(\Omega)} \|2\pi_\varepsilon + \hat{\pi}\|_{\text{L}^2(\Omega)} = \mathcal{O}(\sqrt{\varepsilon}) \rightarrow 0 \quad \text{and} \\ \left| \int_{\Omega} \frac{\varepsilon}{2} b\tilde{\eta}(\tilde{\eta}_\varepsilon + \eta_\varepsilon) \, dx \right| &\leq \frac{\varepsilon}{2} b \|\hat{\eta}\|_{\text{L}^2(\Omega)} \|\tilde{\eta}_\varepsilon + \eta_\varepsilon\|_{\text{L}^2(\Omega)} = \mathcal{O}(\sqrt{\varepsilon}) \rightarrow 0. \end{aligned} \quad (4.12)$$

This allows for the limit passage in the mutual recovery sequence condition (4.7). Using the stability of  $q_\varepsilon$  we know that each term in the limsup is non-negative. Hence we conclude that the right-hand side is non-negative as well. Recalling the special choice of  $(\tilde{u}, \tilde{\pi})$  in (4.8) we have derived the following stability condition for the limit  $(u(t), \pi(t)) \in Q_{\text{PR}}$ :

$$\forall(\hat{u}, \hat{\pi}) \in Q_{\text{PR}}^1 : \quad \mathcal{E}_{\text{PR}}(t, u(t), \pi(t)) \leq \mathcal{E}_{\text{PR}}(t, u(t)+\hat{u}, \pi(t)+\hat{\pi}) + \mathcal{R}_{\text{PR}}(\hat{\pi}), \quad (4.13)$$

where  $Q_{\text{PR}}^1$  is defined in (4.8). Since  $Q_{\text{PR}}^0$  is the (strong) closure of  $Q_{\text{PR}}^1$  in  $Q_{\text{PR}}$  and  $\mathcal{E}_{\text{PR}}$  and  $\mathcal{R}_{\text{PR}}$  are continuous, we have shown that  $(u(t), \pi(t))$  satisfies the upper condition in the assumed property (3.6). Thus, the lower condition holds as well and the desired stability (S) in (1.2a) is established.

*Step 4 (upper energy balance):* Using the energy balance (1.2b) for  $q_\varepsilon$ , for  $f_{\text{ext}}$  defined again by (2.6e) now with  $f = 0$  and  $g = 0$ , we have

$$\begin{aligned} & \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u_\varepsilon(t)) - \pi_\varepsilon(t)) : (e(u_\varepsilon(t)) - \pi_\varepsilon(t)) + \frac{\varepsilon}{2} \mathbb{H} \pi_\varepsilon(t) : \pi_\varepsilon(t) + \frac{\varepsilon}{2} b \eta_\varepsilon(t)^2 dx \\ & + \text{Diss}_{\mathcal{R}}(\pi_\varepsilon, \eta_\varepsilon; [0, t]) - \langle f_{\text{ext}}(t), (u_\varepsilon(t), \pi_\varepsilon(t)) \rangle \leq \int_0^t \langle \dot{f}_{\text{ext}}, (u_\varepsilon, \pi_\varepsilon) \rangle dt \\ & + \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u_0) - \pi_0) : (e(u_0) - \pi_0) + \frac{\varepsilon}{2} \mathbb{H} \pi_0 : \pi_0 + \frac{\varepsilon}{2} b \eta_0^2 dx - \langle f_{\text{ext}}(0), (u_0, \pi_0) \rangle. \end{aligned} \quad (4.14)$$

Note that  $e(u_\varepsilon(t)) - \pi_\varepsilon(t)$  is well defined because of the bound (4.3a) and because  $u_\varepsilon(t)$  is then determined uniquely by minimizing  $\mathcal{E}_\varepsilon(t, \cdot, \pi_\varepsilon(t), \eta_\varepsilon(t))$ . We can pass to the limit in (4.14) by using weak lower semicontinuity of  $\mathcal{E}_{\text{PR}}(t, \cdot, \cdot)$  and that the already mentioned convergence  $e(u_\varepsilon(t)) - \pi_\varepsilon(t) \rightharpoonup e(u(t)) - \pi(t)$  in  $L^2(\Omega; \mathbb{R}^{d \times d})$ , and also by using

$$\liminf_{\varepsilon \rightarrow 0} \text{Diss}_{\mathcal{R}}(\pi_\varepsilon, \eta_\varepsilon; [0, t]) \geq \liminf_{\varepsilon \rightarrow 0} \text{Diss}_{\mathcal{R}_{\text{PR}}}(\pi_\varepsilon; [0, t]) \geq \text{Diss}_{\mathcal{R}_{\text{PR}}}(\pi; [0, t]). \quad (4.15)$$

The first inequality in (4.15) is just by  $\delta_S^*(\pi, \eta) \geq \delta_P^*(\pi)$ , while the second one is by weak lower-semi continuity of  $\text{Diss}_{\mathcal{R}_{\text{PR}}}(\cdot; [0, t])$ . Eventually, we can forget the terms  $\int_{\Omega} \frac{\varepsilon}{2} \mathbb{H} \pi_\varepsilon(t) : \pi_\varepsilon(t) + \frac{\varepsilon}{2} \eta_\varepsilon(t)^2 dx$  while  $\int_{\Omega} \frac{\varepsilon}{2} \mathbb{H} \pi_0 : \pi_0 dx$  converges to 0 because we have assumed  $\pi_0 \in L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$  and also  $\int_{\Omega} \frac{\varepsilon}{2} b \eta_0^2 dx \rightarrow 0$  by  $\eta_0 \in L^2(\Omega)$ . The limit passage in (4.14) results to the upper energy estimate

$$\mathcal{E}_{\text{PR}}(t, u(t), \pi(t)) + \text{Diss}_{\mathcal{R}_{\text{PR}}}(\pi; [0, t]) \leq \mathcal{E}_{\text{PR}}(0, u_0, \pi_0) + \int_0^t \langle \dot{f}_{\text{ext}}, (u, \pi) \rangle dt. \quad (4.16)$$

*Step 5 (lower energy balance):* Recalling the upper energy estimate (4.16) we obtain the desired energy balance (E) in (1.2b), by using the standard argument that stability of  $q(t)$  for all  $t \in \bar{I}$  implies the lower energy inequality, cf. [33, 42].

*Step 6 (improved convergence):* The convergence of the whole sequence  $(\mathbb{C}e(u_\varepsilon) - \pi_\varepsilon)_{\varepsilon > 0}$  is due to the mentioned uniqueness of the elastic stresses. Having the pointwise stability, by abstract arguments from theory of rate-independent processes, the stored energy converges pointwise, i.e.  $\mathcal{E}_\varepsilon(t, u_\varepsilon(t), \pi_\varepsilon(t), \eta_\varepsilon(t)) \rightarrow \mathcal{E}_{\text{PR}}(t, u(t), \pi(t))$  for

any  $t \in [0, T]$ . In view of (4.2) and (3.2c) and using also (4.5), we have

$$\begin{aligned}
& \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u_{\varepsilon}(t)) - \pi_{\varepsilon}(t)) : (e(u_{\varepsilon}(t)) - \pi_{\varepsilon}(t)) \, dx \\
& \leq \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u_{\varepsilon}(t)) - \pi_{\varepsilon}(t)) : (e(u_{\varepsilon}(t)) - \pi_{\varepsilon}(t)) + \frac{\varepsilon}{2} \mathbb{H} \pi_{\varepsilon}(t) : \pi_{\varepsilon}(t) + \frac{\varepsilon}{2} b \eta_{\varepsilon}(t)^2 \, dx \\
& = \lim_{\varepsilon \rightarrow 0} \mathcal{E}_{\varepsilon}(t, u_{\varepsilon}(t), \pi_{\varepsilon}(t), \eta_{\varepsilon}(t)) - \int_{\Omega} \sigma_{\varepsilon}(t) : e(u_{\text{D}}(t)) \, dx \\
& = \mathcal{E}_{\text{PR}}(t, u(t), \pi(t)) - \int_{\Omega} \sigma(t) : e(u_{\text{D}}(t)) \, dx \\
& = \int_{\Omega} \frac{1}{2} \mathbb{C}(e(u(t)) - \pi(t)) : (e(u(t)) - \pi(t)) \, dx. \tag{4.17}
\end{aligned}$$

Thus, by positive definiteness of  $\mathbb{C}$  and using an appropriate equivalent norm on the Hilbert space  $L^2(\Omega; \mathbb{R}^{d \times d})$  induced by  $\mathbb{C}$ , we obtain  $\liminf_{\varepsilon \rightarrow 0} \|e(u_{\varepsilon}(t)) - \pi_{\varepsilon}(t)\|_{L^2(\Omega; \mathbb{R}^{d \times d})} \leq \|e(u(t)) - \pi(t)\|_{L^2(\Omega; \mathbb{R}^{d \times d})}$ . From this and from (4.5), the strong convergence (4.4) follows by the uniform convexity of the Hilbert space  $L^2(\Omega; \mathbb{R}^{d \times d})$ .  $\square$

**Example 4.2** (Dissipation potential). An example for the dissipation potential, considered e.g. in [21, 22, 35, 51] (often with for  $q_{\text{H}} = 1$ ), is

$$\delta_S^*(\dot{\pi}, \dot{\eta}) := \delta_P^*(\dot{\pi}) + \delta_K(\dot{\pi}, \dot{\eta}) \quad \text{with} \quad K := \left\{ (\dot{\pi}, \dot{\eta}) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R} \mid \dot{\eta} \leq -q_{\text{H}} \delta_P^*(\dot{\pi}) \right\} \tag{4.18}$$

for  $P \subset \mathbb{R}_{\text{dev}}^{d \times d}$  a convex, bounded, and closed neighbourhood of the origin and  $q_{\text{H}} > 0$ . We want to identify  $S$  via the definition

$$S = \partial \delta_S^*(0) = \left\{ (\varsigma, \xi) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R} \mid \forall (\dot{\pi}, \dot{\eta}) : \dot{\eta} \leq -q_{\text{H}} \delta_P^*(\dot{\pi}) \Rightarrow \delta_P^*(\dot{\pi}) \geq \dot{\pi} : \varsigma + \dot{\eta} \xi \right\}.$$

As  $\delta_P^*(0)$  is finite, any  $(\varsigma, \xi) \in S$  satisfies implies  $\xi \geq 0$ . Then the maximal value of  $\dot{\pi} : \varsigma + \dot{\eta} \xi$  is attained at  $\dot{\eta} = -q_{\text{H}} \delta_P^*(\dot{\pi})$  and thus equals to  $\dot{\pi} : \varsigma - q_{\text{H}} \delta_P^*(\dot{\pi}) \xi$ . The inequality  $\delta_P^*(\dot{\pi}) \geq \dot{\pi} : \varsigma + \dot{\eta} \xi$  then reduces to  $\dot{\pi} : \varsigma \leq (1 + q_{\text{H}} \xi) \delta_P^*(\dot{\pi})$  for all  $\dot{\pi} \in \mathbb{R}_{\text{dev}}^{d \times d}$ . By the 1-homogeneity of  $\delta_P^*$ , this gives  $\varsigma \in (1 + q_{\text{H}} \xi) \partial \delta_P^*(0)$ . Using  $\partial \delta_P^*(0) = P$ , we obtain the characterization

$$S = \left\{ (\varsigma, \xi) \in \mathbb{R}_{\text{dev}}^{d \times d} \times \mathbb{R} \mid \xi \geq 0 \quad \text{and} \quad \varsigma \in (1 + q_{\text{H}} \xi) P \right\}. \tag{4.19}$$

Note that the constraint  $\dot{\eta} \leq -q_{\text{H}} \delta_P^*(\dot{\pi}) \leq 0$  in (4.18) makes the evolution of the hardening  $\eta$  unidirectional, namely the hardening variable never can increase while  $\xi$  and thus also the set of  $\varsigma$  satisfying  $\delta_{P_0}^*(\varsigma) \leq 1 + q_{\text{H}} \xi$  in (4.19) cannot only decrease. Also (4.1a) obviously holds just with  $P$  from (4.18), and  $C = q_{\text{H}}$  in (4.1b) with  $L = |P| = \text{Lipschitz constant of } \delta_P^*$ , and  $\eta = -\delta_P^*(\pi)$  in (4.1b).

**Remark 4.3.** The isotropic hardening  $\eta$  can be omitted if  $b = 0$  because then there is no driving force for the isotropic hardening and, if  $\eta_0 = 0$ , then identically  $\eta \equiv 0$ . Then the dissipation potential (2.6d) reduces to  $\delta_P^*(\dot{\pi})$ , cf. (3.2d) above.

## 5. NUMERICAL APPROXIMATION AND ITS CONVERGENCE

Numerical convergence results for the problem with hardening are classical, cf. [21, 49, 50, 54] for a survey. However, most of the numerical results are based on the assumption of (often not justified) higher regularity, which then allows us to derive convergence rates, cf. the above references and e.g., [3]. Actually, regularity for perfect plasticity up to the boundary is a difficult problem, requiring high

smoothness of the boundary, as documented in [7, 13, 16, 29]. Besides, the hardening parameters are considered fixed. Moreover, there are some numerical studies directly for Prandtl-Reuss plasticity, e.g. [26, 28, 45, 46]. This suggests a chance for an unconditional convergence jointly with both hardening parameters and numerical discretization parameters approaching 0. This is indeed what we will prove, cf. Theorem 5.1 below.

In the context of rate-independent processes, it is natural to discretize in time just by the fully implicit 1st-order Euler scheme. We consider the constant time-step  $\tau$ , and abbreviate  $t_\tau^k := k\tau$ .

We further consider spatial approximation, using  $C^0$ -conforming P1-elements for the approximation of  $u$  and P0-elements for the approximation of  $\pi$  and  $\eta$ . We will further make a spatial discretization. For this, we assume that we are given a sequence of triangulations  $\{\mathcal{T}_h\}_{h>0}$  of the polyhedral domain  $\Omega$  without hanging nodes but otherwise entirely general. We suppose that  $h > 0$  range over countable sets of positive real numbers with accumulation points at 0, and that  $\max_{E \in \mathcal{T}_h} \text{diam}(E) \leq h$ . The finite-dimensional subspaces of  $L^2(\Omega)$  and  $W^{1,2}(\Omega)$  related to P0- and P1-elements and subordinate to the triangulation  $\mathcal{T}_h$  respectively by  $V_{0,h}$  and  $V_{1,h}$ .

The approximate solution  $q_{\varepsilon\tau h}^k = (u_{\varepsilon\tau h}^k, \pi_{\varepsilon\tau h}^k, \eta_{\varepsilon\tau h}^k)$  is then obtained recursively for  $k = 1, \dots, T/\tau$  as the solution to the incremental problem

$$\left. \begin{array}{l} \text{Minimize} \quad \mathcal{E}_\varepsilon(t_k, u, \pi, \eta) + \mathcal{R}(\pi - \pi_{\varepsilon\tau h}^{k-1}, \eta - \eta_{\varepsilon\tau h}^{k-1}) \\ \text{subject to} \quad u \in V_{1,h}^d, \quad \pi \in V_{0,h}^{d \times d}, \quad \eta \in V_{0,h}, \end{array} \right\} \quad (5.1)$$

with  $\mathcal{E}_\varepsilon$  from (4.2) and  $\mathcal{R}$  from (4.18), when starting from  $q_{\varepsilon\tau h}^0 = q_0$ . It is notable that (5.1) has a unique solution  $q_{\varepsilon\tau h}^k$ ,  $k = 1, \dots, T/\tau$ , which satisfies the two-sided energy inequality

$$\begin{aligned} L_{\varepsilon\tau h}^k &:= \int_{t_{k-1}}^{t_k} \langle \dot{f}_{\text{ext}}(t), q_{\varepsilon\tau h}^k \rangle dt \\ &\leq \mathcal{E}_\varepsilon(t_k, q_{\varepsilon\tau h}^k) - \mathcal{E}_\varepsilon(t_{k-1}, q_{\varepsilon\tau h}^{k-1}) + \text{Diss}_{\mathcal{R}}(q_{\varepsilon\tau h}; [t_{k-1}, t_k]) \\ &\leq \int_{t_{k-1}}^{t_k} \langle \dot{f}_{\text{ext}}(t), q_{\varepsilon\tau h}^{k-1} \rangle dt =: U_{\varepsilon\tau h}^k, \end{aligned} \quad (5.2)$$

and the discrete stability

$$\forall \tilde{q} = (\tilde{u}, \tilde{\pi}, \tilde{\eta}) \in V_{1,h}^d \times V_{0,h}^{d \times d} \times V_{0,h}: \quad \mathcal{E}_\varepsilon(t_k, q_{\varepsilon\tau h}^k) \leq \mathcal{E}_\varepsilon(t_k, \tilde{q}) + \mathcal{R}(\tilde{\pi} - \pi_{\varepsilon\tau h}^{k-1}, \tilde{\eta} - \eta_{\varepsilon\tau h}^{k-1}), \quad (5.3)$$

see [40, 42].

Let us define the piecewise affine interpolant  $(u_{\varepsilon\tau h}, \pi_{\varepsilon\tau h}, \eta_{\varepsilon\tau h})$  by

$$\begin{aligned} [u_{\varepsilon\tau h}, \pi_{\varepsilon\tau h}, \eta_{\varepsilon\tau h}](t) &:= \frac{t - t_\tau^{k-1}}{\tau} (u_{\varepsilon\tau h}^k, \pi_{\varepsilon\tau h}^k, \eta_{\varepsilon\tau h}^k) + \frac{t_\tau^k - t}{\tau} (u_{\varepsilon\tau h}^{k-1}, \pi_{\varepsilon\tau h}^{k-1}, \eta_{\varepsilon\tau h}^{k-1}) \\ &\quad \text{for } t \in [t_\tau^{k-1}, t_\tau^k] \text{ with } k = 0, \dots, K_\tau := T/\tau. \end{aligned} \quad (5.4)$$

Besides, we define also the back-ward piecewise constant interpolant

$$[\bar{u}_{\varepsilon\tau h}, \bar{\pi}_{\varepsilon\tau h}, \bar{\eta}_{\varepsilon\tau h}](t) := (u_{\varepsilon\tau h}^k, \pi_{\varepsilon\tau h}^k, \eta_{\varepsilon\tau h}^k) \quad \text{for } t \in (t_\tau^{k-1}, t_\tau^k], \quad k = 1, \dots, K_\tau. \quad (5.5)$$

For fixed  $\varepsilon > 0$ , taking into account the uniqueness of the energetic solution  $q_\varepsilon = (u_\varepsilon, \pi_\varepsilon, \eta_\varepsilon)$  as stated in Proposition 2.1, one obtains the convergence of the whole sequence  $\tau \rightarrow 0$  and  $h \rightarrow 0$ , cf. [37, Sect.4.2] or also [23]. The strategy is now

to merge this convergence with the limit  $\varepsilon \rightarrow 0$  as stated in Theorem 4.1. Clearly we have the successive convergence of the type

$$\lim_{\varepsilon \rightarrow 0} \lim_{\substack{\tau \rightarrow 0 \\ h \rightarrow 0}} q_{\varepsilon\tau h} = q. \quad (5.6)$$

By arguments as in [31] or [6, Cor.4.8(ii)], one can formulate this result by means of an implicit “stability criterion”, involving functions  $E_1, E_2 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ , such that all cluster points in  $\{(u_{\varepsilon\tau h}, \pi_{\varepsilon\tau h}, \eta_{\varepsilon\tau h})\}_{\varepsilon > 0, \tau > 0, h > 0, \tau \leq E_1(\varepsilon), h \leq E_2(\varepsilon)}$  yield energetic solutions, i.e. one gets the conditional convergence of the type:

$$\lim_{\substack{\varepsilon \rightarrow 0, \tau \rightarrow 0, h \rightarrow 0 \\ \tau \leq E_1(\varepsilon), h \leq E_2(\varepsilon)}} q_{\varepsilon\tau h} = q. \quad (5.7)$$

By a careful inspection of the proof of Theorem 4.1 and a sophisticated construction of the mutual recovery sequence, we can improve (5.7) to the following unconditional convergence result.

**Theorem 5.1** (Unconditional convergence of FEM-discretisation). *Let the assumptions for Theorem 4.1 hold. Then every sequence  $\{(u_{\varepsilon\tau h}, \pi_{\varepsilon\tau h}, \eta_{\varepsilon\tau h})\}_{\varepsilon, \tau, h > 0}$  of the approximate solutions obtained by (5.1) contains a subsequence which converges for  $(\varepsilon, \tau, h) \rightarrow (0, 0, 0)$  weakly\* in the topologies indicated in (4.3a-c), i.e.*

$$\lim_{(\varepsilon, \tau, h) \rightarrow (0, 0, 0)} \bar{u}_{\varepsilon\tau h} = u \quad \text{weakly* in } L^\infty(I; \text{BD}(\Omega; \mathbb{R}^d)), \quad (5.8a)$$

$$\lim_{(\varepsilon, \tau, h) \rightarrow (0, 0, 0)} \bar{\pi}_{\varepsilon\tau h} = \pi \quad \text{weakly* in } \text{BV}(\bar{I}; \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d})), \quad (5.8b)$$

$$\lim_{(\varepsilon, \tau, h) \rightarrow (0, 0, 0)} e(u_{\varepsilon\tau h}) - \pi_{\varepsilon\tau h} = e(u) - \pi \quad \text{weakly* in } L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d})). \quad (5.8c)$$

Any limit  $(u, \pi)$  obtained by this way is an energetic solution in the sense of (1.2) to the Prandtl-Reuss model  $(Q_{\text{PR}}, \mathcal{E}_{\text{PR}}, \mathcal{R}_{\text{PR}}, u_0, \pi_0)$  as defined in (3.2). Moreover, the whole sequence (not only selected subsequences) of stresses  $\{\mathbb{C}(e(u_{\varepsilon\tau h}) - \pi_{\varepsilon\tau h})\}_{\varepsilon, \tau, h > 0}$  converges weakly\* in  $L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}))$ , and even strongly pointwise, i.e. for any  $t \in [0, T]$ , it holds

$$\lim_{(\varepsilon, \tau, h) \rightarrow (0, 0, 0)} \mathbb{C}(e(u_{\varepsilon\tau h}(t)) - \pi_{\varepsilon\tau h}(t)) = \mathbb{C}(e(u(t)) - \pi(t)|_\Omega) \quad \text{strongly in } L^2(\Omega; \mathbb{R}^{d \times d}). \quad (5.9)$$

*Proof.* We highlight only the differences to the proof of Theorem 4.1. It essentially concerns only Step 3, namely a construction and usage of a suitable mutual recovery sequence.

Let us abbreviate  $\mathcal{E}_{\varepsilon h}(t, \cdot, \cdot, \cdot) := \mathcal{E}_\varepsilon(t, \cdot, \cdot, \cdot) + \delta_{V_{1,h}^d \times V_{0,h}^{d \times d} \times V_{0,h}}(\cdot, \cdot, \cdot)$ . Again consider  $(u, \pi) \in Q_{\text{PR}}$  and a stable sequence  $(u_{\varepsilon h}, \pi_{\varepsilon h}) \xrightarrow{*} (u, \pi)$  in  $Q_{\text{PR}}$ , i.e.

$$u_{\varepsilon h} \xrightarrow{*} u \quad \text{in } \text{BD}(\Omega; \mathbb{R}^d), \quad (5.10a)$$

$$\pi_{\varepsilon h} \xrightarrow{*} \pi \quad \text{in } \mathcal{M}(\Omega \cup \Gamma_D; \mathbb{R}_{\text{dev}}^{d \times d}), \quad (5.10b)$$

$$e(u_{\varepsilon h}) - \pi_{\varepsilon h} \rightharpoonup e(u) - \pi \quad \text{in } L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}), \quad (5.10c)$$

with  $(u_{\varepsilon h}, \pi_{\varepsilon h}) \in V_{1,h}^d \times V_{0,h}^{d \times d}$  and an arbitrary  $\eta_{\varepsilon h} \in V_{0,h}$ . To facilitate also the time discretization, a floating time  $t_\tau \rightarrow t$  has to be considered instead of the fixed time considered before in (4.11). Here stability of the sequence  $\{(t_\tau, u_{\varepsilon h}, \pi_{\varepsilon h}, \eta_{\varepsilon h})\}_{\varepsilon, h > 0}$  for the collection of functionals  $\{(\mathcal{E}_{\varepsilon h}, \mathcal{R})\}_{\varepsilon, h > 0}$  means  $\sup_{\varepsilon, h > 0} \mathcal{E}_{\varepsilon h}(t_\varepsilon, u_{\varepsilon h}, \pi_{\varepsilon h}, \eta_{\varepsilon h}) < \infty$  and  $(u_{\varepsilon h}, \pi_{\varepsilon h}, \eta_{\varepsilon h}) \in \mathcal{S}_{\varepsilon h}(t_\tau)$  with  $\mathcal{S}_{\varepsilon h}$  from (2.7) but with  $\mathcal{E}$  replaced by  $\mathcal{E}_{\varepsilon h}$ . In particular,  $\|\pi_{\varepsilon h}\|_{L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})} = \mathcal{O}(\varepsilon^{-1/2})$  and  $\|\eta_{\varepsilon h}\|_{L^2(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})} = \mathcal{O}(\varepsilon^{-1/2})$ .



We consider projectors  $\Pi_{h,1}$  and  $\Pi_{h,0}$  onto the P1- and P0-finite element spaces, respectively, which are assumed to satisfy

$$\Pi_{h,0}f \rightarrow f \text{ in } L^2(\Omega) \quad \text{and} \quad \Pi_{h,1}g \rightarrow g \text{ in } H^1(\Omega) \quad (5.11)$$

for all  $f \in L^2(\Omega)$  and all  $g \in H^1(\Omega)$ . Again consider arbitrary  $\hat{u} \in H^1(\Omega; \mathbb{R}^d)$  with  $\hat{u}|_{\Gamma_D} = 0$  and  $\hat{\pi} \in L^2(\Omega; \mathbb{R}^{d \times d}_{\text{dev}})$  as in (4.8). Then it is always possible to take  $\hat{\eta}_h \in V_{0,h}$  such that  $\delta_S^*(\Pi_{h,0}\hat{\pi}, \hat{\eta}_h) = \delta_P^*(\Pi_{h,0}\hat{\pi})$ . Here we used that  $S$  and  $P$  contain 0, are convex, and do not depend on  $x$  and hence the element-wise averaging involved in  $\Pi_{h,0}$  allows for a point-wise construction on each element. Then, likewise in (4.9), put

$$\tilde{u}_{\varepsilon h} = u_{\varepsilon h} + \Pi_{h,1}\hat{u}, \quad \tilde{\pi}_{\varepsilon h} = \pi_{\varepsilon h} + \Pi_{h,0}\hat{\pi} \quad \text{and} \quad \tilde{\eta}_{\varepsilon h} = \eta_{\varepsilon h} + \hat{\eta}_h. \quad (5.12)$$

Then one has an analog of (4.10):

$$\begin{aligned} \mathcal{R}(\tilde{\pi}_{\varepsilon h} - \pi_{\varepsilon h}, \tilde{\eta}_{\varepsilon h} - \eta_{\varepsilon h}) &= \int_{\Omega} \delta_S^*(\tilde{\pi}_{\varepsilon h} - \pi_{\varepsilon h}, \tilde{\eta}_{\varepsilon h} - \eta_{\varepsilon h}) \, dx = \int_{\Omega} \delta_S^*(\Pi_{h,0}(\tilde{\pi} - \pi), \tilde{\eta}_h) \, dx \\ &= \int_{\Omega} \delta_S^*(\Pi_{h,0}\hat{\pi}, \hat{\eta}_h) \, dx = \int_{\Omega} \delta_P^*(\Pi_{h,0}\hat{\pi}) \, dx \\ &\rightarrow \int_{\Omega} \delta_P^*(\hat{\pi}) \, dx = \mathcal{R}_{\text{PR}}(\hat{\pi}), \end{aligned} \quad (5.13)$$

where the last convergence uses the  $L^2(\Omega)$  convergence assumed in (5.11).

Taking into account the quadratic structure of both  $\mathcal{E}_{\varepsilon h}$  (if restricted on the finite-dimensional FE subspaces) and (5.13), we have

$$\begin{aligned} &\mathcal{E}_{\varepsilon h}(t_{\tau}, \tilde{u}_{\varepsilon h}, \tilde{\pi}_{\varepsilon h}, \tilde{\eta}_{\varepsilon h}) - \mathcal{E}_{\varepsilon h}(t_{\tau}, u_{\varepsilon h}, \pi_{\varepsilon h}, \eta_{\varepsilon h}) \\ &= \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\tilde{u}_{\varepsilon h} - u_{\varepsilon h}) - \tilde{\pi}_{\varepsilon h} + \pi_{\varepsilon h}) : (e(\tilde{u}_{\varepsilon h} + u_{\varepsilon h}) - \tilde{\pi}_{\varepsilon h} - \pi_{\varepsilon h}) \\ &\quad + \frac{\varepsilon}{2} \mathbb{H}(\tilde{\pi}_{\varepsilon h} - \pi_{\varepsilon h}) : (\tilde{\pi}_{\varepsilon h} + \pi_{\varepsilon h}) + \frac{\varepsilon}{2} (\tilde{\eta}_{\varepsilon h} - \eta_{\varepsilon h}) (\tilde{\eta}_{\varepsilon h} + \eta_{\varepsilon h}) \, dx \\ &\quad - \langle f_{\text{ext}}(t_{\tau}), (\tilde{u}_{\varepsilon h} - u_{\varepsilon h}, \tilde{\pi}_{\varepsilon h} - \pi_{\varepsilon h}) \rangle \\ &= \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\Pi_{h,1}\hat{u}) - \Pi_{h,0}\hat{\pi}) : (e(2u_{\varepsilon h} + \Pi_{h,0}\hat{u}) - 2\pi_{\varepsilon h} - \Pi_{h,0}\hat{\pi}) \\ &\quad + \frac{\varepsilon}{2} \mathbb{H}\Pi_{h,0}\hat{\pi} : (2\pi_{\varepsilon h} + \Pi_{h,0}\hat{\pi}) + \frac{\varepsilon}{2} \hat{\eta}_h (\tilde{\eta}_{\varepsilon h} + \eta_{\varepsilon h}) \, dx - \langle f_{\text{ext}}(t_{\tau}), (\Pi_{h,1}\hat{u}, \Pi_{h,0}\hat{\pi}) \rangle. \end{aligned} \quad (5.14)$$

From this, by using the convergence (5.10), (5.13), and the quadratic structure of  $\mathcal{E}_{\text{PR}}$ , one can see that (4.11) modifies as follows:

$$\begin{aligned} &\lim_{(\varepsilon, \tau, h) \rightarrow (0,0,0)} \mathcal{E}_{\varepsilon h}(t_{\tau}, \tilde{u}_{\varepsilon h}, \tilde{\pi}_{\varepsilon h}, \tilde{\eta}_{\varepsilon h}) - \mathcal{E}_{\varepsilon h}(t_{\tau}, u_{\varepsilon h}, \pi_{\varepsilon h}, \eta_{\varepsilon h}) \\ &= \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\hat{u}) - \hat{\pi}) : (e(2u + \hat{u}) - 2\pi - \hat{\pi}) \, dx - \langle f_{\text{ext}}(t), (\hat{u}, \hat{\pi}) \rangle \\ &= \int_{\Omega} \frac{1}{2} \mathbb{C}(e(\tilde{u} - u) - \tilde{\pi} + \pi) : (e(\tilde{u} + u) - \tilde{\pi} - \pi) \, dx - \langle f_{\text{ext}}(t), (\tilde{u} - u, \tilde{\pi} - \pi) \rangle \\ &= \mathcal{E}_{\text{PR}}(t, \tilde{u}, \tilde{\pi}) - \mathcal{E}_{\text{PR}}(t, u, \pi), \end{aligned} \quad (5.15)$$

where we used (5.10c) and also the estimates analogous to (4.12). This allows for the limit passage in the stability condition for the discretised problem with hardening, i.e. in

$$\bar{\mathcal{E}}_{\varepsilon\tau h}(t, \bar{u}_{\varepsilon\tau h}(t), \bar{\pi}_{\varepsilon\tau h}(t), \bar{\eta}_{\varepsilon\tau h}(t)) \leq \bar{\mathcal{E}}_{\varepsilon\tau h}(t, \tilde{u}, \tilde{\pi}, \tilde{\eta}) + \mathcal{R}(\tilde{\pi} - \bar{\pi}_{\varepsilon\tau h}(t), \tilde{\eta} - \bar{\eta}_{\varepsilon\tau h}(t)) \quad (5.16)$$

for any  $t \in [0, T]$  and any  $(\tilde{u}, \tilde{\pi}, \tilde{\eta}) \in V_{1,h}^d \times V_{0,h}^{d \times d} \times V_{0,h}$ , where we used the notation  $\bar{\mathcal{E}}_{\varepsilon\tau h}(t, q) := \mathcal{E}_{\varepsilon h}(t_\tau^k, q)$  for  $t \in (t_\tau^{k-1}, t_\tau^k]$ , cf. (5.3) and the definition (5.5). In fact, one is to use (5.15) with the fact that, for all  $t \in [0, T]$ , one has  $\bar{\mathcal{E}}_{\varepsilon\tau h}(t, q) = \mathcal{E}_{\varepsilon h}(t_\tau, q)$  with  $t_\tau := \min\{t_\tau^k \geq t; k = 0, \dots, T/\tau\} \rightarrow t$  for  $\tau \rightarrow 0$ . Thus we can prove the weaker stability condition (4.13), apply (3.6) to obtain the full stability, and finish as in the proof of Theorem 4.1.  $\square$

## 6. NUMERICAL EXPERIMENTS

We present computational experiments on 2D and 3D Lipschitz domains, so, theoretically, only Theorem 4.1(i-ii) but not (iii) is at our disposal. The two-dimensional setting allows us to use finer discretizations and thereby to consider smaller values of the hardening parameter  $\varepsilon$ . Throughout this section we employ an isotropic material with isotropic hardening and without kinematic hardening. More specifically, we consider the ansatz (4.2) with  $\mathbb{C}e = \lambda(\text{tr } e)\mathbb{I} + 2\mu e$  and with  $\mathbb{H} = 0$ . Having in mind materials like steel, we consider the Lamé constants  $\lambda = 79$  GPa and  $\mu = 52.7$  GPa, which correspond to Young's modulus 137 GPa and Poisson's ratio 0.3. Moreover, we employ the hardening parameter  $b = \mu$  in (4.2) and study different choices of  $\varepsilon$ . We further consider the dissipation determined by  $S$  in (4.19) with  $P = \{\varsigma \in \mathbb{R}_{\text{dev}}^{d \times d} \mid |\varsigma| \leq \sigma_y\}$  with  $\sigma_y := 450$  MPa and  $q_{\mathbb{H}} = 1/\sigma_y$ . The coercivity condition (2.9) holds even if  $\mathbb{H} = 0$  because  $|\pi| \rightarrow \infty$  makes here also  $\eta \rightarrow -\infty$ ; more specifically,  $\eta \leq -q_{\mathbb{H}}\sigma_y|\pi|$  and thus the infimum in (2.9) is exactly  $bq_{\mathbb{H}}^2\sigma_y^2/(1+q_{\mathbb{H}}^2\sigma_y^2)$ . The nonlinear systems of equations arising in each time step were solved by a Newton method using ideas from [8].

**6.1. Two-dimensional experiment.** We consider a rectangular specimen with slightly generalized Dirichlet boundary conditions:

$$\Omega = (-5\frac{L}{2}, 5\frac{L}{2}) \times (-\frac{L}{2}, \frac{L}{2}), \quad (6.1a)$$

$$\Gamma_{\text{D}} = \Gamma_{\text{D}}^{\text{top}} \cup \Gamma_{\text{D}}^{\text{bottom}}, \quad \Gamma_{\text{D}}^{\text{top}} = (-5\frac{L}{2}, \frac{L}{2}) \times \{\frac{L}{2}\}, \quad (6.1b)$$

$$\Gamma_{\text{D}}^{\text{bottom}} = (-\frac{L}{2}, 5\frac{L}{2}) \times \{-\frac{L}{2}\}, \quad (6.1c)$$

$$u_{\text{D}}(t, x) = 0 \quad \text{for } x \in \Gamma_{\text{D}}^{\text{bottom}}, \text{ and} \quad (6.1d)$$

$$(u_{\text{D}})_2(t, x) = -t \cdot 2 \cdot 10^{-3} \text{ m/s} \quad \text{for } x \in \Gamma_{\text{D}}^{\text{top}}, \quad (6.1e)$$

i.e., only the normal displacement on  $\Gamma_{\text{D}}^{\text{top}}$  is prescribed (and gradually increasing in time). Allowing the top side gliding facilitates the development of a shear band better than if it would be fixed, although this requires a modification of the arguments presented in Sections 2-5; in particular, instead of  $u \otimes \nu \, d\mathcal{H}^{d-1} + \pi = 0$  in (3.2b), one should prescribe  $u \cdot \nu \, d\mathcal{H}^{d-1} + \nu \cdot \pi \cdot \nu = 0$  on this “gliding” boundary. Moreover, we set  $T = 1$  s but note that, in fact, this physical unit is only related to (6.1e), otherwise it is irrelevant in the considered rate-independent case.




FIGURE 1. Geometry for the 2D example in Sect. 6.1

The employed triangulations are so-called red-refinements of mesh-width  $h_\ell = 2^{-\ell} \cdot 10^{-2}\text{m}$  of the coarse triangulation with  $h_0 = L = 10^{-2}\text{m}$  and 10 triangles displayed in Figure 1. We emphasize that the triangulations do not match the expected  $45^\circ$ -oriented slip band in this example. We always employ the time step

$$\tau = \tau_\ell = h_\ell/v_0, \quad \text{with } v_0 = 0.04\text{m/s}. \quad (6.2)$$

Snapshots of the discrete evolution with  $\ell = 7$ , i.e., a triangulation with  $4^7 \cdot 10 = 163\,840$  triangles, and  $\varepsilon = 1/64$  for  $t = 0.25\text{s}$ ,  $0.5\text{s}$ ,  $0.75\text{s}$ , and  $1\text{s}$  are shown in Figure 2, where we plotted the hardening variable  $\eta_h(\cdot, t)$  together with the magnified discrete displacement field  $u_h(\cdot, t)$ . We observe that a plastic region develops which connects those points on the upper and lower boundary of  $\Omega$  at which the type of the boundary conditions change.




FIGURE 2. Snapshots of the time evolution of one slip-band depicted via the hardening variable  $\eta_h$  with  $\varepsilon = 1/64$  in Example 6.1;  $t = 0.25\text{s}$ ,  $0.5\text{s}$ ,  $0.75\text{s}$ , and  $1\text{s}$ . The displacement is magnified by a factor 4.

To visualize the dependency of  $\eta_h$  on  $\varepsilon$ , we displayed for  $\ell = 7$  and  $\varepsilon = 4^{-m}$ ,  $m = 0, 1, 2, 3$ , in the left and right plot of Figure 3 the functions

$$t \mapsto \|\eta_h(\cdot, t)\|_{L^2(\Omega)}, \quad x_K^{\text{cs}} \mapsto -\eta_h(x_K^{\text{cs}}, T), \quad (6.3)$$

where  $x_K^{\text{cs}}$  ranges over a neighbourhood of the cross-section indicated in Figure 1; more specifically,  $x_K^{\text{cs}}$  are those midpoints of elements whose distance from this cross-section is less than  $10^{-4}\text{m}$ .

The entire functions  $\eta_h(\cdot, T)$  at the end of the evolution are displayed for  $\varepsilon = 4^{-m}$ ,  $m = 0, 1, 2, 3$  and  $\ell = 7$  in Figure 4. We observe a diffuse plastic region for large hardening parameters  $\varepsilon b$  and, on the other hand, a narrower region with sharper interfaces (= a slip band) for the smallest value of  $\varepsilon b$ .

We experimentally studied the validity of the two-sided energy inequality discussed in Section 5. Figure 5 displays the discrete upper and lower bounds for discretizations defined through  $\ell = 4, 5, 6, 7$  uniform refinements with corresponding time-step sizes and for fixed  $\varepsilon = 1/16$ . We see in the left plot of Figure 5 that the discrete upper bound dominates the lower bound for  $\ell = 4$ . This is predicted by the theoretical estimates (5.2) which assume an exact solution of the nonlinear systems




FIGURE 3. Illustration of development of a shear band when  $\varepsilon \downarrow 0$ : the  $L^2(\Omega)$ -norm of  $\eta_h(\cdot, t)$  for time  $0 \leq t \leq T = 1$  showing the blow-up tendency  $\mathcal{O}(1/\sqrt{\varepsilon})$  predicted in (4.3f) (left) and a spatial profile of  $-\eta_h(\cdot, T)$  in Example 6.1 along the middle cross-section indicated in Fig. 1 at the terminal time  $t = T$  (right).




FIGURE 4. Displacement  $u_h$  and hardening variable  $\eta_h$  ( $\sim |\text{plastic strain}|$ ) for  $t = T$  and decreasing hardening parameter  $\varepsilon = 1, 1/4, 1/16,$  and  $1/64$ , showing how a diagonal slip-band becomes more and more pronounced. The underlying triangulation has 163 840 triangles, the discrete displacement is displayed magnified by a factor 4.

of equations. Let us remark that Figure 5(left) refers to the problem with transformed Dirichlet condition and the displayed curves depend on the specific choice of  $u_D$  and do not represent the real physical power; more specifically, our choice of  $u_D$  was just to be the (approximate) elasto-plastic solution of the original problem, which makes the “transformed” power zero until the plasticification starts. To compare the upper and lower bound for different discretization parameters we scaled them by  $1/\tau$  and in the right plot of Figure 5 we plotted their difference with a logarithmic scale on the  $y$ -axis. We see that the difference decays linearly to 0 with the spatial and temporal discretization parameters.




FIGURE 5. Rates (=power) of the upper and the lower bounds in (5.2) for  $h = 2^{-4}\text{cm}$  and  $\tau \doteq 16 \cdot 10^{-3}\text{s}$  (left) and difference between them in a semi-logarithmic scaling for different discretization parameters, indicating convergence like  $\mathcal{O}(\tau)$  (right);  $\varepsilon = 1/16$ .

**6.2. Three-dimensional experiment.** Although we know that approximations converge unconditionally to the unique solution, the qualitative behaviour of the discrete solutions may depend on the employed triangulations in a preasymptotic range. In our three-dimensional experiment the underlying triangulations match the expected geometry of the slip band.

We let

$$\Omega = (-2L, 2L) \times (-\frac{L}{2}, \frac{L}{2})^2, \quad (6.4a)$$

$$\Gamma_D = \Gamma_D^{\text{top}} \cup \Gamma_D^{\text{bottom}}, \quad \Gamma_D^{\text{top}} = (0, \frac{L}{2}) \times (-\frac{L}{2}, \frac{L}{2}) \times \{\frac{L}{2}\}, \quad (6.4b)$$

$$\Gamma_D^{\text{bottom}} = (-2L, 0) \times (-\frac{L}{2}, \frac{L}{2}) \times \{-\frac{L}{2}\}, \quad (6.4c)$$

$$u_D(t, x) = 0 \quad \text{for } x \in \Gamma_D^{\text{bottom}}, \text{ and} \quad (6.4d)$$

$$(u_D)_3(t, x) = -t \cdot 2 \cdot 10^{-3} \text{m/s} \quad \text{for } x \in \Gamma_D^{\text{top}}, \quad (6.4e)$$

for  $L = 1 \cdot 10^{-2}\text{m}$ , i.e., again only the normal displacement is prescribed on the top part of  $\Gamma_D$ , and  $\Gamma_N = \partial\Omega \setminus \Gamma_D$




FIGURE 6. Geometry for the 3D example in Sect. 6.2

The employed triangulations are red-refinements of mesh-width  $h_\ell = 2^{-\ell} \cdot 10^{-2}\text{m}$  with  $8^\ell \cdot 24$  tetrahedra of the coarse triangulation with  $h_0 = 10^{-2}\text{m}$  displayed in Figure 6. We always used the time-step size  $\tau = 0.025 \cdot 2^{-\ell}\text{s}$ .

For  $\ell = 4$  and  $\varepsilon = 4^{-m}$ ,  $m = 0, 1, 2$ , we displayed in the left and right plot of Figure 7 the functions

$$t \mapsto \|\eta_h(\cdot, t)\|_{L^2(\Omega)}, \quad x_K^{\text{ax}} \mapsto -\eta_h(x_K^{\text{ax}}, T),$$

where, like in (6.3),  $x_K^{\text{ax}}$  ranges over a neighbourhood of the horizontal axis in Figure 6; more specifically,  $x_K^{\text{ax}}$  are those midpoints of elements whose distance from this axis is less than  $10^{-4}\text{m}$ .




FIGURE 7. The  $L^2(\Omega)$ -norm of  $\eta_h(\cdot, t)$  for time  $0 \leq t \leq T = 1$  (left) and a spatial profile of  $\eta_h(T, \cdot)$  along the horizontal axis of  $\Omega$  in Example 6.2

The functions  $\eta_h(T, \cdot)$  are displayed for  $\ell = 7$  and  $\varepsilon = 4^{-m}$ ,  $m = 0, 1, 2, 3$  in Figure 8. We observe that for  $\varepsilon$  tending to zero, the developed plastic region becomes narrower and the interfaces less diffuse.




FIGURE 8. Displacement  $u_h$  and hardening variable  $\eta_h$  ( $\sim$  |plastic strain|) for  $t = T$  and  $\varepsilon = 1, 1/4, \text{ and } 1/16$ , showing how a vertical slip-band becomes more and more pronounced. The underlying triangulation has 98 304 tetrahedra, the displacement is magnified by a factor 4.

**Remark 6.1** (Regularity, rate of convergence). If  $\Omega$  is smooth and the loading has additional regularity in time, it has been shown in [29] that the energetic solution  $(u_\varepsilon, z_\varepsilon)$  admits additional regularity properties, namely

$$u_\varepsilon \in L^\infty(I; W^{3/2-\alpha}(\Omega; \mathbb{R}^d)), \quad z \in L^\infty(I; W^{1/2-\alpha}(\Omega; \mathbb{R}^{d \times d} \times \mathbb{R})), \quad \text{for some } \alpha > 0. \quad (6.5)$$

This can further be used, cf. [30], to derive the rate of convergence:

$$\|u_\varepsilon - \bar{u}_{\varepsilon\tau h}\|_{L^\infty(I; H^1(\Omega; \mathbb{R}^d))} + \|z_\varepsilon - \bar{z}_{\varepsilon\tau h}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d} \times \mathbb{R}))} \leq C(\alpha, \varepsilon)(\sqrt{\tau} + {}^4\sqrt{h}). \quad (6.6)$$

The constant  $C(\alpha, \varepsilon)$  naturally depends on the positive-definiteness constant  $\varepsilon$  of  $\mathcal{E}^\varepsilon(t, \cdot, \cdot)$  and one can identify from [30] the explicit upper bound in the form  $C(\alpha, \varepsilon) =$

$C_\alpha \varepsilon^{-2} e^{c/\varepsilon}$  with some  $c, C \in \mathbb{R}$ . Obviously,  $C(\alpha, \varepsilon)$  grows essentially like  $e^{c/\varepsilon}$ . In smooth cases, one thus gets an explicit hint how to converge simultaneously with hardening and discretisation, namely

$$\tau = o(e^{-2c/\varepsilon}), \quad h = o(e^{-(4-\alpha)c/\varepsilon}). \quad (6.7)$$

This extreme upper bound cannot be used practically, but it has the advantage that it is mathematically rigorous. Nevertheless we hope that it is too pessimistic and that it will be possible to improve the bounds considerably, at least in particular cases.

*Acknowledgment:* The authors are thankful to Dorothee Knees for making  $C(\alpha, \varepsilon)$  in (6.6) more explicit and to Filip Rindler for discussion concerning approximation issues and for the intuition about Conjecture 3.1. Moreover, the authors acknowledge partial support from Deutsche Forschungsgemeinschaft via SFB 611 ‘‘Singular phenomena and scaling in mathematical models’’ and MATHEON project C18, from the grants A 100750802 (GA AV R), 201/09/0917 and 201/10/0357 (GA R), LC 06052 and MSM 21620839 (MŠMT R), and from the research plan AV0Z20760514 ‘‘Complex dynamical systems in thermodynamics, mechanics of fluids and solids’’ (R). A.M. thanks for the hospitality of the Neas Center for Mathematical Modeling and similarly T.R. thanks for the hospitality of Weierstrass Institut fur angewandte Analysis und Stochastik in Berlin and Rheinische Friedrich-Wilhelms-Universitat Bonn, where parts of this research were performed.

#### REFERENCES

- [1] H.-D. ALBER: *Materials with Memory*, L.N. in Math. **1682**, Springer-Verlag, Berlin, 1998.
- [2] H.-D. ALBER: Mathematical analysis of constitutive equations: existence and collapse of solutions. In: *Solid Materials at Large Strains* (C.Miehe, ed.), Kluwer 2003, pp.45–54.
- [3] J. ALBERTY, C. CARSTENSEN: Numerical analysis of time-depending primal elastoplasticity with hardening. *SIAM J. Numer. Anal.* **37** (2000), 1271–1294.
- [4] L. AMBROSIO, N. FUSCO, D. PALLARA: *Functions of Bounded Variations and Free-Discontinuity Problems*. Oxford Mathematical Monographs, Oxford University Press, 2000.
- [5] G. ANZELLOTTI, S. LUCKHAUS: Dynamical evolution of elasto-perfectly plastic bodies. *Appl. Math. Optim.* **15** (1987), 121–140.
- [6] S. BARTELS, T. ROUBÍEK: Thermo-visco-elasticity with rate-independent plasticity in isotropic materials undergoing thermal expansion. (Preprint no.486, SFB 611, INS, Univ. Bonn, 2009) M2AN, accepted.
- [7] M. BULÍEK, J. FREHSE, J. MALEK: On boundary regularity for the stress in problems of linearized elasto-plasticity. *Intl. J. Advances in Engr. Sci. Appl. Math.* **1** (2010), 141–156.
- [8] C. CARSTENSEN, R. KLOSE: Elastoviscoplastic finite element analysis in 100 lines of Matlab. *J. Numer. Math.* **10** (2002), 157–192.
- [9] K. CHELMIŃSKI: Coercive approximation of viscoplasticity and plasticity. *Asymptot. Anal.*, **26** (2001), 105–133.
- [10] K. CHELMIŃSKI: Perfect plasticity as a zero relaxation limit of plasticity with isotropic hardening. *Math. Methods Appl. Sci.* **24** (2001), 117–136.
- [11] G. DAL MASO, A. DESIMONE, M.G. MORA: Quasistatic evolution problems for linearly elastic-perfectly plastic materials. *Arch. Ration. Mech. Anal.* **180** 2006, 237–291.
- [12] G. DAL MASO, R. TOADER: Quasistatic crack growth in elasto-plastic materials: the two-dimensional case. *Archive Rat. Mech. Anal.* **196** (2010), 867–906.
- [13] A. DEMYANOV: Regularity of stresses in Prandtl-Reuss perfect plasticity. *Calc. Var. P.D.E.* **34** (2009), 23–72.
- [14] F. EBOBISSE, B.D. REDDY: Some mathematical problems in perfect plasticity, *Computer Meth. Appl. Mech. Engr.* **193** (2004), 5071–5094.
- [15] G. FRANCFORT, A. MIELKE: An existence result for a rate-independent material model in the case of nonconvex energies. *J. reine u. angew. Math.* **595** (2006), 55–91.
- [16] J. FREHSE, J. MALEK: Boundary regularity results for models of elasto-perfect plasticity. *Math. Models Meth. Appl. Sci.* **9** (1999), 1307–1321

- [17] A. GIACOMINI, L. LUSSARDI: Quasistatic evolution for a model in strain gradient plasticity. *SIAM J. Math. Anal.* **40** (2008), 1201–1245.
- [18] P. GRUBER, D. KNEES, S. NESENEKO, M. THOMAS: Analytical and numerical aspects of time-dependent problems with internal variables. *Zeitsch. Angew. Math. Mech.* **90** (2010), 861–902.
- [19] P. GRUBER, J. VALDMAN: Solution of one-time-step problems in elastoplasticity by a slant Newton method. *SIAM J. Sci. Comput.* **31** (2008/09), 1558–1580.
- [20] K. GRÖGER: Zur Theorie des quasi-statischen Verhaltens von elastisch-plastischen Körpern. *Zeitsch. Angew. Math. Mech.* **58** (1978), 81–88.
- [21] W. HAN, B.D. REDDY: *Plasticity (Mathematical Theory and Numerical Analysis)*. Springer-Verlag, New York, 1999.
- [22] W. HAN, B.D. REDDY: Convergence analysis of discrete approximations of problems in hardening plasticity. *Comput. Methods Appl. Mech. Engrg.* **171** (1999), 327–340.
- [23] W. HAN, B.D. REDDY: Convergence of approximations to the primal problem in plasticity under conditions of minimal regularity. *Numer. Math.* **87** (2000), 283–315.
- [24] C. JOHNSON: On plasticity with hardening, *J. Math. Anal. Appl.* **62**, 1978, 325–336.
- [25] C. JOHNSON: Existence theorems for plasticity problems, *J. Math. Pures Appl.* **55** (1976), 431–444.
- [26] C. JOHNSON: On finite-element methods for plasticity problems. *Numer. Math.* **26** (1976), 79–84.
- [27] C. JOHNSON: A mixed finite-element method for plasticity problems with hardening. *SIAM J. Numer. Math.* **14** (1977), 575–583.
- [28] C. JOHNSON, R. SCOTT: A finite element method for problems in perfect plasticity using discontinuous trial functions. In: *Nonlin. Finite Element Anal. in Structural Mech.* (W.Wunderlich, E.Stein, K.-J.Bathe, eds.) Springer, Berlin, 1981, pp.307-324.
- [29] D. KNEES: On global spatial regularity in elasto-plasticity with linear hardening. *Calc. Var.* **36** (2009), 611–625.
- [30] D. KNEES: On global spatial regularity and convergence rates for time dependent elasto-plasticity. *Math. Models Methods Appl. Sci.* **20** (2010), 1823–1858.
- [31] M. KRUŽÍK, A. MIELKE, T. ROUBÍČEK: Modelling of microstructure and its evolution in shape-memory-alloy single-crystals, in particular in CuAlNi. *Meccanica* **40** (2005), 389–418.
- [32] J. LEMAITRE, J.L. CHABOCHE: *Mechanics of solid materials*. Cambridge, 1990 (Translated from French original *Mécanique des matériaux solides*, Dunod, 1985).
- [33] A. MAINIK, A. MIELKE: Existence results for energetic models for rate-independent systems. *Cal. Var. PDE* **22** (2005), 73–99.
- [34] G.A. MAUGIN: *The Thermomechanics of Plasticity and Fracture*. Cambridge Univ. Press, Cambridge, 1992.
- [35] A. MIELKE: Evolution of rate-independent systems. In: *Handbook of Differential Equations: Evolut. Diff. Eqs.* (Eds. C.Dafermos, E.Feireisl), pp. 461–559, Elsevier, Amsterdam (2005).
- [36] A. MIELKE, A. PETROV, J.A.C. MARTINS: Convergence of solutions of kinetic variational inequalities in the rate-independent quasistatic limit. *J. Math. Anal. Appl.* **348** (2008), 1012–1020.
- [37] A. MIELKE, T. ROUBÍČEK: Numerical approaches to rate-independent processes and applications in inelasticity. *Math. Modelling Numer. Anal.* **43** (2009), 399–428.
- [38] A. MIELKE, T. ROUBÍČEK, U. STEFANELLI:  $\Gamma$ -limits and relaxations for rate-independent evolutionary problems. *Calc. Var. PDE* **31** (2008), 387–416.
- [39] A. MIELKE, T. ROUBÍČEK, M. THOMAS: From damage to delamination in nonlinearly elastic materials at small strains. (Preprint No.1542, WIAS, Berlin, 2010.) *J. Elasticity*, submitted.
- [40] A. MIELKE, F. THEIL: A mathematical model for rate-independent phase transformations with hysteresis. In: *Models of Continuum Mechanics in Analysis and Engineering*. (Eds.: H.-D.Alber, R.Balean, R.Farwig), Shaker Ver., Aachen, pp.117-129 (1999).
- [41] A. MIELKE, F. THEIL: On rate-independent hysteresis models. *Nonlin. Diff. Eq. Appl.* **11** (2004), 151–189.
- [42] A. MIELKE, F. THEIL, V.I. LEVITAS: A variational formulation of rate-independent phase transformations using an extremum principle. *Archive Rat. Mech. Anal.* **162** (2002), 137–177.
- [43] D. ORNSTEIN: A non-inequality for differential operators in the  $L^1$ -norm. *Arch. Rat. Mech. Anal.* **11** (1962), 40–49.
- [44] W. PRAGER: A new method of analyzing stress and strains in work hardening solids. *ASME J. Appl. Mech.* **23** (1956), 493–496.



- [45] R. RANNACHER, F.T. SUTTMEIER: A posteriori error estimation and mesh adaptation for finite element models in elasto-plasticity. *Computer Methods Appl. Mech.* **176** (1999), 333-361.
- [46] S.I. REPIN: Errors of finite element method for perfectly elasto-plastic problems. *Math. Models Methods Appl. Sci.* **6** (1996), 587-607.
- [47] S.I. REPIN, J. VALDMAN: Functional a posteriori error estimates for incremental models in elasto-plasticity. *Central Euro. J. Math.* **7** (2009), 506-519.
- [48] F. RINDLER: In preparation.
- [49] J.C. SIMO: Numerical analysis and simulation of plasticity. In: *Handbook of Numerical Analysis* **6** (Eds. Ciarlet, P.G., Lions, J.L.), North-Holland, Amsterdam, 1998, pp.183-499.
- [50] J.C. SIMO, J.R. HUGHES: *Computational Inelasticity*. Springer, Berlin, 1998.
- [51] U. STEFANELLI: A variational principle for hardening elastoplasticity. *SIAM J. Math. Anal.* **40** (2008), 623-652.
- [52] P.-M. SUQUET: Existence et régularité des solutions des équations de la plasticité parfaite. *C. R. Acad. Sci. Paris Sér.A* **286** (1978), 1201-1204.
- [53] P.-M. SUQUET: Sur les équations de la plasticité: existence et régularité des solutions. *J. Mécanique* **20** (1981), 3-39.
- [54] R. TEMAM: *Mathematical Problems in Plasticity*. Gauthier-Villars, Paris, 1985.
- [55] H. ZIEGLER: A modification of Prager's hardening rule. *Quart. Appl. Math.* **17** (1959), 55-65.

INSTITUT FÜR NUMERISCHE SIMULATION, RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT BONN, WEGELERSTRASSE 6, D-53115 BONN, GERMANY

*E-mail address:* bartels@ins.uni-bonn.de

WEIERSTRASS-INSTITUT FÜR ANGEWANDTE ANALYSIS UND STOCHASTIK, MOHRENSTR.39, D-10117 BERLIN, AND

INSTITUT FÜR MATHEMATIK, HUMBOLDT UNIVERSITÄT ZU BERLIN, RUDOWER CHAUSSEE 25, D-12489 BERLIN, GERMANY.

*E-mail address:* mielke@wias-berlin.de

MATHEMATICAL INSTITUTE, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, CZ-186 75 PRAHA 8, AND

INSTITUTE OF THERMOMECHANICS OF THE ASCR, DOLEJŠKOVA 5, CZ-182 00 PRAHA 8, CZECH REPUBLIC.

*E-mail address:* roubicek@karlin.mff.cuni.cz

Bestellungen nimmt entgegen:

Sonderforschungsbereich 611  
der Universität Bonn  
Endenicher Allee 60  
D - 53115 Bonn

Telefon: 0228/73 4882

Telefax: 0228/73 7864

E-Mail: astrid.link@ins.uni-bonn.de

<http://www.sfb611.iam.uni-bonn.de/>

### Verzeichnis der erschienenen Preprints ab No. 460

460. Brenier, Yann; Otto, Felix; Seis, Christian: Upper Bounds on Coarsening Rates in Demixing Binary Viscous Liquids
461. Bianchi, Alessandra; Bovier, Anton; Ioffe, Dmitry: Pointwise Estimates and Exponential Laws in Metastable Systems Via Coupling Methods
462. Basile, Giada; Bovier, Anton: Convergence of a Kinetic Equation to a Fractional Diffusion Equation; erscheint in: Review Markov Processes and Related Fields
463. Bartels, Sören; Roubíček, Tomáš: Thermo-Visco-Elasticity with Rate-Independent Plasticity in Isotropic Materials Undergoing Thermal Expansion
464. Albeverio, Sergio; Torbin, Grygoriy: The Ostrogradsky-Pierce Expansion: Probability Theory, Dynamical Systems and Fractal Geometry Points of View
465. Capella Kort, Antonio; Otto, Felix: A Quantitative Rigidity Result for the Cubic to Tetragonal Phase Transition in the Geometrically Linear Theory with Interfacial Energy
466. Philipowski, Robert: Stochastic Particle Approximations for the Ricci Flow on Surfaces and the Yamabe Flow
467. Kuwada, Kazumasa; Philipowski, Robert: Non-explosion of Diffusion Processes on Manifolds with Time-dependent Metric; erscheint in: Mathematische Zeitschrift
468. Bacher, Kathrin; Sturm, Karl-Theodor: Ricci Bounds for Euclidean and Spherical Cones
469. Bacher, Kathrin; Sturm, Karl-Theodor: Localization and Tensorization Properties of the Curvature-Dimension Condition for Metric Measure Spaces
470. Le Peutrec, Dorian: Small Eigenvalues of the Witten Laplacian Acting on  $p$ -Forms on a Surface
471. Wirth, Benedikt; Bar, Leah; Rumpf, Martin; Sapiro, Guillermo: A Continuum Mechanical Approach to Geodesics in Shape Space
472. Berkels, Benjamin; Linkmann, Gina; Rumpf, Martin: An SL (2) Invariant Shape Median
473. Bartels, Sören; Schreier, Patrick: Local Coarsening of Triangulations Created by Bisections

474. Bartels, Sören: A Lower Bound for the Spectrum of the Linearized Allen-Cahn Operator Near a Singularity
475. Frehse, Jens; Löbach, Dominique: Improved  $L_p$ -Estimates for the Strain Velocities in Hardening Problems
476. Kurzke, Matthias; Melcher, Christof; Moser, Roger: Vortex Motion for the Landau-Lifshitz-Gilbert Equation with Spin Transfer Torque
477. Arguin, Louis-Pierre; Bovier, Anton; Kistler, Nicola: The Genealogy of Extremal Particles of Branching Brownian Motion
478. Bovier, Anton; Gayraud, Véronique: Convergence of Clock Processes in Random Environments and Ageing in the  $p$ -Spin SK Model
479. Bartels, Sören; Müller, Rüdiger: Error Control for the Approximation of Allen-Cahn and Cahn-Hilliard Equations with a Logarithmic Potential
480. Albeverio, Sergio; Kusuoka, Seiichiro: Diffusion Processes in Thin Tubes and their Limits on Graphs
481. Arguin, Louis-Pierre; Bovier, Anton; Kistler, Nicola: Poissonian Statistics in the Extremal Process of Branching Brownian Motion
482. Albeverio, Sergio; Pratsiovyta, Iryna; Torbin, Grygoriy: On the Probabilistic, Metric and Dimensional Theories of the Second Ostrogradsky Expansion
483. Bulíček, Miroslav; Frehse, Jens:  $C^{\alpha}$ -Regularity for a Class of Non-Diagonal Elliptic Systems with  $p$ -Growth
484. Ferrari, Partik L.: From Interacting Particle Systems to Random Matrices
485. Ferrari, Partik L.; Frings, René: On the Partial Connection Between Random Matrices and Interacting Particle Systems
486. Scardia, Lucia; Zeppieri, Caterina Ida: Line-Tension Model as the  $\Gamma$ -Limit of a Nonlinear Dislocation Energy
487. Bolthausen, Erwin; Kistler, Nicola: A Quenched Large Deviation Principle and a Parisi Formula for a Perceptron Version of the Grem
488. Griebel, Michael; Harbrecht, Helmut: Approximation of Two-Variate Functions: Singular Value Decomposition Versus Regular Sparse Grids
489. Bartels, Sören; Kruzik, Martin: An Efficient Approach of the Numerical Solution of Rate-independent Problems with Nonconvex Energies
490. Bartels, Sören; Mielke, Alexander; Roubicek, Tomas: Quasistatic Small-strain Plasticity in the Limit of Vanishing Hardening and its Numerical Approximation