

Literaturverzeichnis

- Arber, W. und Linn, S. (1969) DNA modification and restriction. *Annu. Rev. Biochem.* 38, 467-500.
- Arnold, F.H. (1991) Metal-affinity separation: A new dimension in protein processing. *Bio/Technology* 9, 151-156.
- Bain, J.D., Switzer, C., Chamberlin, A.R. und Benner, S.A. (1992) Ribosome-mediated incorporation of a non-standard amino acid into a peptide through expansion of the genetic code. *Nature* 356, 537-539.
- Bayer, E.A., Ben Hur, H., Gitlin, G. und Wilchek, M. (1986) An improved method for the single-step purification of streptavidin. *J. Biochem. Biophys. Methods* 13, 103.
- Bayer, E.A. und Wilchek, M. (1990) Isolation and properties of streptavidin. *Methods Enzymol.* 184, 80-89.
- Berger, S.L. (1987) Isolation of cytoplasmic RNA: ribonucleoside-vanadyl complexes. *Methods Enzymol.* 152, 227-234.
- Bieberich, E., Kapitonov, D., Tencomnao, T. und Yu, R.K. (2000) Protein-ribosome-mRNA display: affinity isolation of enzyme-ribosome-mRNA complexes and cDNA cloning in a single-tube reaction. *Anal. Biochem.* 287, 294-298.
- Billich, S., Wissel, T., Kratzin, H., Hahn, U., Hagenhoff, B., Lezius, A.G. und Spender, F. (1988) Cloning of a full-length complementary DNA for fatty-acid-binding protein from bovine heart. *Eur. J. Biochem.* 175, 549-556.
- Birnboim, H.C. und Doly, J. (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids Res.* 7, 1513-1523.
- Boder, E.T. und Wittrup, K.D. (1997) Yeast surface display for screening combinatorial polypeptide libraries. *Nat. Biotechnol.* 15, 553-557.
- Bouvet, P. und Belasco, J.G. (1992) Control of RNase E mediated RNA degradation by 5'-terminal base pairing in *E. coli*. *Nature* 360, 488.
- Burks, E.A., Chen, G., Georgiou, G. und Iverson, B.L. (1997) *In vitro* scanning saturation mutagenesis of an antibody binding pocket. *Proc. Natl. Acad. Sci. USA* 94, 412-417.
- Cadwell, R.C. und Joyce, G.F. (1992) Randomization of genes by PCR mutagenesis. *PCR Methods Appl.* 2, 28-33.
- Cadwell, R.C. und Joyce, G.F. (1994) Mutagenic PCR. *PCR Methods Appl.* 3, 136-140.
- Chapeville, F., Lipmann, F., von Ehrenstein, G., Weisblum, B., Ray, W.J. und Benzer, S. (1962) On the role of soluble ribonucleic acid encoding for amino acid. *Proc. Natl. Acad. Sci. USA* 48, 1086-1092.
- Chien, C.T., Bartel, P.L., Sternglanz, R. und Fields, S. (1991) The two-hybrid system: a method to identify and clone genes for proteins that interact with a protein of interest. *Proc. Natl. Acad. Sci. USA* 88, 9578-9582.
- Cohen, S.N., Chang, A.C.Y. und Hsu, L. (1972) Nonchromosomal antibiotic resistance in bacteria: Genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. USA* 70, 1293-1297.
- Daugherty, P.S., Olsen, M.J., Iverson, B.L. und Georgiou, G. (1999) Development of an optimized expression system for the screening of antibody libraries displayed on the *E. coli* surface. *Protein Eng.* 12, 613-621.
- De Pasquale, C. und Kanduc, D. (1998) Modulation of HPV16E7 translation by tRNAs in eukaryotic cell-free translation systems. *Biochem. Mol. Biol. Int.* 45, 1005-1009.
- Devlin, J.J., Panganiban, L.C. und Devlin, P.E. (1990) Random peptide libraries: a source of specific protein binding molecules. *Science* 249, 404-406.
- De Vries, J.K. und Zubay, G. (1967) DNA-directed peptide synthesis. II. The synthesis of the alpha-fragment of the enzyme beta-galactosidase. *Proc. Natl. Acad. Sci. USA* 57, 1010-1012.
- Doi, N. und Yanagawa, H. (1999) STABLE: protein-DNA fusion system for screening of combinatorial protein libraries *in vitro*. *FEBS Lett.* 457, 227-230.
- Eckert, K.A. und Kunkel, T.A. (1990) High fidelity DNA synthesis by the *Thermus aquaticus* DNA polymerase. *Nucleic Acids Res.* 18, 3739-3744.
- Eckert, K.A. und Kunkel, T.A. (1991) DNA polymerase fidelity and the polymerase chain reaction. *PCR Methods Appl.* 1, 17-24.
- Ellington, A.D. und Szostak, J.W. (1990) *In vitro* selection of RNA molecules that bind specific ligands. *Nature* 346, 818-822.
- Erdmann, V.A., Stiege, W., Henze, P.P. und Ulbrich, N. (1989) in: *Tharmacological interventions on central cholinergic mechanisms in senile dementia*. W. Zuckschwerdt Verlag, München, 45-54.
- Erdmann, V.A., Fuchs, U. und Stiege, W. (1994) The Protein-Bioreactor: Its Potentials for the synthesis of proteins in biotechnology, medicine and molecular biology. *Biotech 2000*, Seoul, Korea, 57-70.
- Fields, S. und Song, O. (1989) A novel genetic system to detect protein-protein interactions. *Nature* 340, 245-246.
- Freitag, S., Trong, I.L., Klumb, L.A., Chu, V., Chilkoti, A., Stayton, P.S. und Stenkamp, R.E. (1999) X-ray crystallographic studies of streptavidin mutants binding to biotin. *Biomol. Eng.* 16, 13-19.
- Frolova, L., Le Goff, X., Rasmussen, H.H., Cheperegin, S., Dugeon, G., Kress, M., Arman, I., Haenni, A.L., Celis, J.E., Philippe, M., *et al.* (1994) A highly conserved eukaryotic protein family possessing properties of polypeptide chain release factor. *Nature* 372, 701-703.

- Gates, C.M., Stemmer, W.P.C., Kaptein, R. und Schatz, P.J. (1996) Affinity selective isolation of ligands from peptide libraries through display on a *lac* repressor „headpiece dimer“. *J. Mol. Biol.* 255, 373-386.
- Georgiou, G., Poetschke, H.L., Stathopoulos, C. und Francisco, J.A. (1993) Practical applications of engineering gram-negative bacterial cell surfaces. *Trends Biotechnol.* 11, 6-10.
- Gersuk, G.M., Corey, M.J., Corey, E., Stray, J.E., Kawasaki, G.H. und Vessella, R.L. (1997) High-affinity peptide ligands to prostate specific antigen identified by polysome selection. *Biochem. Biophys. Res. Commun.* 232, 578-582.
- Gold, L.M. und Schweiger, M. (1969) Synthesis of phage-specific α - and β -glucosyl transferase directed by T-even DNA *in vitro*. *Proc. Natl. Acad. Sci. USA* 62, 892-898.
- Gold, L., Polisky, B., Uhlenbeck, O. und Yarus, M. (1995) Diversity of oligonucleotide functions. *Ann. Rev. Biochem.* 64, 763-797.
- Green, N.M. (1990) Avidin and Streptavidin. *Methods Enzymol.* 184, 51-67.
- Grentzmann, G., Brechemier-Baey, D., Heurgue-Hamard, V. und Buckingham, R.H. (1995) Function of polypeptide chain release factor RF-3 in *E. coli*. RF-3 action in termination is predominantly at UGA-containing stop signals. *J. Biol. Chem.* 270, 10595-10600.
- Hajnsdorf, E., Braun, F., Haugel-Nielsen, J., Le Derout, J. und Regnier, P. (1996) Multiple degradation pathways of the rpsO mRNA of *E. coli*. RNase E interacts with the 5' and 3' extremities of the primary transcript. *Biochimie*. 78, 416-424.
- Hanes, J. und Plückthun, A. (1997) *In vitro* selection and evolution of functional proteins by using ribosome display. *Proc. Natl. Acad. Sci. USA* 94, 4937-4942.
- Hanes, J., Jermutus, L., Weber-Bornhauser, S., Bosshard, H.R. und Plückthun, A. (1998) Ribosome display efficiently selects and evolves high-affinity antibodies *in vitro* from immune libraries. *Proc. Natl. Acad. Sci. USA* 95, 14130-14135.
- Hanes, J., Schaffitzel, C., Knappik, A. und Plückthun, A. (2000) Picomolar affinity antibodies from a fully synthetic naive library selected and evolved by ribosome display. *Nat. Biotechnol.* 18, 1287-1292.
- He, M. und Taussig, M.J. (1997) Antibody-ribosome-mRNA (ARM) complexes as efficient selection particles for *in vitro* display and evolution of antibody combining sites. *Nucleic Acids Res.* 25, 5132-5134.
- He, M., Menges, M., Groves, M.A.T., Corps, E., Liu, H., Brüggemann, M. und Taussig, M.J. (1999) Selection of a human anti-progesterone antibody fragment from a transgenic mouse library by ARM ribosome display. *J. Immunol. Methods* 231, 105-117.
- Henkel, T. und Baeuerle, P.A. (1993) Functional analysis of mutated cDNA clones by directed use of PCR products in *in vitro* transcription/translation reactions. *Anal. Biochem.* 214, 351-352.
- Hermann, T. und Patel, D.J. (2000) Adaptive recognition by nucleic acid aptamers. *Science* 287, 820-825.
- Herrlich, P. und Schweiger, M. (1974) DNA- and RNA-directed synthesis *in vitro* of phage enzymes. *Methods Enzymol.* 30, 654-669.
- Hochuli, E., Bannwarth, W., Döbeli, H., Gentz, R. und Stüber, D. (1988) Genetic approach to facilitate purification of recombinant proteins with a novel metal chelate adsorbent. *Bio/Technology*. 6, 1321-1325.
- Huppa, J.B. und Ploegh, H.L. (1997) *In vitro* translation and assembly of a complete T cell receptor-CD3 complex. *J. Exp. Med.* 186, 393-403.
- Irving, R.A., Coia, G., Roberts, A., Nuttall, S.D. und Hudson, P.J. (2001) Ribosome display and affinity maturation: from antibodies to single V-domains and steps towards cancer therapeutics. *J. Immunol. Methods* 248, 31-45.
- Ish-Horowicz, D. und Burke, J.F. (1981) Rapid and efficient cosmid cloning. *Nucl. Acids Res.* 9, 2989-2998.
- Jackson, S.E. und Fersht, A.R. (1994) Contribution of residues in the reactive site loop of chymotrypsin inhibitor 2 to protein stability and activity. *Biochemistry* 33, 13880-13887.
- Janosi, L., Shimizu, I. und Kaji, A. (1994) Ribosome recycling factor (ribosome releasing factor) is essential for bacterial growth. *Proc. Natl. Acad. Sci. USA* 91, 4249-4253.
- Jäschke, A. 1998. Catalysis of organic reactions by RNA – strategies for the selection of catalytic RNAs, in: *The many faces of RNA*, pp. 179-190. San Diego, Academic Press.
- Jermutus, L., Ryabova, L.A. und Plückthun, A. (1998) Recent advances in producing and selecting functional proteins by using cell-free translation. *Curr. Opin. Biotechnol.* 9, 534-548.
- Jermutus, L., Honegger, A., Schwesinger, F., Hanes, J. und Plückthun, A. (2000) Tailoring *in vitro* evolution for protein affinity or stability. *Proc. Natl. Acad. Sci. USA* 98, 75-80.
- Kain, K.C., Orlandi, P.A. und Lanar, D.E. (1991) Universal promoter for gene expression without cloning: expression-PCR. *Biotechniques* 10, 366-374.
- Kay, B.K., Adey, N.B., He, Y.S., Manfredi, J.P., Mataragnon, A.H. und Fowlkes, D.M. (1993) An M13 phage library displaying random 38-amino-acid peptides as a source of novel sequences with affinity to selected targets. *Gene* 128, 59-65.
- Keefe, A.D. und Szostak, J.W. (2001) Functional proteins from a random-sequence library. *Nature* 410, 715-718.

- Keefe, A.D., Wilson, D.S., Seelig, B. und Szostak, J.W. (2001a) One-step purification of recombinant proteins using a nanomolar-affinity streptavidin-binding peptide, the SBP-tag. *Protein Expression and Purification* 23, 440-446.
- Keiler, K.C., Waller, P.R.H. und Sauer, R.T. (1996) Role of a peptide tagging system in degradation of proteins synthesized from damaged messenger RNA. *Science* 271, 990-993.
- Keohavong, P. und Thilly, W.G. (1989) Fidelity of DNA polymerase in DNA amplification. *Proc. Natl. Acad. Sci. USA* 86, 9253-9257.
- Kieke, M.C., Cho, B.K., Boder, E.T., Kranz, D.M. und Wittrup, K.D. (1997) Isolation of anti-T cell receptor scFv mutants by yeast surface display. *Protein Eng.* 10, 1303-1310.
- Kigawa, T., Yabuki, T., Yoshida, Y., Tsutsui, M., Ito, Y., Shibata, T. und Yokoyama, S. (1999) Cell-free production and stable-isotope labeling of milligram quantities of protein. *FEBS Lett.* 442, 15-19.
- Kim, D.M., Kigawa, T., Choi, C.Y. und Yokoyama, S. (1996) A highly efficient cell-free protein synthesis system from *E. coli*. *Eur. J. Biochem.* 239, 881-886.
- Kim, D.M. und Choi, C.Y. (1996a) A semicontinuous prokaryotic coupled transcription/ translation system using a dialysis membrane. *Biotechnol. Prog.* 12, 645-649.
- Kim, D.M. und Swartz, J.R. (2000) Prolonging cell-free protein synthesis by selective reagent additions. *Biotechnol. Prog.* 16, 385-390.
- Klußmann, S., Nolte, A., Bald, R., Erdmann, V.A. und Fürste, J.P. (1996) Mirror-image RNA that binds D-adenosine. *Nat. Biotechnol.* 14, 1112-1115.
- Komar, A.A., Lesnik, T. und Reiss, C. (1999) Synonymous codon substitutions affect ribosome traffic and protein folding during *in vitro* translation. *FEBS Lett.* 462, 387-391.
- Komine, Y., Kitabatake, M., Yokogawa, T. und Nishikawa, K. (1994) A tRNA-like structure is present in 10Sa RNA, a small stable RNA from *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 91, 9223-9227.
- Komissarov, A.A., Marchbank, M.T. und Deutscher S.L. (1996) The use of Ni-nitrilotriacetic acid agarose for estimation of affinities of hexahistidine-tagged Fab to single stranded DNA. *Anal. Biochem.* 247, 123-129.
- Korman, A.J., Knudsen, P.J., Kaufman, J.F. und Strominger, J.L. (1982) cDNA clones for the heavy chain of HLA-DR antigen obtained after immunopurification of polysomes by monoclonal antibody. *Proc. Natl. Acad. Sci. USA* 79, 1844-1848.
- Kozak, M. (1984) Point mutations close to the AUG initiator codon affect the efficiency of translation of rat preproinsulin *in vivo*. *Nature* 308, 241-246.
- Kozak, M. (1989) Circumstances and mechanisms of inhibition of translation by secondary structure in eucaryotic mRNAs. *Mol. Cell. Biol.* 9, 5134-5142.
- Kraus, J.P. und Rosenberg, L.E. (1982) Purification of low abundance messenger RNAs from rat liver by polysome immunoadsorption. *Proc. Natl. Acad. Sci. USA* 79, 4015-4019.
- Kurz, M., Gu, K. und Lohse, P.A. (2000) Psoralen photo-crosslinked mRNA-puromycin conjugates: a novel template for the rapid and facile preparation of mRNA-protein fusions. *Nucleic Acids Res.* 28, e83.
- Kurzchalia, T.V., Wiedmann, M., Breter, H., Zimmermann, W., Bauschke, E. und Rapoport, T.A. (1988) tRNA-mediated labelling of proteins with biotin. A nonradioactive method for the detection of cell-free translation products. *Eur. J. Biochem.* 172, 663-668.
- LaBean, T.H. und Kauffman, S.A. (1993) Design of synthetic gene libraries encoding random sequence proteins with desired ensemble characteristics. *Protein Sci.* 2, 1249-1254.
- Laemmli, U.K. (1970) Cleavage of the structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.
- Lam, K.S., Salmon, S.E., Hersh, E.M., Hruby, V.J., Kazmierski, W.M. und Knapp, R.J. (1991) A new type of synthetic peptide library for identifying ligand-binding activity. *Nature* 354, 82-84.
- Lamla, T. und Erdmann, V.A. (2001) *In vitro* selection of other proteins than antibodies by means of ribosome display. *FEBS Lett.* 502, 35-40.
- Lamla, T., Mammeri, K. und Erdmann, V.A. (2001a) The cell-free protein biosynthesis-applications and analysis of the system. *Acta Biochim. Pol.* 48, 453-465.
- Lamla, T. und Erdmann, V.A. (2002) Improved batch translation system based on *E. coli* extract, in: *Cell-free translation systems*, pp. 23-39. Springer Verlag, ISBN 3-540-42050-9.
- Lancet, D., Sadovsky, E. und Seidemann, E. (1993) Probability model for molecular recognition in biological receptor repertoires: significance to the olfactory system. *Proc. Natl. Acad. Sci. USA* 90, 3715-3719.
- Lassen, D., Lucke, C., Kveder, M., Mesgarzadeh, A., Schmidt, J.M., Specht, B., Lezius, A., Spener, F. und Ruterjans, H. (1995) Three-dimensional structure of bovine heart fatty-acid-binding protein with bound palmitic acid, determined by multidimensional NMR spectroscopy. *Eur. J. Biochem.* 230, 266-280.
- Lesley, S.A., Brow, M.A. und Burgess, R.R. (1991) Use of *in vitro* protein synthesis from polymerase chain reaction-generated templates to study interaction of *E. coli* transcription factors with core RNA polymerase and for epitope mapping of monoclonal antibodies. *J. Biol. Chem.* 266, 2632-2638.
- Li, M. (2000) Applications of display technology in protein analysis. *Nat. Biotechnol.* 18, 1251-1256.
- Ling, L.L., Keohavong, P., Dias, C. und Thilly, W.G. (1991) Optimization of the polymerase chain reaction with regard to fidelity: modified T7, *Taq*, and *Vent* polymerases. *PCR Methods Appl.* 1, 63-69.

- Liu, R., Barrick, J.E., Szostak, J.W. und Roberts, R.W. (2000) Optimized synthesis of RNA-protein fusions for *in vitro* protein selection. *Methods Enzymol.* 318, 268-293.
- Low, N.M., Holliger, P.H. und Winter, G. (1996) Mimicking somatic hypermutation: affinity maturation of antibodies displayed on bacteriophage using a bacterial mutator strain. *J. Mol. Biol.* 260, 359-368.
- Madin, K., Sawasaki, T., Ogasawara, T. und Endo, Y. (2000) A highly efficient and robust cell-free protein synthesis system prepared from wheat embryos: plants apparently contain a suicide system directed at ribosomes. *Proc. Natl. Acad. Sci. USA* 97, 559-564.
- Makeyev, E.V., Kolb, V.A. und Spirin, A.S. (1996) Enzymatic activity of the ribosome-bound nascent polypeptide. *FEBS Lett.* 378, 166-170.
- Malkin, L.I. und Rich, A. (1967) Partial resistance of nascent polypeptide chains to proteolytic digestion due to ribosomal shielding. *J. Mol. Biol.* 26, 329.
- Mattheakis, L.C., Bhatt, R.R. und Dower, W.J. (1994) An *in vitro* polysome display system for identifying ligands from very large peptide libraries. *Proc. Natl. Acad. Sci. USA* 91, 9022-9026.
- Mattheakis, L.C., Dias, J.M. und Dower, W.J. (1996) Cell-free synthesis of peptide libraries displayed on polysomes. *Methods Enzymol.* 267, 195-207.
- McPhalen, C.A. und James, M.N. (1987) Crystal and molecular structure of the serine proteinase inhibitor CI-2 from barley seeds. *Biochemistry* 26, 261-269.
- Merk, H., W. Stiege, W., Tsumoto, K., Kumagai, I. und Erdmann, V.A. (1999) Cell-free expression of two single-chain monoclonal antibodies against lysozyme: Effect of domain arrangement on the expression. *J. Biochem.* 125:328-333.
- Merk, H. (2001) Steigerung der Effizienz des *Escherichia coli in vitro* Translationssystems durch Optimierung der Nukleinsäurekomponenten. *Dissertation*, FU Berlin.
- Monro, R.E. und Marcker, K.A. (1967) Ribosome-catalysed reaction of puromycin with a formylmethionine-containing oligonucleotide. *J. Mol. Biol.* 25, 347-350.
- Mullis, K.B. und Faloona, F.A. (1987) Specific synthesis of DNA *in vitro* via a polymerase-catalysed chain reaction. *Methods Enzymol.* 155, 335-350.
- Nemoto, N., Miyamoto-Sato, E., Husimi, Y. und Yanagawa, H. (1997) *In vitro* virus: bonding of mRNA bearing puromycin at the 3'-terminal end to the C-terminal end of its encoded protein on the ribosome *in vitro*. *FEBS Lett.* 414, 405-408.
- Nieba, L., Nieba-Axmann, S.E., Persson, A., Hämäläinen, M., Edebratt, F., Hansson, A., Lidholm, J., Magnusson, K., Karlsson, A.F. und Plückthun, A. (1997) BIACORE analysis of histidine-tagged proteins using a chelating NTA sensor chip. *Anal. Biochem.* 252, 217-228.
- Nirenberg, N.W. und Matthaei, J.H. (1961) The dependence of cell-free protein synthesis in *E. coli* on naturally occurring or synthetic polyribonucleotides. *Proc. Natl. Acad. Sci. USA* 47, 1588-1602.
- Noren, C.J., Anthony-Cahill, S.J., Griffith, M.C. und Schultz, P.G. (1989) A general method for site-specific incorporation of unnatural amino acids into proteins. *Science* 244, 182-188.
- Olins, P.O. und Rangwala, S.H. (1989) A novel sequence element derived from bacteriophage T7 mRNA acts as an enhancer of translation of the *lacZ* gene in *Escherichia coli*. *J. Biol. Chem.* 264, 16973-16976.
- Osborne, S.E., Matsumura, I. und Ellington, A.D. (1997) Aptamers as therapeutic and diagnostic reagents: problems and prospects. *Curr. Opin. Chem. Biol.* 1, 5-9.
- Patnaik, R. und Swartz, J.R. (1998) *E. coli*-based *in vitro* transcription/translation: *in vivo*-specific synthesis rates and high yields in a batch system. *Biotechniques* 24, 862-868.
- Pelletier, J.N., Campbell-Valois, F.X. und Michnick, S.W. (1998) Oligomerization domain-directed reassembly of active dihydrofolate reductase from rationally designed fragments. *Proc. Natl. Acad. Sci. USA* 95, 12141-12146.
- Plückthun, A., Schaffitzel, C., Hanes, J. und Jermutus, L. (2000) *In vitro* selection and evolution of proteins. *Adv. Protein Chem.* 55, 367-403.
- Ray, B.K. und Apirion, D. (1979) Characterization of 10S RNA: a new stable RNA molecule from *E. coli*. *Mol. Gen. Genet.* 174, 25-32.
- Roberts, R.W. und Szostak, J.W. (1997). RNA-peptide fusions for the *in vitro* selection of peptides and proteins. *Proc. Natl. Acad. Sci. USA* 94, 12297-12302.
- Roberts, R.W. (1999) Totally *in vitro* protein selection using mRNA-protein fusions and ribosome display. *Curr. Opin. Chem. Biol.* 3, 268-273.
- Roberts, R.W. und Ja, W.W. (1999) *In vitro* selection of nucleic acids and proteins: What are we learning? *Curr. Opin. Struct. Biol.* 9, 521-529.
- Roche, E.D. und Sauer, R.T. (1999) SsrA-mediated peptide tagging caused by rare codons and tRNA scarcity. *EMBO J.* 18, 4579-4589.
- Saiki, R.K., Scharf, S., Faloona, F.A., Mullis, K.B., Horn, G.T., Erlich, H.A., Norman, A. (1985) Enzymatic amplification of β -globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230, 1350-1354.
- Saiki, R.K., Gelfand, S., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, S.J. und Ehrlich, H.A. (1988) Primer directed amplification of DNA with thermostable DNA polymerase. *Science* 239, 487-491.

- Sambrook, J., Fritsch, E.F. und Maniatis, T. (1989) Molecular cloning: A laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S. und Coulson, A.R. (1977) DNA Sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
- Schaffitzel, C., Berger, I., Postberg, J., Hanes, J., Lipps, H.J. und Plückthun, A. (2001) *In vitro* generated antibodies specific for telomeric guanine-quadruplex DNA react with *Stylonychia lemnae* macronuclei. *Proc. Natl. Acad. Sci. USA* 98, 8572-8577.
- Schmidt, T.G.M. und Skerra, A. (1993) The random peptide library-assisted engineering of a C-terminal affinity peptide, useful for the detection and purification of a functional Ig Fv fragment. *Protein Eng.* 6, 109-122.
- Schmidt, T.G.M. und Skerra, A. (1994) One-step affinity purification of bacterially produced proteins by means of the „Strep tag“ and immobilized recombinant core streptavidin. *J. Chromtog. Sect. A*, 676, 337-345.
- Schmidt, T.G.M., Koepke, J., Frank, R. und Skerra, A. (1996) Molecular interaction between the Strep-tag affinity peptide and its cognate target, Streptavidin. *J. Mol. Biol.* 255, 753-766.
- Shapiro, D.J. (1981) Quantitative ethanol precipitation of nanogram quantities of DNA and RNA. *Anal. Biochem.* 10, 229-231.
- Sharp, P.A., Sugden, B. und Sambrook, J. (1973) Detection of two restriction endonuclease activities in *Haemophilus parainfluenzae* using analytical agarose-ethidium bromide electrophoresis. *Biochemistry* 12, 3055-3063.
- Shaw, G.L., Davis, B., Keeler, J. und Fersht, A.R. (1995) Backbone dynamics of chymotrypsin inhibitor 2: effect of breaking the active site bond and its implications for the mechanism of inhibition of serine proteases. *Biochemistry* 34, 2225-2233.
- Shine, J. und Dalgarno, L. (1975) Growth-dependent changes in terminal heterogeneity involving 3'-adenylate of bacterial 16S ribosomal RNA. *Nature* 256, 232-233.
- Smith, G.P. (1985) Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science* 228, 1315-1317.
- Smith, W.P., Tai, P.C. und Davis, B.D. (1978) Interaction of secreted nascent chains with surrounding membrane in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* 75, 5922.
- Stemmer, W.P.C. (1994) Rapid evolution of a protein *in vitro* by DNA shuffling. *Nature* 370, 389-391.
- Stiege, W. und Erdmann, V.A. (1995) The potentials of the *in vitro* protein biosynthesis system. *J. of Biotechnology* 41, 81-90.
- Studier, F.W., Rosenberg, A.H., Dunn, J.J. und Dubendorff, J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.* 185, 60-89.
- Tabor, S. und Richardson, C.C. (1987) DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc. Natl. Acad. Sci. USA* 84, 4767-4771.
- Takahashi, F., Ebihara, T., Mie, M., Yanagida, Y., Endo, Y., Kobatake, E. und Aizawa, M. (2002) Ribosome display for selection of active dihydrofolate reductase mutants using immobilized methotrexate on agarose beads. *FEBS Lett.* 514, 106-110.
- Tate, W.P. und Brown, C.M. (1992) Translation termination: „stop“ for protein synthesis or „pause“ for regulation of gene expression. *Biochemistry* 31, 2443-2450.
- Tawfik, D.S. und Griffiths, A.D. (1998) Man-made cell-like compartments for molecular evolution. *Nat. Biotechnol.* 16, 652-656.
- Tuerk, C. und Gold, L. (1990) Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 249, 505-510.
- Tuite, M.F. und Stansfield, I. (1994) Termination of protein synthesis. *Mol. Biol. Rep.* 19, 171-181.
- Vogelstein, B. und Gillespie, D. (1979) Preparative and analytical purification of DNA from agarose. *Proc. Natl. Acad. Sci. USA* 76, 615-619.
- Voss, S. und Skerra, A. (1997) Mutagenesis of a flexible loop in streptavidin leads to higher affinity for the Strep-tag II peptide and improved performance in recombinant protein purification. *Protein Eng.* 10, 975-982.
- Weber, K. und Osborn, M. (1969) The reliability of molecular weight determination by dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Biol. Chem.* 244, 4406-4412.
- Weber, P.C., Ohlendorf, D.H., Wendoloski, J.J. und Salemme, F.R. (1989) Structural origins of high-affinity biotin binding to streptavidin. *Science*, 243, 85.
- Weiss, B., Jacquemin-Sablon, A., Live, T.R., Fareed, G.C. und Richardson, C.C. (1968) Enzymatic breakage and joining of deoxyribonucleic acid. I. Further purification and properties of polynucleotide ligase from *Escherichia coli* infected with bacteriophage T4. *J. Biol. Chem.* 243, 4543-4555.
- Wilson, D.S., Keefe, A.D. und Szostak, J.W. (2001) The use of mRNA display to select high-affinity protein-binding peptides. *Proc. Natl. Acad. Sci. USA* 98, 3750-3755.
- Winter, G., Griffith, A.D., Hawkins, R.E. und Hoogenboom, H.R. (1994) Making antibodies by phage display technology. *Ann. Rev. Immunol.* 12, 433-455.
- Yanisch-Perron, C., Vieira, J. und Messing, J. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33, 103-119.

-
- Yu, A., Barreiro, V. und Haggard-Ljungquist, E. (1994) Regulation of int gene expression in bacteriophage P2. *J. Virol.* 68, 4220-4226.
- Yuan, R. (1981) Structure and mechanism of multifunctional restriction endonucleases. *Annu. Rev. Biochem.* 50, 285-319.
- Zhao, H., Giver, L., Shao, Z., Affholter, J.A. und Arnold, F.H. (1998) Molecular evolution by staggered extension process (StEP) *in vitro* recombination. *Nat. Biotechnol.* 16, 258-261.
- Zhouravleva, G., Frolova, L., Le Goff, X., Le Guellec, R., Inge-Vechtomov, S., Kisselev, L. und Philippe, M. (1995) Termination of translation in eukaryotes is governed by two interacting polypeptide chain release factors, eRF1 and eRF3. *EMBO* 14, 4065-4072.
- Zubay, G. (1973) *In vitro* synthesis of protein in microbial systems *Annu. Rev. Genet.* 7, 267-287.