

**Singular Probability Distributions and Fractal Properties  
of Sets of Real Numbers Defined by the Asymptotic  
Frequencies of their  $S$ -Adic Digits**

**Sergio Albeverio, Mykola Pratsiovytyi, Grygoriy Torbin**

**no. 231**

Diese Arbeit ist mit Unterstützung des von der Deutschen Forschungsgemeinschaft getragenen Sonderforschungsbereiches 611 an der Universität Bonn entstanden und als Manuskript vervielfältigt worden.

Bonn, Juni 2005

Dedicated to V.S.Korolyuk on occasion of his 80-th birthday

**SINGULAR PROBABILITY DISTRIBUTIONS  
AND FRACTAL PROPERTIES OF SETS OF REAL NUMBERS  
DEFINED BY THE ASYMPTOTIC FREQUENCIES OF THEIR S-ADIC DIGITS**

SERGIO ALBEVERIO<sup>1,2,3,4</sup>, MYKOLA PRATSIOVYTYI<sup>5</sup>, GRYGORIY TORBIN<sup>6,7</sup>

ABSTRACT. Properties of the set  $T_s$  of "particularly non-normal numbers" of the unit interval are studied in details ( $T_s$  consists of real numbers  $x$ , some of whose  $s$ -adic digits have the asymptotic frequencies in the nonterminating  $s$ -adic expansion of  $x$ , and some do not). It is proven that the set  $T_s$  is residual in the topological sense (i.e., it is of the first Baire category) and it is generic in the sense of fractal geometry ( $T_s$  is a superfractal set, i.e., its Hausdorff-Besicovitch dimension is equal to 1). A topological and fractal classification of sets of real numbers via analysis of asymptotic frequencies of digits in their  $s$ -adic expansions is presented.

<sup>1</sup> Institut für Angewandte Mathematik, Universität Bonn, Wegelerstr. 6, D-53115 Bonn (Germany) <sup>2</sup> SFB 611, Bonn, BiBoS, Bielefeld - Bonn <sup>3</sup> CERFIM, Locarno and Acc. Arch., USI (Switzerland) <sup>4</sup> IZKS, Bonn; e-mail: albeverio@uni-bonn.de

<sup>5</sup> National Pedagogical University, Pyrogova str. 9, 01030 Kyiv (Ukraine); e-mail: prats@ukrpost.net

<sup>6</sup> National Pedagogical University, Pyrogova str. 9, 01030 Kyiv (Ukraine), <sup>7</sup> Institute for Mathematics of NASU, Tereshchenkivs'ka str., 3, 01601 Kyiv (Ukraine); e-mail: torbin@imath.kiev.ua

**AMS Subject Classifications (2000): 11K55, 28A80, 60G30.**

**Key words:** normal numbers, non-normal numbers, essentially non-normal numbers, particularly non-normal numbers, Hausdorff-Besicovitch dimension, fractals, Baire category.

## 1. INTRODUCTION

Let us consider the classical  $s$ -adic expansion of  $x \in [0, 1]$  :

$$x = \sum_{n=1}^{\infty} s^{-n} \alpha_n(x) = \Delta^s \alpha_1(x) \alpha_2(x) \dots \alpha_k(x) \dots, \alpha_k(x) \in A = \{0, 1, \dots, (s-1)\},$$

and let  $N_i(x, k)$  be the number of digits "i" among the first  $k$  digits of the  $s$ -adic expansion of  $x$ ,  $i \in A$ . If the limit  $v_i(x) = \lim_{k \rightarrow \infty} \frac{N_i(x, k)}{k}$  exists, then the number  $v_i(x)$  is said to be the frequency of the digit "i" (or the asymptotic frequency of "i") in the  $s$ -adic expansion of  $x$ .

A property of an element  $x \in M$  is usually said to be "normal" if "almost all" elements of  $M$  have this property. There exist many mathematical notions (e.g., cardinality, measure, Hausdorff-Besicovitch dimension, Baire category) allowing us to interpret the words "almost all" in a rigorous mathematical sense. "Normal" properties of real numbers are deeply connected with the asymptotic frequencies of their digits in some systems of representation.

The set

$$N_s = \{x | (\forall i \in A) \lim_{k \rightarrow \infty} \frac{N_i(x, k)}{k} = \frac{1}{s}\}$$

is said to be the set of  $s$ -normal numbers (or the set of real numbers which are normal with respect to the base  $s$ ). It is well known (E.Borel, 1909), that the sets  $N_s$  and the set  $N^* = \bigcap_{s=2}^{\infty} N_s$  are of full Lebesgue measure (i.e., they have Lebesgue measure 1).

The unit interval  $[0, 1]$  can be decomposed in the following way:

$$[0, 1] = E_s \cup D_s,$$

where

$$E_s = \{x | v_i(x) \text{ exists}, \forall i \in A\},$$

$$D_s = \{x | \exists i \in A, \lim_{k \rightarrow \infty} \frac{N_i(x, k)}{k} \text{ does not exist}\}.$$

The set  $D_s$  is said to be the set of non-normal real numbers. Each of the subsets  $E_s$  and  $D_s$  can be decomposed in the following natural way.

The set

$$W_s = \{x | (\forall i \in A) : v_i(x) \text{ exists, and } (\exists j \in A) : v_j(x) \neq \frac{1}{s}\}$$

is said to be the set of *quasinormal* numbers. It is evident that

$$E_s = W_s \cup N_s, W_s \cap N_s = \emptyset.$$

The set

$$L_s = \{x | (\forall i \in A) \lim_{k \rightarrow \infty} \frac{N_i(x, k)}{k} \text{ does not exist}\}$$

is said to be the set of *essentially non-normal* numbers.

The set

$$T_s = \{x | (\exists i \in A) : \lim_{k \rightarrow \infty} \frac{N_i(x, k)}{k} \text{ does not exist, and } (\exists j \in A) : \lim_{k \rightarrow \infty} \frac{N_j(x, k)}{k} \text{ exists}\}$$

is said to be the set of *particularly non-normal* numbers.

It is evident that

$$D_s = L_s \cup T_s, L_s \cap T_s = \emptyset.$$

The sets  $N_s, W_s, T_s, L_s$  are everywhere dense sets, because the frequencies  $v_i(x)$  do not depend on any finite number of  $s$ -adic symbols of  $x$ . It is also not hard to prove that these sets have the cardinality of the continuum.

The main purpose of the paper is to fill in completely the following table, which reflects the metric, topological and fractal properties of the corresponding sets:

	Lebesgue measure	Hausdorff dimension	Baire category
$N_s$			
$W_s$			
$L_s$			
$T_s$			

Let  $\mathbf{v} = (v_0, v_1, \dots, v_{s-1})$  be a stochastic vector and let

$$W_s[\mathbf{v}] = \{x : x = \Delta^s \alpha_1(x) \alpha_2(x) \dots \alpha_k(x) \dots, \lim_{k \rightarrow \infty} \frac{N_i^*(x, k)}{k} = v_i, \forall i \in A\}.$$

The well known Besicovitch-Eggleston's theorem (see, e.g., [4, 6]) gives the following formulae for the determination of the Hausdorff-Besicovitch dimension  $\alpha_0(W_s[\mathbf{v}])$  of the set  $W_s[\mathbf{v}]$ :

$$\alpha_0(W_s[\mathbf{v}]) = \frac{\sum_{i=0}^{s-1} v_i \log v_i}{-\log s}.$$

From the latter formulae it easily follows that the set  $W_s$  of all quasinormal numbers is a superfractal set, i.e.,  $W_s$  is a set of zero Lebesgue measure with full Hausdorff-Besicovitch dimension ( $\alpha_0(W_s) = 1$ ).

Properties of subsets of the set of non-normal numbers have been intensively studied during recent years (see, e.g., [12, 8, 9, 10] and references therein). Some interesting subsets of  $D_s$  were studied in [8] by using the techniques and results from the theory of multifractal divergence points. In [12] it has been proven that the set  $D_s$  is superfractal.

In the paper [1] of the authors it has been proven that the set  $L_s$  of essentially non-normal numbers is also superfractal and it is of the second Baire category. Moreover, it has been proven that the set  $L_s$  contains an everywhere dense  $G_\delta$ -set. So, the sets  $N_s, W_s, T_s$  are of the first Baire category. From these results it follows that essentially non-normal numbers are generic in the topological sense as well as in the sense of fractal geometry; nevertheless, the set  $L_s$  is small from the point of view of Lebesgue measure.

The main goal of the present paper is the investigation of fractal properties of the set  $T_s$  of particularly non-normal numbers. To this end we apply a probabilistic approach for the calculation of the Hausdorff dimension of subsets. More precisely, we apply the results of fine fractal analysis of singular continuous probability distributions.

The first step of the fractal analysis of a singular continuous measure  $\nu$  is the investigation of metric, topological and fractal properties of the corresponding topological support  $S_\nu$  (i.e. the minimal closed set supporting the measure). These are good characteristics only for the class of uniform Cantor-type singular measures. But, in general, they are only "external characteristics", because there exist essentially different singular continuous measures concentrating on the common topological support. The main idea of the paper [1] consisted in the construction of singular continuous measures whose topological supports coincide with some subsets of the set of essentially non-normal numbers.

The second step of the fractal analysis of a singular continuous measure  $\nu$  is the determination of the Hausdorff dimension  $\alpha_0(\nu)$  (and the local Hausdorff dimension) of the measure, i.e., roughly speaking, finding the Hausdorff dimension of the minimal (in the fractal dimension sense) supports (which are not necessarily closed) of the measure. This problem is much more complicated than the previous one (see, e.g., [3]), especially in the case of essentially superfractal measures.

In Section 2 we prove that for all  $s \geq 3$  the set  $T_s$  is of full Hausdorff dimension. To prove the main result we construct a sequence of singular continuous measures  $\mu_p$  such that the corresponding minimal dimensional supports consist of only particularly non-normal numbers, and apply the results of [3] to perform a fine fractal analysis of these supports.

## 2. FRACTAL PROPERTIES OF THE SET OF PARTICULARLY NON-NORMAL NUMBERS

Let us study the sets  $T_s$  of particularly non-normal numbers which were defined in Section 1. It is easy to see that the set  $T_2$  is empty, because from the existence of the asymptotic frequency  $v_i(x)$  for some  $i \in \{0, 1\}$  the existence of another asymptotic frequency follows.

**Theorem 1.** *For any positive integer  $s \geq 3$  the set  $T_s$  of particularly non-normal real numbers is superfractal, i.e., the Hausdorff-Besicovitch dimension of the set  $T_s$  equals 1.*

*Proof.* To prove the theorem we shall construct a superfractal set  $G \subset T_s$ .

In the sequel we usually shall not use the indices  $s$  in the notation of the corresponding subsets, since  $s$  will be an arbitrary fixed natural number greater than 2. Let us consider the classical  $s$ -adic expansion of  $x \in [0, 1] : x = \sum_{n=1}^{\infty} s^{-n} \alpha_n(x) = \Delta^s \alpha_1(x) \alpha_2(x) \dots \alpha_k(x) \dots$ . If  $x$  is an  $s$ -adic rational number, then we shall use the representation without the period "s-1".

For a given  $p \in \mathbb{N}$  and for any  $x \in [0, 1)$  we define the following mapping  $\varphi_p$ :

$$\begin{aligned} \varphi_p(x) &= \varphi_p(\Delta^s \alpha_1(x) \alpha_2(x) \dots \alpha_k(x) \dots) = \\ &= \Delta^s \overbrace{00 \dots 0}^{s-1} \overbrace{11 \dots 1}^{s-1} \dots \overbrace{(s-2)(s-2) \dots (s-2)}^{s-1} (s-1) \alpha_1(x) \alpha_2(x) \dots \alpha_{s^2 p}(x) \\ &\quad \overbrace{00 \dots 0}^{2(s-1)} \overbrace{11 \dots 1}^{2(s-1)} \dots \overbrace{(s-2)(s-2) \dots (s-2)}^{2(s-1)} (s-1)(s-1) \alpha_{s^2 p+1}(x) \alpha_{s^2 p+2}(x) \dots \alpha_{s^2 p+2s^2 p}(x) \dots \\ &\quad \overbrace{00 \dots 0}^{2^{k-1}(s-1)} \overbrace{11 \dots 1}^{2^{k-1}(s-1)} \dots \overbrace{(s-2)(s-2) \dots (s-2)}^{2^{k-1}(s-1)} \overbrace{(s-1)(s-1) \dots (s-1)}^{2^{k-1}} \alpha_{(2^{k-1}-1)s^2 p+1}(x) \dots \alpha_{(2^{k-1}-1)s^2 p}(x) \dots \end{aligned}$$

Let us explain the construction of  $\varphi_p$ . First of all we divide the  $s$ -adic expansion of  $x$  into groups in the following way: the  $k$ -th group consists of the sequence  $(\alpha_{(2^{k-1}-1)s^2 p+1}(x) \dots \alpha_{(2^k-1)s^2 p}(x))$ ,  $k \in \mathbb{N}$ . The  $s$ -adic expansion of  $y = \varphi_p(x)$  is constructed from the  $s$ -adic expansion of  $x$  via inserting (before the  $k$ -th group) the following series of fixed symbols  $(0 \dots 01 \dots 1 \dots (s-2) \dots (s-2) (s-1) \dots (s-1))$ , where each symbol "i" ( $0 \leq i \leq s-2$ ) occurs  $2^{k-1}(s-1)$  times, but the symbol "s-1" occurs  $2^{k-1}$  times.

Let  $M_p = \varphi_p([0, 1)) = \{y : y = \varphi_p(x), x \in [0, 1)\}$ .

For a given  $p \in N$  and for any  $y \in M_p$  we define the mapping  $\psi_p(y)$  in the following way: if  $y = \varphi_p(x) =$

$$\begin{aligned} &= \Delta^s \overbrace{00\dots 0}^{s-1} \dots \overbrace{(s-2)\dots(s-2)}^{s-1} (s-1)\alpha_1(x)\alpha_2(x)\dots\alpha_{s^{2p}}(x) \\ &\quad \overbrace{0\dots 0}^{2(s-1)} \dots \overbrace{(s-2)\dots(s-2)}^{2(s-1)} (s-1)(s-1)\alpha_{s^{2p+1}}(x)\alpha_{s^{2p+2}}(x)\dots\alpha_{s^{2p+2s^{2p}}}(x) \dots, \end{aligned}$$

then  $z = \psi_p(y) =$

$$\begin{aligned} &= \Delta^s \overbrace{0\dots 0}^{s-1} (s-1) \dots \overbrace{(s-2)\dots(s-2)}^{s-1} (s-1) (s-1) (01\dots(s-2)) \alpha_1(x)\alpha_2(x)\dots\alpha_{s^{2p}}(x) \\ &\quad \overbrace{00\dots 0}^{(s-1)} \overbrace{(s-1)\dots(s-1)}^{(s-1)} \dots \overbrace{(s-2)(s-2)\dots(s-2)}^{(s-1)} (s-1) \overbrace{(s-2)(s-2)\dots(s-2)}^{(s-1)} (s-1) \end{aligned}$$

$$(s-1) (01\dots(s-2)) (s-1) (01\dots(s-2)) \alpha_{s^{2p+1}}(x)\alpha_{s^{2p+2}}(x)\dots\alpha_{s^{2p+2s^{2p}}}(x) \dots, x \in [0, 1),$$

i.e., the s-adic expansion of  $z = \psi_p(y)$  can be obtained from the s-adic expansion of  $y = \varphi(x)$  by using the following algorithm:

- 1) after any fixed symbol "(s-1)" we insert the following series of symbols: (01...(s-2));
- 2) after any subseries consisting of (s-1) fixed symbols "i",  $0 \leq i \leq s-2$  we insert the symbol "s-1".

Let  $f_p = \psi_p(\varphi_p)$  and let

$$S_p = f_p([0, 1)) = \{z : z = f_p(x), x \in [0, 1)\} = \{z : z = \psi_p(y), y \in M_p\},$$

$$G_p = f_p([0, 1)) = \{z : z = f_p(x), x \in N_s\}.$$

The following two lemmas will describe some properties of the constructed sets  $G_p$ .

**Lemma 1.** For any  $z = \sum_{n=1}^{\infty} s^{-n} \alpha_n(z) \in G_p$  the limit  $\lim_{n \rightarrow \infty} \frac{N_i(z, n)}{n}$  does not exist for any  $i \in \{0, 1, \dots, s-2\}$ , and  $\lim_{n \rightarrow \infty} \frac{N_{s-1}(z, n)}{n} = \frac{1}{s}$ .

*Proof.* The set  $G_p$  has the following structure:

$$G_p = \{z : z = \Delta^s \underbrace{\overbrace{0\dots 0}^{s-1} (s-1) \dots \overbrace{(s-2)\dots(s-2)}^{s-1} (s-1) (s-1) (01\dots(s-2)) \alpha_1(x)\alpha_2(x)\dots\alpha_{s^{2p}}(x)}_{\text{first group}}\}$$

$$\underbrace{0\dots 0(s-1) 0\dots 0(s-1)\dots(s-2)\dots(s-2)(s-1) (s-2)\dots(s-2)(s-1) (s-1)(01\dots(s-2)) (s-1)(01\dots(s-2))\alpha_{s^2 p+1}(x)\alpha_{s^2 p+2}(x)\dots\alpha_{s^2 p+2s^2 p}(x)}_{\text{second group}} \dots, x \in N_s \}.$$

From  $x \in N_s$  it follows that the symbol "s-1" has the asymptotic frequency  $\frac{1}{s}$  in the sequence  $\{\alpha_k(x)\}$  and the equality  $\lim_{n \rightarrow \infty} \frac{N_{s-1}(z,n)}{n} = \frac{1}{s}$  follows from the construction of the set  $G_p$ .

Let  $l_k$  be the number of the position at which the above k-th group of symbols ended, i.e.,  $l_k = s^2(p+1)(2^k - 1)$ .

Let  $m'_k(i)$  be the number of the position at which the k-th series of the fixed symbols "i" and "(s-1)" ( $0 \leq i \leq s-2$ ) ended, i.e.,  $m'_{k+1}(i) = s^2(p+1)(2^k - 1) + s(i+1)2^k$ .

Let  $m''_k(i)$  be the number of the position at which the k-th series of the fixed symbols "i" ( $0 \leq i \leq s-2$ ) started, i.e.,  $m''_{k+1}(i) = s^2(p+1)(2^k - 1) + si2^k + 1$ .

If  $z \in G_p$ , then there are  $s(2^{k+1} - 1) + d_k$  symbols "i" ( $0 \leq i \leq s-2$ ) among the first  $m'_{k+1}(i)$  symbols of the s-adic expansion of  $z$ , where  $d_k$  is the quantity of the symbol "i" among the first  $(2^k - 1)s^2 p$  s-adic symbols  $\alpha_i(x)$  in the expansion of  $x = f_p^{-1}(z)$ . Since  $x$  is an s-normal number, we have:  $d_k = (2^k - 1)sp + o(2^k)$ .

$$\text{So, } \lim_{n \rightarrow \infty} \frac{N_i(z, m'_{k+1}(i))}{m'_{k+1}(i)} = \lim_{n \rightarrow \infty} \frac{(2^{k+1}-1)s+(2^k-1)sp+s^{-1}o(2^k)}{s^2(p+1)(2^k-1)+s(i+1)2^k} = \frac{p+2}{s(p+1)+i+1}.$$

If  $z \in G_p$ , then there are  $s(2^k - 1) + d_k$  symbols "i" ( $0 \leq i \leq s-2$ ) among the first  $m''_{k+1}(i) - 1$  symbols of the s-adic expansion of  $z$ .

$$\text{So, } \lim_{n \rightarrow \infty} \frac{N_i(z, m''_{k+1}(i)-1)}{m''_{k+1}(i)-1} = \lim_{n \rightarrow \infty} \frac{(2^k-1)s+(2^k-1)sp+s^{-1}o(2^k)}{s^2(p+1)(2^k-1)+si2^k} = \frac{p+1}{s(p+1)+i} < \frac{p+2}{s(p+1)+i+1}.$$

Therefore, for any  $z \in G_p$  and for any  $i \in \{0, 1, \dots, s-2\}$  the limit  $\lim_{n \rightarrow \infty} \frac{N_i(z,n)}{n}$  does not exist.  $\square$

The following Corollary is immediate, using the definitions of  $G_p, T_s$  and Lemma 1:

**Corollary.**  $G_p \subset T_s, \forall p \in N$ .

**Lemma 2.** *The Hausdorff-Besicovitch dimension of the set  $G_p$  is equal to  $\frac{p}{p+2}$ .*

*Proof.* Let  $B_p(i)$  be the subset of  $N$  with the following property:  $\forall k \in N, k \in B_p(i)$  if and only if  $\alpha_k(f_p(x)) = i$  for any  $x \in [0, 1)$ , i.e.,  $B_p(i)$  consists of the numbers of positions with the fixed symbols "i" in the s-adic expansion of any  $z \in S_p$ . Let  $B_p = \bigcup_{i=0}^{s-1} B_p(i)$ , and let  $C_p = N \setminus B_p$ .

Let us consider the following random variable  $\xi^{(p)}$  with independent  $s$ -adic digits:

$$\xi^{(p)} = \sum_{k=1}^{\infty} s^{-k} \xi_k^{(p)},$$

where  $\xi_k^{(p)}$  are independent random variables with the following distributions: if  $k \in B_p(i)$ , then  $\xi_k^{(p)}$  takes the value "i" with probability 1. If  $k \in C_p$ , then  $\xi_k^{(p)}$  takes the values  $0, 1, \dots, (s-1)$  with probabilities  $\frac{1}{s}, \frac{1}{s}, \dots, \frac{1}{s}$ .

It is evident that the set  $S_p$  is the topological support of the distribution of the random variable  $\xi^{(p)}$ . Actually, the corresponding probability measure  $\mu_p = P_{\xi^{(p)}}$  is the image of Lebesgue measure on  $[0, 1)$  under the mapping  $f_p = \psi_p(\varphi_p)$ , i.e.,  $\forall E \subset \mathcal{B} : \mu_p(E) = \mu_p(E \cap S_p) = \lambda(f_p^{-1}(E \cap S_p))$ .

**A)** Firstly we prove that  $\alpha_0(G_p) \leq \frac{p}{p+2}$ . Since  $G_p \subset S_p$ , it is sufficient to show that  $\alpha_0(S_p) \leq \frac{p}{p+2}$ . To this end we consider the sequence  $\{B_i^{(k)}\} (k \in N, i \in \{1, 2, \dots, s^{s^2 p(2^{k-1}-1)}\})$  of special coverings of the set  $S_p$  by  $s$ -adic closed intervals of the rank  $m_k = l_k - 2^{k-1}s^2 p = s^2(p+1)(2^k - 1) - 2^{k-1}s^2 p$ . For any  $k \in N$  the covering  $\{B_i^{(k)}\}$  consists of the  $s^{s^2 p(2^{k-1}-1)}$  closed  $s$ -adic intervals of  $m_k$ -th rank with length  $\varepsilon_k = s^{-(s^2(p+1)(2^k-1)-2^{k-1}s^2 p)}$ .

The  $\alpha$ - volume of the covering  $\{B_i^{(k)}\}$  is equal to

$$I_{\varepsilon_k}^{\alpha}(S_p) = s^{s^2 p(2^{k-1}-1)} \cdot s^{-\alpha(s^2(p+1)(2^k-1)-2^{k-1}s^2 p)} = s^{(p-\alpha(p+2))2^{k-1}s^2} \cdot s^{\alpha(p+1)-p}.$$

For the Hausdorff premeasure  $h_{\varepsilon_k}^{\alpha}$  we have:  $h_{\varepsilon_k}^{\alpha}(S_p) \leq I_{\varepsilon_k}^{\alpha}(S_p)$  for any  $k \in N$ . So, for the Hausdorff measure  $H_{\alpha}$  we have  $H_{\alpha}(S_p) \leq \lim_{k \rightarrow \infty} I_{\varepsilon_k}^{\alpha}(S_p) = 0$  if  $\alpha > \frac{p}{p+2}$ .

Hence,  $\alpha_0(S_p) \leq \frac{p}{p+2}$ .

**B)** Secondly we prove that  $\alpha_0(G_p) \geq \frac{p}{p+2}$ . To this end we shall analyze the internal fractal properties of the singular continuous measure  $\mu_p$ .

For any probability measure  $\nu$  one can introduce the notion of the Hausdorff dimension of the measure in the following way:

$$\alpha_0(\nu) = \inf_{E \in N(\nu)} \{\alpha_0(E), E \in \mathcal{B}\},$$

where  $N(\nu)$  is the class of all "possible supports" of the measure  $\nu$ , i.e.,

$$N(\nu) = \{E : E \in \mathcal{B}, \nu(E) = 1\}.$$

An explicit formula for the determination of the Hausdorff dimension of the measures with independent  $Q^*$ -symbols has been found in [3]. Applying this formula to our case ( $q_{ik} = \frac{1}{s}, \forall k \in N, \forall i \in \{0, 1, \dots, s-1\}$ ), we have

$$\alpha_0(\mu_p) = \lim_{n \rightarrow \infty} \frac{H_n}{n \ln s}$$

where  $H_n = \sum_{j=1}^n h_j$ , and  $h_j$  are the entropies of the random variables  $\xi_j^{(p)} : h_j = - \sum_{i=0}^{s-1} p_{ij} \ln p_{ij}$ .

If  $j \in B_p$ , then  $h_j = 0$ . If  $j \in C_p$ , then  $h_j = \ln s$ .

So,

$$\alpha_0(\mu_p) = \lim_{n \rightarrow \infty} \frac{H_n}{n \ln s} = \lim_{k \rightarrow \infty} \frac{H_{m_k}}{m_k \ln s} = \lim_{k \rightarrow \infty} \frac{s^2 p (2^{k-1} - 1) \ln s}{(s^2 (p+1) (2^k - 1) - p s^2 2^{k-1}) \ln s} = \frac{p}{p+2}.$$

The above defined set  $G_p = f_p(N_s)$  is a support of the measure  $\mu_p$ , because  $\mu_p = \lambda(f_p^{-1})$  and the Lebesgue measure of the set  $N_s$  of  $s$ -normal numbers of the unit interval is equal to 1.

Since  $G_p \in N(\mu_p)$  and  $\alpha_0(\mu_p) = \frac{p}{p+2}$ , we get  $\alpha_0(\mu_p) \geq \frac{p}{p+2}$ , which proves Lemma 2.  $\square$

**Corollary.** *The set  $G_p$  is the minimal dimensional support of the measure  $\mu_p$ , i.e.,  $\alpha_0(G_p) \leq \alpha_0(E)$  for any other support  $E$  of the measure  $\mu_p$ .*

Finally, let us consider the set  $G = \bigcup_{p=1}^{\infty} G_p$ . From Lemma 1 it follows that  $G \subset T_s$ . From Lemma 2 and from the countable stability of the Hausdorff dimension it follows that  $\alpha_0(G) = \sup_p \alpha_0(G_p) = 1$ . So,  $\alpha_0(T_s) = 1$ , which proves Theorem 1.  $\square$

Summarizing the results of Sections 1 and 2, we have for  $s > 2$ :

	Lebesgue measure	Hausdorff dimension	Baire category
$N_s$	1	1	first
$W_s$	0	1	first
$T_s$	0	1	first
$L_s$	0	1	second

For the case  $s = 2$  we have a corresponding result, but the Hausdorff dimension of the set  $T_s$  is equal to 0, because the set  $T_s$  is empty for  $s = 2$ .

## Acknowledgment

This work was partly supported by DFG 436 UKR 113/78, DFG 436 UKR 113/80, INTAS 00-257 and SFB-611 projects. The last two named authors gratefully acknowledge the hospitality of the Institute of Applied Mathematics and of the IZKS of the University of Bonn.

## REFERENCES

- [1] Albeverio S., Pratsiovytyi M., Torbin G., Topological and fractal properties of subsets of real numbers which are not normal, to appear in Bulletin des Sciences Mathématiques ( SFB 611 Preprint Nr.208 (2004), Universität Bonn).
- [2] Albeverio S., Pratsiovytyi M., Torbin G., Fractal probability distributions and transformations preserving the Hausdorff-Besicovitch dimension, Ergodic Theory and Dynamical Systems, **24** (2004 ), 1-16.
- [3] Albeverio S. , Torbin G., Fractal properties of singular continuous probability distributions with independent  $Q^*$ -digits, Bulletin des Sciences Mathématiques, 129(2005) No.4, 356-367.
- [4] Besicovitch A., On the sum of digits of real numbers represented in the dyadic systems, Math. Ann., 110 (1934), 321-330.
- [5] Billingsley P., *Ergodic theory and information*, John Wiley & Sons, Inc., New York-London-Sydney, 1965.
- [6] Eggleston H.G., The fractional dimension of a set defined by decimal properties, Quart.J.Math. Oxford Ser. 20 (1949), 31-36.
- [7] Falconer K.J., *Fractal geometry*, John Wiley & Sons, 1995.
- [8] Olsen L., Applications of multifractal divergence points to some sets of  $d$ -tuples of numbers defined by their  $N$ -adic expansion. Bull.Sci.Math., 128 (2004), 265-289.
- [9] Olsen L., Applications of multifractal divergence points to sets of numbers defined by their  $N$ -adic expansion. Math. Proc. Cambridge Philos. Soc. 136 (2004), No. 1, 139–165.
- [10] Olsen L., Winter S., Normal and non-normal points of self-similar sets and divergence points of self-similar measures. J. London Math. Soc. (2) 67 (2003), No. 1, 103–122.
- [11] Pratsiovytyi M., *Fractal approach to investigations of singular distributions*, National Pedagogical Univ., Kyiv, 1998 (In Ukrainian).
- [12] Pratsiovytyi M., Torbin G., Superfractality of the set of numbers having no frequency of  $n$ -adic digits, and fractal probability distributions. Ukrainian Math. J. 47(1995),No. 7, 971-975.
- [13] Schweiger F., *Ergodic theory of fibred systems and metric number theory*, Oxford: Clarendon Press, 1995.
- [14] Turbin A.F., Pratsiovytyi M .V., *Fractal sets, functions, and distributions*, Naukova Dumka, Kiev, 1992 (In Russian).

Bestellungen nimmt entgegen:

Institut für Angewandte Mathematik  
der Universität Bonn  
Sonderforschungsbereich 611  
Wegelerstr. 6  
D - 53115 Bonn

Telefon: 0228/73 3411

Telefax: 0228/73 7864

E-mail: anke@iam.uni-bonn.de

Homepage: <http://www.iam.uni-bonn.de/sfb611/>

### Verzeichnis der erschienenen Preprints ab No. 200

200. Albeverio, Sergio; Ayupov, Shavkat A.; Omirov, Bakhrom A.: Cartan Subalgebras and Criterion of Solvability of Finite Dimensional Leibniz Algebras; eingereicht bei: Journal of Lie Theory
201. not published
202. Sturm, Karl-Theodor: Convex Functionals of Probability Measures and Nonlinear Diffusions on Manifolds; erscheint in: J. Math. Pures Appl.
203. Sturm, Karl-Theodor: On the Geometry of Metric Measure Spaces
204. Müller, Jörn; Müller, Werner: Regularized Determinants of Laplace Type Operators, Analytic Surgery and Relative Determinants
205. Mandrekar, Vidyadhar; Rüdiger, Barbara: Lévy Noises and Stochastic Integrals on Banach Spaces
206. Albeverio, Sergio; Proskurin, Daniil; Turowska, Lyudmila: On  $*$ -Representations of the  $\lambda$ -Deformation of a Wick Analogue of the CAR Algebra
207. Albeverio, Sergio; Fei, Shao-Ming; Song, Tong-Qiang: Multipartite Entangled State Representation and its Squeezing Transformation
208. Albeverio, Sergio; Pratsiovytyi, Mykola; Torbin, Grygoriy: Topological and Fractal Properties of Real Numbers which are not Normal; eingereicht bei: Bull. Sci. Math.
209. Albeverio, Sergio; Cattaneo, Laura; Fei, Shao-Ming; Wang, Xiao-Hong: Multipartite States Under Local Unitary Transformations
210. Albeverio, Sergio; Bozhok, Roman; Dudkin, Mykola; Koshmanenko, Volodymyr: Dense Subspaces in Scales of Hilbert Spaces
211. Otto, Felix; Westdickenberg, Michael: Eulerian Calculus for the Contraction in the Wasserstein Distance; eingereicht bei: SIAM J. Math. Anal.
212. Albeverio, Sergio; Smorodina, Nataliya V.: The Multiple Stochastic Integrals and the Transformations of the Poisson Measure
213. Albeverio, Sergio; Pustyl'nikov, Lev D.: Some Properties of Dirichlet  $L$ -Functions Associated with their Nontrivial Zeros I.

214. Ebmeyer, Carsten: Regularity in Sobolev Spaces for the Fast Diffusion and the Porous Medium Equation; erscheint in: J. Math. Anal. Appl.
215. Albeverio, Sergio; Korolyuk, Volodymyr S.; Bratiychuk, Mykola S.: Asymptotic Behaviour of the Ruin Probability for Stochastic Risk Models
216. Albeverio, Sergio; Korolyuk, Volodymyr S.; Lebedev, E.A.; Chechelnitsky, A.A.: Functional Limit Theorems for Multi-Channel Networks
217. Albeverio, Sergio; Binding, Paul; Hryniv, Rostyslav; Mykytyuk, Yaroslav: Inverse Spectral Problems for Coupled Oscillating Systems
218. Kornhuber, Ralf; Krause, Rolf: Robust Multigrid Methods for Vector-Valued Allen-Cahn Equations with Logarithmic Free Energy
219. Albeverio, Sergio; Konstantinov, Alexei; Koshmanenko, Volodymyr: Remarks on the Inverse Spectral Theory for Singularly Perturbed Operators
220. Otto, Felix; Rump, Tobias; Slepčev, Dejan: Coarsening Rates for a Droplet Model: Rigorous Upper Bounds
221. Gozzi, Fausto; Marinelli, Carlo: Stochastic Optimal Control of Delay Equations Arising in Advertising Models
222. Griebel, Michael; Oeltz, Daniel; Vassilevsky, Panayot: Space-Time Approximation with Sparse Grids
223. Arndt, Marcel; Griebel, Michael; Novák, Václav; Šittner, Petr; Roubíček, Tomáš: Martensitic Transformation in NiMnGa Single Crystals: Numerical Simulation and Experiments
224. DeSimone, Antonio; Knüpfer, Hans; Otto, Felix: 2-d Stability of the Néel Wall
225. Griebel, Michael; Metsch, Bram; Oeltz, Daniel; Schweitzer, Marc Alexander: Coarse Grid Classification: A Parallel Coarsening Scheme for Algebraic Multigrid Methods
226. De Santis, Emilio; Marinelli, Carlo: Stochastic Games with Infinitely many Interacting Agents
227. Cantero-Álvarez, Rubén; Otto, Felix: The Concertina Pattern: A Supercritical Bifurcation in Ferromagnetic Thin Films
228. Verleye, Bart; Klitz, Margrit; Croce, Roberto; Roose, Dirk; Lomov, Stepan; Verpoest, Ignaas: Computation of Permeability of Textile Reinforcements; erscheint in: Proceedings, Scientific Computation IMACS 2005, Paris (France), July 11-15, 2005
229. Albeverio, Sergio; Pustyl'nikov, Lev; Lokot, Tatiana; Pustylnikov, Roman: New Theoretical and Numerical Results Associated with Dirichlet  $L$ -Functions
230. Garcke, Jochen; Griebel, Michael: Semi-Supervised Learning with Sparse Grids; erscheint in: Proceedings of 22<sup>nd</sup> Int. Conference on Machine Learning
231. Albeverio, Sergio; Pratsiovytyi, Mykola; Torbin, Grygoriy: Singular Probability Distributions and Fractal Properties of Sets of Real Numbers Defined by the Asymptotic Frequencies of their  $S$ -Adic Digits